



Protein glycation inhibitory activities of *Lawsonia inermis* and its active principles

Nighat Sultana, Muhammed Iqbal Choudhary & Ambrin Khan

To cite this article: Nighat Sultana, Muhammed Iqbal Choudhary & Ambrin Khan (2009) Protein glycation inhibitory activities of *Lawsonia inermis* and its active principles, Journal of Enzyme Inhibition and Medicinal Chemistry, 24:1, 257-261, DOI: [10.1080/14756360802057500](https://doi.org/10.1080/14756360802057500)

To link to this article: <https://doi.org/10.1080/14756360802057500>



Published online: 01 Feb 2009.



Submit your article to this journal [↗](#)



Article views: 1787



View related articles [↗](#)



Citing articles: 4 View citing articles [↗](#)

Protein glycation inhibitory activities of *Lawsonia inermis* and its active principles

NIGHAT SULTANA¹, MUHAMMED IQBAL CHOUDHARY², & AMBRIN KHAN²

¹PCSIR Laboratories Complex, Pharmaceutical Research Center, Karachi 75280, Pakistan, and ²International Center for Chemical Sciences, H. E. J. Research Institute of Chemistry, University of Karachi, Karachi 75270, Pakistan

(Received 19 January 2008; accepted 22 February 2008)

Abstract

The protein glycation inhibitory activity of ethanolic extract of *Lawsonia inermis* (henna) plant tissues was evaluated *in vitro* using the model system of bovine serum albumin and glucose. Protein oxidation and glycation are posttranslational modifications that are implicated in the pathological development of many age-related disease processes. This study investigated the effects of *Lawsonia inermis* ethanolic extract and its components, on protein damage induced by a free radical generator in *in vitro* assay system. We found that alcoholic extract of *Lawsonia inermis* can effectively protect against protein damage and showed that its action is mainly due to Lawsone. In addition, the presence of gallic acid also plays an important role in the protective activity against protein oxidation and glycation. Two known compounds, namely, Lawsone and gallic acid previously isolated from this plant were subjected to glycation bioassay for the first time. It was found that the alcoholic extract, lawsone (1) and gallic acid (2) showed significant inhibition of Advanced Glycated End Products (AGEs) formation and exhibit 77.95%, 79.10% and 66.98% inhibition at a concentration of 1500 µg/mL, 1000 µg/mL and 1000 µM respectively. *Lawsonia inermis*, compounds 1 and 2 were found to be glycation inhibitors with IC₅₀ 82.06 ± 0.13 µg/mL, 67.42 ± 1.46 µM and 401.7 ± 6.23 µM respectively. This is the first report on the glycation activity of these compounds and alcoholic extract of *Lawsonia inermis*.

Keywords: *Lawsonia inermis*, lawsone, gallic acid, antioxidant, glycation, inhibitors

Introduction

The accumulation of the reaction products of protein glycation (nonenzymatic reaction of proteins with glucose and other reducing sugars) in living organisms leads to structural and functional modifications of tissue proteins. Many studies have shown a significant role for glycation in the progress of normal ageing and the pathogenesis of age-related diseases, such as diabetes, atherosclerosis, end-stage renal disease, rheumatoid arthritis, and neurodegenerative diseases [1]. The present study was undertaken to find out the glycation inhibitory activity of the alcoholic suspension of *Lawsonia Inermis* and its constituents.

Lawsonia inermis is a shrub of Asian origin, indigenous to Egypt, Arabia and India [2]. It is cultivated in many tropical countries and warm

temperate region as a hedge plant. In the West and the Middle East, the leaves are utilized locally for cosmetic purposes [3]. It highlights hair especially as tint for hands, feet and nails [4–6] and is also employed as a deodorant. It is an excellent conditioning agent and used as an ingredient in shampoos, hair dyes, conditioners and rinses. The dye is also used to stain leather and hides in various dye industries for commercial use.

Medicinally, *L. inermis* is used as an astringent, antihemorrhagic, intestinal antineoplastic, cardio-inhibitor, hypotensive, sedative, anti-inflammatory and antioxidant [7,8]. It has also been used as a folk remedy against amoebiasis, soothe fevers, headache, jaundice, hysteria, nervous disorder and leprosy [4,9–12]. Henna extracts show antibacterial, antifungal and ultraviolet light screening activity. It has exhibited

Correspondence: N. Sultana, PCSIR Laboratories Complex, Pharmaceutical Research Center, Karachi 75280, Pakistan. Tel: 92 21 8141832 35. Fax: 92 21 8141847. E-mail: nighat2001us@yahoo.com

antifertility activity in animals and may induce menstruation. In Malaysia, the leaf decoction is used after child birth and for beri-beri, rheumatism, skin disorders, stomach disorders and venereal disease [10]. Some of the tribals are using this plant leaves in a lower concentration for body pain, skin infections to reduce the lesions after bee sting, allergy, infections, inflammations [13]. *L. inermis* leaves are used as a remedy in skin diseases in the form of paste or decoction against boils burns, bruises and skin inflammation. Leaves in the form of paste are used as external application in head ache and rubbed over the soles of the burning feet. Decoction of the leaves is used as gargle in sore throat [14]. Crude extract of *L. inermis* (Linn) leaves shows dose dependent analgesic, antipyretic effect in rats [15]. Leaves are also useful to bring down the severity of many medical problems like dysentery, diseases of the spleen, lumbago, bronchitis and syphilitic eye infection [16,17]. *Lawsonia* shows anticancer against sarcoma 180 in mice and walker 256 carcinosarcoma.

A number of components such as lawsone [18], xanthenes [19,20], isoplumbagin, phenolic glycosides, coumarins, gallic acid [21], xanthenes, quinoids, organo-chlorine compounds, β -sitosterol glucosides, flavonoids including luteolin and its 7-O-glycosides, fats, resinm, henna-tanin. and triterpenoids [22,23] have been isolated from various parts of the plant. The leaf was found to contain thirty-six components, which constituted 80.4% of the oil, were identified. The major components were ethyl hexadecanoate (24.4%), (E)-methyl cinnamate (11.4%), isocaryophyllene (8.1%), (E)- β -ionone (5.8%) and methyl linolenate (4.1%). The flower oil was found to contain among other secondary metabolites, (Z)-2-hexenol, linalool and β -ionone [24]. The commercial essential oil comprised of mainly α -terpineol, terpinolene, δ -3-carene and γ -terpineol [25]. *Lawsonia* leaves contains a coloring compound 2-OH-1, 4 naphthaquinone (Lawson) **1** in higher concentration [26] which has proved to have analgesic, genotoxicity [27], anti inflammatory and antipyretic effects in rat models [16] and also some inhibitory effect against *Proteus* and *Staphylococcus aureus* [28]. Gallic acid slightly inhibits *Streptococcus aureus*. It contributes to significant inhibition of colon, esophageal, liver, lung, tongue and skin cancers. Based on these observations, compounds **1**, **2** and alcoholic extract of the aerial parts of *Lawsonia inermis* were screened to determine the possible antiglycation activities.

Materials and methods

General experimental procedures

The mass spectra were recorded on a Jeol HX-110 instrument. The ^1H and ^{13}C NMR spectra were recorded in CDCl_3 at 500, 400 and 125, 75 MHz,

respectively, on a Bruker AM-500, 400 NMR spectrometer. The UV and IR spectra were recorded on Shimadzu UV-240 and JASCO A-320 spectrophotometers, respectively. Optical rotations were measured on a polatronic D Polarimeter. The purity of the compounds was checked on TLC (Si-gel, Merck PF₂₅₄, 0.25 mm thickness). Melting points were determined in glass capillary tubes using a Buchi 535 and a Gallenkamp 30/MF-370 melting point apparatus. Bovine serum albumin (BSA) was purchased from Research Organics Cleveland, while others chemicals {glucose anhydrous, trichloroacetic acid sodium azide, dimethyl sulfoxide, sodium dihydrogen phosphate, sodium chloride, disodium hydrogen phosphate, potassium chloride, potassium dihydrogen phosphate, and sodium hydroxide} were purchased from Sigma Aldrich. Sodium phosphate buffer (pH 7.4), was prepared by mixing Na_2HPO_4 and NaH_2PO_4 (67 mM) containing sodium azide (3 mM), phosphate buffer saline (PBS) pH 10 was prepared by mixing NaCl (137 mM) + Na_2HPO_4 (8.1 mM) + KCl (2.68 mM) + KH_2PO_4 (1.47 mM). pH 10 was adjusted with NaOH (0.25 mM), while BSA (10 mg/mL) and anhydrous glucoses (50 mg/mL) solutions were prepared in sodium phosphate buffer.

Plant material

L. inermis leaves were collected in summer (April and May) locally and identified by a taxonomist, Mr. Abid Askari, PSO at the Botany Section of PCSIR Laboratories Complex, Karachi. After the species identification was done, Coimbatore were allowed to dry in open air in the shaded area for few weeks. Air dried *L. inermis* leaves were powdered mechanically. At first the dried leaf powder was tested for the presence of contamination before and after autoclaving.

Extraction and isolation

Air-dried powdered aerial parts of *L. inermis* (20 kg dry weight) were extracted with EtOH (100 L). The EtOH extract was concentrated to a gum (822 gm), dissolved in distd. water and extracted thoroughly with pet. ether (45 L). The pet. ether soluble portion was evaporated under reduced pressure to yield a gum (66.92 gm).

The remaining aqueous layer was acidified with acetic acid to pH 3, and then extracted with CHCl_3 . The remaining aqueous acidic layer was made alkaline with NH_4OH to pH 12 and extracted with CHCl_3 (40 L). The CHCl_3 soluble portion was dried as a crude mixture (74.96 gm), which was chromatographed on a si-gel column (Merck, 70–230 mesh, 2015.01 gm). Elution of this column with 96% CHCl_3 -MeOH (15 L) yielded an impure mixture (11.02 gm, Fr.20–26, each fraction 500 ml each) containing compounds **1** and **2**. This mixture was chromatographed on a SiO_2 gel

column (2.5 cm × 70 cm, Merck, 70–230 mesh, 322.11 gm) which was first eluted with CHCl₃, and then with 5:95 MeOH: CHCl₃. Fractions from 55–90 (500 ml each), 3.96 gm obtained with 5:95 MeOH: CHCl₃ (3 L) were again subjected to CC over silica gel (70–230 mesh size, 99.24 gm). The column (1.5 cm × 50 cm) was initially eluted with CHCl₃-MeOH (96:4, 9 L) to afford eighteen fractions. These were combined and further purified by repeated TLC plates (Merck, PF 254, 0.5 mm) using CHCl₃: MeOH (92:8) to afford **1** (18.13 mg, $9.0 \times 10^{-5}\%$ yield with $R_f = 0.32$). Fractions 90–96 obtained on elution with 80:20, 3.5 L, CHCl₃-MeOH contain **2**, which was purified by preparative tlc plates using 85:15 CHCl₃-MeOH (20 mg, $1.0 \times 10^{-4}\%$ yield with $R_f = 0.47$).

Lawsonone (1): Yellow crystals, M. P. 191–193°C, EIMS: m/z 174.15, UV/vis (MeOH), IR (CHCl₃), ¹H NMR (500 MHz, CDCl₃): δ ¹³C NMR (125 MHz, CD₃OD) δ : [29–33].

Gallic acid (2): White powder, EIMS: m/z 170.12, UV/vis (MeOH), IR (CHCl₃), ¹H NMR (500 MHz, CDCl₃): δ : ¹³C NMR (125 MHz, CD₃OD) δ [29–33].

In vitro glycation inhibition assay

Sample preparation. Samples were prepared in DMSO for plant extracts (1 mg /mL), for pure compounds (2 mM).

Non-enzymatic glycation of protein. According to the method of Vinson and Howard, bovine serum albumin (BSA) (10 mg/ml) in 40 μ L phosphate buffer, the

reaction mixture was mixed with an aqueous solution of test sample. Six concentrations were prepared for *Lawsonia inermis* alcoholic extract, its components lawsone and gallic acid. After incubating at 37°C for two weeks, the fluorescent reaction products were assayed on a fluorescence spectrophotometer with an excitation wavelength of 370 nm and an emission wavelength of 340 nm. The data were expressed in terms of percentage inhibition, calculated from a control measurement of the fluorescence intensity of the reaction mixture with no test sample.

Methodology. In 96-well plate assays, each well contain 60 μ L reaction mixtures (20 μ L BSA (10 mg/mL + 20 μ L of glucose anhydrous (50 mg/mL) + 20 μ L test sample) [34]. Glycated control contain 20 μ L BSA + 20 μ L glucose + 20 μ L sodium phosphate buffer (pH 7.4, 67 mM), while blank control contains 20 μ L BSA and 40 μ L sodium phosphate buffer. Reaction mixture was incubated at 37°C for 7-days [35]. After incubation, 6 μ L (100%) of TCA was added in each well and centrifuged (15,000 rpm) for 4 minutes at 4°C [36]. After centrifugation, the pellets were rewashed with 60 μ L (10%) of TCA [36]. The supernatant containing glucose, inhibitor and interfering substance was removed and pellet contains AGE-BSA were dissolved in 60 μ L PBS [28]. Assessment of fluorescence spectrum (ex. 370 nm), and change in fluorescence intensity (ex. 370 nm to em. 440 nm) based on AGEs were monitored by using a spectrofluorimeter (RF-1500, Shimadzu, Japan) [1]. %

Table I. *In vitro* quantitative inhibition of glycation by compounds 1, 2 and an alcoholic extract of *Lawsonia inermis*.

COMPOUND NAME	FLURECENCE (FU)	% INHIBITION	CONCENTRATION	
			(μ M/ml)	IC ₅₀ VALUE
<i>Lawsonia inermis</i> alcoholic extract	195 FU	77.95	1500 μ g	82.06 \pm 0.13
	226 FU	74.28	1000 μ g	
	288 FU	67.32	500 μ g	
	296 FU	66.62	250 μ g	
	328 FU	62.62	100 μ g	
	611 FU	36.67	31.25 μ g	
Gallic acid (2)	338 FU	66.98	1000 μ M	401.7 \pm 6.23
	383 FU	58.83	500 μ M	
	535 FU	36.51	250 μ M	
	625 FU	34.88	100 μ M	
	629 FU	30.62	62.5 μ M	
	681 FU	26.87	31.25 μ M	
Lawsonone (1)	176 FU	79.1	1000 μ M	67.42 \pm 1.46
	228 FU	72.9	500 μ M	
	213 FU	74.7	250 μ M	
	236 FU	72	100 μ M	
	530 FU	37.25	50 μ M	
STANDARD (1 mM/mL)	594 FU	29.6	31.25 μ M	41.9 \pm 2.3 μ M
RUTIN	127 FU	85.80%	100 μ M	

Inhibition was calculated through the following formula.

$$\% \text{ Inhibition} = [1 - (\text{Fluorescence of sample} / \text{Fluorescence of glycated})] \times 100$$

Results and discussion

In this study we tested *Lawsonia inermis* plant tissue and its two constituents Lawsone (1) and gallic acid (2) for their inhibitory activity on protein glycation. Lawsone (1) was isolated as a yellow crystalline compound from the alcoholic extract of leaves of *Lawsonia inermis* by column and thin layer chromatography. Compound 1 ($\text{C}_{10}\text{H}_6\text{O}_3$) was identified by comparison of its data with those reported earlier [20]. ^1H -NMR and ^{13}C -NMR of compound 1 were the same as reported previously [29–33]. Compound 2 ($\text{C}_7\text{H}_6\text{O}_5$) was isolated as white powder from alcoholic extract of the same plant. It was identified by comparison of its data with those reported earlier, which was originally isolated from *Onosma hispidum*. The structures were established by spectroscopic studies [29–33,37–46]. Table I shows the effect of *Lawsonia inermis* extract and its components on Advance Glycated ends Product (AGEs) formation after incubation for 2 weeks. *L. inermis* alcoholic extract inhibited the reaction 77.95%, 74.28%, 67.32%, 66.62%, 62.62% and 36.67%, significantly at concentrations of 1500, 1000, 500, 250, 100 and 31.25 $\mu\text{g}/\text{mL}$ respectively and sustained dose dependency. Its IC_{50} value was $82.06 \pm 0.13 \mu\text{g}/\text{mL}$ (Figure 1).

Compound 1 inhibited the reaction 79.1%, 72.9%, 74.7%, 72.0%, 37.25% and 29.6%, (Figure 3) significantly at concentrations of 1000, 500, 250, 100, 50 and 31.25 $\mu\text{M}/\text{ml}$ respectively and sustained dose dependency. Its IC_{50} value was $67.42 \pm 1.46 \mu\text{M}$. Compound 2 inhibited the reaction 66.98%, 58.83%, 36.51%, 34.88%, 30.62% and 26.87%, significantly at concentrations of 1000, 500, 250, 100, 62.5 and 31.25 $\mu\text{M}/\text{ml}$ respectively (Figure 2) and sustained dose dependency. Its IC_{50} value was $401.7 \pm 6.23 \mu\text{M}$.

The most active compound was lawsone. Its IC_{50} value was $67.42 \pm 1.46 \mu\text{M}$. and approached that




Figure 1. Glycation inhibition activity of *Lawsonia inermis*.




Figure 2. Glycation inhibition activity of gallic acid.




Figure 3. Glycation inhibition activity of 2-hydroxy-1,4-naphthoquinone.

of *Rutin* at $41.9 \pm 2.3 \mu\text{M}$. *L. inermis* (leaves) was the second most bioactive extract followed by gallic acid. The IC_{50} values of these were above 67 μM .

Lawsonia inermis, compound 1 and 2 exhibited *in vitro* glycation activity with IC_{50} $82.06 \pm 0.13 \mu\text{M}$, $67.42 \pm 1.46 \mu\text{M}$ and $401.7 \pm 6.23 \mu\text{M}$ respectively. *Rutin* (IC_{50} $41.9 \pm 2.3 \mu\text{M}$) was used as a positive control [1]. The antioxidant activities of *Lawsonia inermis* and compounds 1 and 2 were earlier reported through pharmacological methods. The glycation inhibitory activity was significantly correlated with the antioxidative potency of the extracts. There is growing interest in natural products with combined anti-glycation and antioxidant properties as they may have reduced toxicity. The positive glycation inhibitory and antioxidant activity of this plant might suggest a possible role in targeting ageing and diabetic complications.

Acknowledgements

The authors wish to thank the Ministry of Science and Technology, Government of Pakistan for providing the financial support for the current study.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

- [1] Hye YK, Kyong K. Protein glycation and antioxidative activity of some plant extracts *in vitro*. *J Agric Food Chem* 2003;51: 1586–1591.
- [2] Amelio D, Botanicals FS. A phytocosmetic desk reference. Boca Raton: CRC Press; 1999. p 126–135.
- [3] Ong HC, Norzalina J. Malay Herbal Medicine in Gemencheh, Negri, Sembilan, Malaysia. *Fitoterapia* 1997;70:10–14.
- [4] Keay RWJ. Trees in Nigeria. Oxford: Clarendon Press; 1989. p 50.
- [5] Mabberley DJ. The plant book. London: Cambridge University Press; 1990. p 318.
- [6] El-Kamali HH, Khalid SA. The Most Common Herbal Remedies in Central Sudan. *Fitoterapia* 1996;68:301–366.
- [7] Ali BH, Bashir AK, Tanira MO. Anti-inflammatory, antipyretic, and analgesic effects of *Lawsonia inermis* L. (henna) in rats. *Pharmacology* 1995;51(6):356–363.
- [8] Mikhaeil BR, Badria FA, Maatooq GT, Amer MM. Antioxidant and immunomodulatory constituents of henna leaves. *Z Naturforsch [C]* 2004;59(7–8):468–476.
- [9] Bep-Oliver-Bever. Medicinal plants in tropical west Africa. London: Cambridge University Press; 1986. p 53.
- [10] Dalziel JM. The useful plants of west tropical Africa. London: Crown Agents; 1937. p 40–41.
- [11] Burkill HM, The Useful Plants of West Tropical Africa. Families J–L, Royal Botanic Gardens 3 1995. p 562–564.
- [12] KI-Merzabani A, El-Hazer AA, Attia MA, Al Duweini AK, Ghazal AM. Screening system for Egyptian plants with potential anti-tumor activity. *Planta Med* 1976;36:150–155.
- [13] Zargari A. Medicinal plants., Vol.2 Tehran University Publication; 1991. p 353–358.
- [14] Dhur UN. Chopra's indigenous drugs of India, 2nd ed. Calcutta: Dhur UN and sons Pvt. Ltd; New Delhi, 1958. p 48–55.
- [15] Evans WC. Trease and evans pharmacognosy. 4th ed. London: WB Saunders Co. Ltd. 1996. p 248.
- [16] Warriar PK, Nambiar VP, Ramankutty C, editors. Indian medicinal plants, compendium of 500 species. Chennai: Orient Longmann Pvt. Ltd. 1995. p 303–306.
- [17] Duke JA. Handbook of medicinal herbs. Haemolytic activity and nephrotoxicity of 2-hydroxy-1,4-naphthoquinone in rats. Boca Raton: CRC Press; 1985. p 274.
- [18] Bhardway DK, Murari R, Seshadri TR, Singh R. Lacoumarin from *Lawsonia inermis*. *Phytochemistry* 1976;15:1789–1792.
- [19] Bhardway DK, Seshadri TR, Singh R. Xanthoness from *Lawsonia inermis*. *Phytochemistry* 1977;16:1616–1617.
- [20] Bhardway DK, Jain RK, Jain BC, Mehta CK. 1-Hydroxy-3,7-dimethoxy-6-acetoxanthone, A new xanthone from *Lawsonia inermis*. *Phytochemistry* 1978;17:1440–1441.
- [21] Stanley RH, Gilbert AF. Hair dyes and hair- dyeing chemistry and technique. London: Great Britain Whitefriars press Ltd; 1939. p 72–77.
- [22] Sarita G, Mohammed A, Sarwar AM, Masatake N, Tatsuko S. Isolation and characterization of a dihydroxysterol from *Lawsonia inermis*. *Nat Prod Lett* 1994;4:195–201.
- [23] Handa G, Kapil A, Sharma S, Singh J. Lawnermis acid: A new anticomplementary triterpenoid from *Lawsonia inermis* seeds. *Indian J Chem* 1977;VB(36B):252–256.
- [24] Wong KC, Teng YE. Volatile components of *Lawsonia inermis* L Flowers. *J Essent oil Res* 1995;7:425–428.
- [25] Reichling VJ, Harkenthal M, Brandt H, Bayerl C. Temporare HennaTaottoos. *Deut Apoth Ztg* 1999;33:35–41.
- [26] Bhuvaneswari K, Gnana Poongothai S, Kuruvilla A, Appala Raju B. Inhibitory concentrations of lawsonia innermis dry powder for urinary pathogens, *Indian Journal of Pharmacology* 2002; 34: 260–263.
- [27] Kirkland D, Marzin D. An assessment of the genotoxicity of 2-hydroxy-1, 4-naphthoquinone, the natural dye ingredient of Henn. *Mutat Res* 2003;537(2):183–199.
- [28] Pandey DK, Cahandra H, Sexena AR, Dixit SN. *Indian Drugs* 1982;19:459.
- [29] Yan Y, Guo HW, Jian WP, Yang KE, Tong SL, Fang CG, Li Q, Chang X. Catalytic Conversion of 2-naphthol to 2-hydroxy-1,4-naphthoquinone under mild conditions, *Chinese Journal of Chemistry* 2004; 22: 487–491.
- [30] Morris R, Nagarkatti J, Heaton J, Lane C. Aldrich catalog handbook of fine chemical 1996–1997. Aldrich Chemical Co., Inc. 1996. p 829.
- [31] Zhao YX, Sun XY. Spectrum analysis and structure identifying of organic compounds (Chinese). ed. I Chinese Science and Technology University Press; June 1992. p 315–325.
- [32] Pouchert CJ. The Aldrich library of infrared spectra. ed. III Aldrich Chemical Company, Inc; 1981. p 897A.
- [33] Silverstein RM, Bassler GC, Morrill TC. Spectrometric identification of organic compounds. ed. 3 John Wiley; 1977. p 65.
- [34] Takako N, Takako Y, Katsutoshi T, Seui S, Lekh RJ. *J Agric Food Chem* 2002;50:2418–2422.
- [35] Fumio Y, Toshiaki A, Yoshihiro Y, Hiroyuki N. Antioxidative and anti- glycation activity of Garcinol from *Garcinia indica* fruit rind. *J Agric Food Chem* 2000;48:180–185.
- [36] Nobuyasu M, Tadashi A, Chihiro S, Hiroyuki K, Mitsuharu O, Junichi H, Makoto U. Screening system for the Maillard reaction inhibitors from natural product extracts. *J Health Sci* 2002;48(6):520–526.
- [37] Atta-ur-Rahman, Sultana N, Jahan S, Choudhary MI. Studies on the constituents of *Commiphora mukul*. *Z Naturforsch* 2005; 60b:1202–1206.
- [38] Atta-ur-Rahman, Sultana N, Jahan S, Choudhary MI. Phytochemical studies on *Skimmia laureola*. *Nat Prod Lett* 1998;12(3):223.
- [39] Fatima K, Sultana N. Studies on bioassay directed antifungal activity of medicinal plants of *Skimmia laureola*, *Calotropis procera* and *P. pterocarpum*. *J Chem Soc Pak* 2003;25:328.
- [40] Atta-ur-Rahman, Sultana N, Choudhary MI, Shah PM, Khan MR. Isolation and structural studies on the chemical constituents of *Skimmia laureola*. *J Nat Prod* 1998;61(6):713.
- [41] Atta-ur-Rahman. Nuclear magnetic resonance spectroscopy, basic principles. New York: Springer-Verlag; 1986. p 227.
- [42] Atta-ur-Rahman. One-and two-dimentional NMR spectroscopy. Amsterdam: Elsevier Science Publishers; 1989. p 406.
- [43] Sultana N, Atta-ur-Rahman, Khalid A. New natural cholinesterase inhibiting and calcium channel blocking quinoline alkaloids. *J Enz Inhib Med Chem* 2006;21(6):703.
- [44] Sultana N, Atta-ur-Rahman, Khalid A. A new fatty ester and a new triterpene from *Skimmia laureola* *Nat Prod Res* 2008., (in press)
- [45] Sultana N, Atta-ur-Rahman, Khan TH. Tyrosinase inhibitor fatty ester and a quinoline alkaloid from *Skimmia laureola*. *Z Naturforsch B* 2005;60:1186.
- [46] Atta-ur-Rahman, Sultana N. Phytochemical studies on *Adhatoda vasica*. *Nat Prod Lett* 1997;10:249.