

Acta Oncologica



ISSN: 0284-186X (Print) 1651-226X (Online) Journal homepage: informahealthcare.com/journals/ionc20

Linking measured intercellular oxygen concentration to human cell functions

Peter Ebbesen, Kai-Uwe Eckardt, Fedor Ciampor & Erik O. Pettersen

To cite this article: Peter Ebbesen, Kai-Uwe Eckardt, Fedor Ciampor & Erik O. Pettersen (2004) Linking measured intercellular oxygen concentration to human cell functions, Acta Oncologica, 43:6, 598-600, DOI: 10.1080/02841860410020220

To link to this article: https://doi.org/10.1080/02841860410020220



Published online: 08 Jul 2009.



🖉 Submit your article to this journal 🗹





View related articles



Citing articles: 4 View citing articles 🕑

Linking Measured Intercellular Oxygen Concentration to Human Cell Functions

Peter Ebbesen, Kai-Uwe Eckardt, Fedor Ciampor and Erik O. Pettersen

From Aalborg University, Laboratory for Stem Cell Research, Aarhus, Denmark (P. Ebbesen), University of Erlangen-Nuremberg, Nephrology/Hypertension Section (K.-U. Eckardt), Presidium of Slovak Academy of Sciences, Bratislava, Slovakia (F. Ciampor), University of Oslo, Department of Physics, Oslo, Norway (E.O. Pettersen)

Correspondence to: Professor Peter Ebbesen, the Laboratory for Stem Cell Research, Aalborg University, Gustav Wieds Vej 10B, 8000 Aarhus C, Denmark. E-mail: ebbesen@lst.aauc.dk

Acta Oncologica Vol. 43, No. 6, pp. 598-600, 2004

Received 19 March 2004 Accepted 9 June 2004

To the Editor

The cell responses to a specific pericellular oxygen concentration depend on a number of factors including cell type, duration of the particular oxygen concentration, cell cycle stage, pH and concentration of nutrients and regulatory molecules.

The isolated effect of low oxygen, however, seems decisive enough to allow some general statements about cell regulatory responses to various pericellular oxygen concentrations.

Quantification of oxygen tension around the cells is therefore an important indicator of the cell phenotype (1).

The developing understanding is that responses to hypoxia are orchestrated by a number of overlapping regulatory processes. These processes result in a sequence of changes in the cell phenotype according to changes in the oxygen concentration.

Reliable oxygen measurements in both in vitro model systems (2-5) and in vivo (6-9) are indicative of which phenotype the cell will exhibit within a given range of oxygen concentrations. The present short survey deals with these relationships.

When the intercellular oxygen concentration in tissues is reduced from normoxia $(4-1\% O_2 \approx 4-1 \text{ kPa} \text{ and corresponding to } 40\,000-10\,000 \text{ ppm})$ to *moderate hypoxia* $(1-0.1\% O_2 \text{ and kPa, correspond$ $ing to 10\,000-1000 \text{ ppm})$, respiration, and as a result the ATP production, is largely unaffected (2, 10). For a long time, this observation has stood in the way of realizing that this is not the case for many other cell functions. In fact, there is a striking effect on gene expression due to transcription, mRNA stability, protein synthesis, and post-translational modifications.

In the range of 0.1-1% O₂/kPa O₂ (1000-10000 ppm O₂), the central hypoxia-responsive pathways of the hypoxia-inducible factor (HIF) are functionally very important. These pathways function by binding to hypoxia-responsive elements on promoters of more than 60, but not all, hypoxia-regulated genes (11). The activity of this HIF transcription regulator is itself governed by prolyl hydroxylases. In the presence of a sufficiently high concentration of molecular oxygen, these prolyl hydroxylases convert specific prolyl residues in the HIF-1 α subunits into hydroxyproline. This allows capture by the von Hippel-Lindau tumour suppressor E₃ ligase complex, leading to ubiquitination and destruction, thus preventing HIF activity at higher oxygen concentrations. Below 1% O₂/1 kPa (10000 ppm O₂), oxygenation is too small to

induce degradation of the HIF-1 α subunit, and HIF-1 accumulates instantaneously and influences specific promoter regions (12, 13). A second mechanism regulating HIF-1 activity is mediated by an asparaginyl hydroxylase, which modulates transactivations by HIF-1.

Downstream of HIF, carbonic anhydrase IX (CAIX) is an important example of a factor with a number of other functions. It is furthermore responsive to an upward shifted range of hypoxia $(0.1-5\% \text{ O}_2/\text{kPa} \text{ or } 1\,000-50\,000 \text{ ppm})$ and has a longer half-life.

Moderate hypoxia first results in a general translation inhibition, which later switches to a more specific, cap-dependent translation inhibition. Certain genes, including HIF-1 and VEGF, contain an internal ribosome entry site (IRES) allowing ribosomes to bind directly to the mRNA, thus bypassing the cap-dependent transcription repression (14).

In vitro stress proteins, e.g. HSP70 and HSP90, may become expressed, but may also appear as a response to reoxygenation (15) and function as a nuclear transfer factor by HIF-1.

The functional goal of these different cell activities can be seen as a restoration of the oxygen supply by HIF-1 mediated upregulation of the growth factor (VEGF), the platelet-derived endothelial cell growth factor vasodilating NO and erythropoietin production (16).

From 0.1% O_2 (1000 ppm) down to 0.001% O_2 (10 ppm) is the range of severe hypoxia where cell survival is affected after a variable duration of hypoxia. 0.1% O_2 is about the oxygen concentration where aerobic glucolysis yields to ATP production by anaerobic glucolysis (2, 10). The low-energy output by anaerobic glucolysis is in normal tissue and during short periods of severe hypoxia partly countered by increased glucose uptake by the cells, but not necessarily so in tumour tissue (17).

Protein accumulation is greatly diminished, partly due to reduced overall protein synthesis as a result of phosphorylation of the eukaryotic initiation factor eIF2R and partly due to a slight upregulation of overall protein degradation (18). mRNA splicing may be altered at hypoxia (19). The inhibitor of HIF-1, iPAS, is a splicing variant of HIF-3.

At the low end of severe hypoxia, mRNA translation – the most energy-costly process in the cell – is much suppressed, as is ATP production. Respiration is reduced, and there is a limit as to how long cells can stay alive. However, there are two regulatory processes aiming at opposite goals: On the one hand, through cell cycle regulatory processes, cells are able to increase their probability of survival. On the other hand, through induction of apoptosis, damaged cells commit suicide, thereby protecting tissues from problems later on due to multiplication of damaged cells. At 0.05% O₂ (500 ppm O₂), cells are still cycling. However, from 0.05–0.5% O₂ (500–5000 ppm), the ribonucleotide reductase activity is less than maximal owing to lack of molecular oxygen. This results in reduction in DNA synthesis and cell cycling.

In the range between 0.005 and 0.05% O_2 (50 and 500 ppm), cell division will continue until reaching the late G_1 phase (20).

The functional goal of many of the above adaptations can be seen as protection against hypoxia-induced cell death.

Below 0.001% O_2/I Pa (10 ppm) we have the extreme hypoxia (anoxia), at which cell survival is a matter of hours or a few days, and no respiration takes place (2, 21).

We are talking about a state of oxygen deficiency which in vivo is known to exist in areas of solid tumours and infarct tissue (22). Cells introduced abruptly to this condition are arrested at certain restriction points. Cells in S phase stop DNA synthesis momentarily and are lethally damaged after a few hours, while cells in early G1 progress to a restriction point in G1 where they are arrested, but able to survive for many hours. Cells in G2 may complete cell division before they are arrested in the subsequent G1 with a similar ability to survive as G1 cells. Cells may die in the energyrequiring apoptosis or due to other mechanisms. Generally, cells in S phase will survive for less than 10 hours, while cells in G₁ may survive for up to 24-48 hours (21).

The cells' different responses to different oxygen concentrations offer new possibilities for *therapy*.

For tissue with moderate hypoxia, angiogenesis is a target in clinical anticancer trials, and at severe hypoxia bioreductive drugs have proved effective in tumour therapy. In the oxygen concentration range in between, it is to be expected that targeting HIF-1 and downstream processes for enhancement or inhibition



Fig. 1. Relationships between oxygen concentrations and important cell functions. $1\% O_2$ is roughly equal to 1 kPa and 1 kPa equals 10000 ppm at sea level.

will become of major importance in cardiovascular and solid cancer management (13).

A first estimate of relationships between oxygen concentration ranges and important cell functions is given in Fig. 1.

We would like to stress that the present estimates of pericellular oxygen tension, especially in vitro, are hampered by a lack of measurements and use of electrodes measuring at a distance from the cells. Working out better and practical techniques for measuring oxygen is a pressing task.

ACKNOWLEDGEMENTS

This work was supported by EU grant No. LSHC-CT-2003-502932.

REFERENCES

- Ebbesen P, Pettersen EO, Denekamp J, et al. Hypoxia, normoxia and hyperoxia: terminology for medical in vitro cell biology [letter]. Acta Oncol 2000; 39: 247–8.
- 2. Froese G. The respiration of ascites tumour cells at low oxygen concentrations. Biochim Biophys Acta 1962; 57: 509–19.
- 3. Koch CJ, Kruuv J. Measurements of very low oxygen tensions in unstirred liquids. Anal Chem 1972; 44: 1258–63.
- Mamchaoui K, Saumon G. A method for measuring the oxygen consumption of intact cell monolayers. Am J Physiol Lung Cell Mol Physiol 2000; 278: L858–63.
- Baumgardner JE, Otto CM. In vitro intermittent hypoxia: challenges for creating hypoxia in cell culture. Respir Physiol Neurobiol 2003; 136: 131–9.
- Rofstad EK, Sundfor K, Lyng H, Trope CG. Hypoxia-induced treatment failure in advanced squamous cell carcinoma of the uterine cervix is primarily due to hypoxia-induced radiation resistance rather than hypoxia-induced metastasis. Br J Cancer 2000; 83: 354–9.
- Acker H, Lubbers DW, Purves MJ. Local oxygen tension field in the glomus caroticum of the cat and its change at changing arterial PO₂. Pflugers Arch 1971; 329: 136–55.
- Vaupel P, Schlenger K, Knoop C, Hockel M. Oxygenation of human tumors: evaluation of tissue oxygen distribution in breast cancers by computerized O₂ tension measurements. Cancer Res 1991; 51: 3316–22.
- Horsman MR, Nordsmark M, Overgaard J. Techniques to assess the oxygenation of human tumors. State of the art. Strahlenther Onkol 1998; 174(Suppl 4): 2–5.
- Boag JW. Cell respiration as a function of oxygen tension. Int J Radiat Biol Relat Stud Phys Chem Med 1970; 18: 475–8.
- Williams KJ, Telfer BA, Airley RE, et al. A protective role for HIF-1 in response to redox manipulation and glucose deprivation: implications for tumorigenesis. Oncogene 2002; 21: 282– 90.
- Maxwell PH, Wiesener MS, Chang GW, et al. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature 1999; 399: 271-5.
- Semenza GL. Hypoxia-inducible factor 1: control of oxygen homeostasis in health and disease. Pediatr Res 2001; 49: 614–7.
- Koumenis C, Naczki C, Koritzinsky M, et al. Regulation of protein synthesis by hypoxia via activation of the endoplasmic reticulum kinase PERK and phosphorylation of the translation initiation factor eIF2alpha. Mol Cell Biol 2002; 22: 7405– 16.
- Fink T, Kazlauskas A, Poellinger L, Ebbesen P, Zachar V. Identification of a tightly regulated hypoxia-response element in the promoter of human plasminogen activator inhibitor-1. Blood 2002; 99: 2077–83.

- Shweiki D, Itin A, Soffer D, Keshet E. Vascular endothelial growth factor induced by hypoxia may mediate hypoxiainitiated angiogenesis. Nature 1992; 359: 843–5.
- Thews O, Kelleher DK, Esser N, Kraus S, Vaupel P. Lack of association between tumor hypoxia, GLUT-1 expression and glucose uptake in experimental sarcomas. Adv Exp Med Biol 2003; 510: 57–61.
- Kraggerud SM, Sandvik JA, Pettersen EO. Regulation of protein synthesis in human cells exposed to extreme hypoxia. Anticancer Res 1995; 15: 683–6.
- Makino Y, Kanopka A, Wilson WJ, Tanaka H, Poellinger L. Inhibitory PAS domain protein (IPAS) is a hypoxia-inducible splicing variant of the hypoxia-inducible factor-3alpha locus. J Biol Chem 2002; 277: 32405–8.
- Graff P, Amellem O, Andersson KK, Pettersen EO. Role of ribonucleotide reductase in regulation of cell cycle progression during and after exposure to moderate hypoxia. Anticancer Res 2002; 22: 59–68.
- 21. Amellem O, Pettersen EO. Cell inactivation and cell cycle inhibition as induced by extreme hypoxia: the possible role of

cell cycle arrest as a protection against hypoxia-induced lethal damage. Cell Prolif 1991; 24: 127–41.

 Wang Z, Su MY, Nalcioglu O. Applications of dynamic contrast enhanced MRI in oncology: measurement of tumor oxygen tension. Technol Cancer Res Treat 2002; 1: 29–38.

ADDENDUM

The SI unit for pressure is 1 Pascal (Pa) =1 Newton per m², 1 atmosphere =760 mm Hg =1.013 × 10⁵ Pa or approx. 100 kPa. The fraction of oxygen molecules in air is 21%, corresponding to 210 000 ppm (parts per million). Partial pressure of oxygen under 1 atmosphere =21.2 kPa. Dissolved oxygen in water in contact with air under 1 atmosphere pressure, 21.2 kPa, is 220 μ M (micromolar). The lower limit for oxygen partial pressure for full respiration at 37°C (measured with Ehrlich ascites cells) is 0.13 kPa, corresponding to 1.4 μ M oxygen dissolved in the extracellular fluid (Ebbesen et al. 2000 (1)).