



**Pharmaceutical Biology** 

ISSN: 1388-0209 (Print) 1744-5116 (Online) Journal homepage: informahealthcare.com/journals/iphb20

## Screening of Native Plants from Yucatan for Anti-Giardia lamblia. Activity

Sergio R. Peraza-Sánchez, Seydi Poot-Kantún, Luis W. Toores-Tapia, Filogonio May-Pat, Paulino Simá-Polanco & Roberto Cedillo-Rivera

To cite this article: Sergio R. Peraza-Sánchez, Seydi Poot-Kantún, Luis W. Toores-Tapia, Filogonio May-Pat, Paulino Simá-Polanco & Roberto Cedillo-Rivera (2005) Screening of Native Plants from Yucatan for Anti-Giardia lamblia. Activity, Pharmaceutical Biology, 43:7, 594-598, DOI: 10.1080/13880200500301720

To link to this article: https://doi.org/10.1080/13880200500301720



Published online: 07 Oct 2008.



Submit your article to this journal 🕑

Article views: 962



View related articles



# Screening of Native Plants from Yucatan for Anti-*Giardia lamblia* Activity

Sergio R. Peraza-Sánchez<sup>1</sup>, Seydi Poot-Kantún<sup>1</sup>, Luis W. Torres-Tapia<sup>1</sup>, Filogonio May-Pat<sup>1</sup>, Paulino Simá-Polanco<sup>1</sup>, and Roberto Cedillo-Rivera<sup>2</sup>

<sup>1</sup>Centro de Investigación Científica de Yucatán, Mérida, México; <sup>2</sup>Unidad Interinstitucional de Investigación en Salud, Instituto Mexicano del Seguro Social/Universidad Autónoma de Yucatán, Mérida, México

## Abstract

The *in vitro* activity of 10 methanol extracts prepared from native plants collected in the Yucatan Peninsula [*Byrsonima crassifolia* (L.) Kunth, *Cupania dentata* DC., *Diphysa carthagenensis* Jacq., *Dorstenia contrajerva* L., *Gliricidia sepium* (Jacq.) Kunth ex Walp., *Justicia spicigera* Schldl., *Pluchea odorata* (L.) Cass., *Spigelia anthelmia* L., *Tridax procumbens* L., and *Triumfetta semitriloba* Jacq.] was evaluated against *Giardia lamblia* trophozoites. All the extracts showed activity against *G. lamblia* trophozoites. *T. procumbens* was most active, with an IC<sub>50</sub> of  $6.34 \mu g/ml$ , followed by *C. dentata*, 7.59  $\mu g/ml$ , *D. carthagenensis*, 11.53  $\mu g/ml$ , and *B. crassifolia*, 15.55  $\mu g/ml$ .

**Keywords:** Antiprotozoal activity, *Giardia lamblia*, infectious diarrhea, Yucatecan medicinal plants.

## Introduction

*Giardia lamblia* is the most commonly identified protozoan parasite that causes gastrointestinal infection in young children worldwide (Jones, 1991). Clinical manifestations include acute or persistent diarrhea, intestinal malabsorption, and retardation of growth and development (Babb, 1995). The highest incidence of giardiasis (up to 40%) occurs particularly in tropical countries with poor sanitation (Juckett, 1996), and an estimated 200 million people are infected annually by *G. lamblia* in Africa, Asia, and South America (Warren, 1988; Walsh, 1989). In Mexico, the prevalence is 3–60%, depending on the geographic zone and population age (Cedillo-Rivera, 2001). A study carried out in 2002 among people grouped by age in the state of Yucatan, Mexico, showed a total of 3512 cases per 100,000 inhabitants with this infection (Secretaría de Salubridad y Asistencia, 2002). There are several drugs that are effective for the treatment of giardiasis such as metronidazole, tinidazole, guinacrine, furazolidone, albendazole, and nitazoxanide, but most of them cause adverse side effects. In addition, there are reports of drug resistance of G. lamblia to these drugs (McEvoy, 1997; Gardner & Hill, 2001; Upcroft & Upcroft, 2001; Wright et al., 2003). Thus, the discovery of new, effective, and safe antigiardial agents is imperative from other sources. In the current study, a group of selected plants used by Mayan communities that inhabit the Yucatan Peninsula to cure their diarrheal diseases was subjected to a pharmacological evaluation against G. lamblia, the protozoan that causes giardiasis.

## **Materials and Methods**

## Plant materials

Plants were all collected in the Yucatan Peninsula in the period July–October 2001; voucher specimens were authenticated by P. Simá-Polanco and F. May-Pat and deposited at the herbarium of the Centro de Investigación Científica de Yucatán (CICY) under the following code numbers: *Byrsonima crassifolia* (L.) Kunth (Malpighiaceae, FM-1985), *Cupania dentata* DC. (Sapindaceae, PS-2587), *Diphysa carthagenensis* Jacq. (Fabaceae, PS-2577), *Dorstenia contrajerva* L.

Accepted: July 15, 2005

Address correspondence to: Sergio R. Peraza-Sánchez, Unidad de Biotecnología, Centro de Investigación Científica de Yucatán, Calle 43 #130, Chuburná, Mérida, Yucatán, México 97200. Tel.: (52-999) 981-3923 ext. 264; Fax: (52-999) 981-3900; E-mail: speraza@cicy.mx

| Botanical species   | Plant part<br>used   | $IC_{50} (\mu g/ml)^b$   |
|---|--|--|
| Byrsonima crassifolia<br>Cupania dentata<br>Diphysa carthagenensis<br>Dorstenia contrajerva<br>Clinici dia carting                              | Leaves<br>Bark<br>Leaves<br>Whole plant                            | 15.55 (15.33–15.78)<br>7.59 (7.54–7.64)<br>11.53 (11.43–11.62)<br>34.38 (33.47–35.34)  |
| Guricidia sepium<br>Justicia spicigera<br>Pluchea odorata<br>Spigelia anthelmia<br>Tridax procumbens<br>Triumfetta semitriloba<br>Metronidazole | Aerial part<br>Aerial part<br>Whole plant<br>Whole plant<br>Leaves | $\begin{array}{c} 62.66 \ (62.13-65.21) \\ 117.41 \ (117.00-117.79) \\ 59.81 \ (59.59-60.05) \\ 113.27 \ (112.36-114.21) \\ 6.34 \ (6.17-6.50) \\ 46.41 \ (43.16-50.16) \\ 0.21 \ (0.15-0.27) \end{array}$ |

Table 1. Antiprotozoal activity of the tested methanolic extracts.<sup>a</sup>

<sup>*a*</sup>Potency is expressed as IC<sub>50</sub> against axenic trophozoites of *Giardia lamblia in vitro*.

<sup>b</sup>95% confidence interval is expressed in parentheses.

(Moraceae, FM-1984), Gliricidia sepium (Jacq.) Kunth ex Walp. (Fabaceae, FM-1991), Justicia spicigera Schldl. (Acanthaceae, FM-1957), Pluchea odorata (L.) Cass. (Asteraceae, PS-2573), Spigelia anthelmia L. (Loganiaceae, FM-1958), Tridax procumbens L. (Asteraceae, FM-1955), and Triumfetta semitriloba Jacq. (Tiliaceae, PS-2673). These plants were selected by searching in the literature for books and publications on herbal medicine developed by the Mayan people and their descendants. Plant species used in the treatment of diarrhea were selected primarily from a general list (ca. 284 species) provided by Morton (1981). From this list, the number of species was narrowed down to include those that grow in the area of interest, using references given by Roys (1931), Orellana (1987), and Berlin and Berlin (1996). From a survey using computer databases with free access on the World Wide Web (e.g., MEDLINE) and from the records of the NAPRALERT database, a final listing was produced on plant extracts with promising antigiardial activity. Small amounts (100-500 g) of each plant part were collected for biological testing.

#### Preparation of extracts

Powdered and air-dried plant material (100 g) was macerated three-times with methanol (MeOH) at room temperature and evaporated under vacuum in a rotary evaporator to give a semisolid extract. After determining the yield, part of each sample was solubilized in dimethyl sulfoxide (DMSO) just before the biological assay.

#### Giardia lamblia trophozoites

*Giardia lamblia* IMSS:0696:1 isolate, obtained from an individual with symptomatic giardiasis, was used (Cedillo-Rivera et al., 2003). Trophozoites were cultured

in TYI-S-33 modified medium, supplemented with 10% calf serum, and subcultured twice a week; for the assay, trophozoites were tested in their log phase of growth (Cedillo-Rivera et al., 1991).

#### Antiprotozoal assay

The assay has been described in the literature (Cedillo-Rivera & Muñoz, 1992; Cedillo-Rivera et al., 1992; Calzada et al., 1999). The extracts (20 mg) dissolved in 4.0 ml of DMSO were added to tubes containing 1.5 ml of medium in order to get a range of concentrations between 2.5 and 200 µg/ml. This solution was inoculated with G. lamblia to achieve an inoculum of  $5 \times 10^4$  trophozoites/ml. The test included metronidazole (Sigma-Aldrich) as the drug of reference, a control (culture medium with trophozoites and DMSO), and a blank (culture medium). After 48 h at 37°C, parasites were detached by chilling, and 50 µl of each culture tube were subcultured in fresh medium without extracts or drug and incubated for 48 h at 37°C. Cell proliferation was measured with a hemocytometer, and the percentage of trophozoite growth inhibition was calculated by comparison with the controls. The percentage of inhibition calculated for each concentration was transformed into Probit units. The plot of Probit against log concentration was made; the best straight line was determined by regression analysis, and the 50% inhibitory concentration  $(IC_{50})$  values were calculated. The experiments were done in duplicate and repeated at least three times.

#### **Results and Discussion**

The plant that showed the most potent effect against *G. lamblia* was *T. procumbens* (IC<sub>50</sub> =  $6.34 \mu g/ml$ ) (Table 1). This perennial herb grows wild in many tropical countries and is known popularly as hierba del toro ("bull's herb") in Yucatan, Mexico. The entire plant is used by the indigenous people to treat gastrointestinal disorders, stomach pain, and diarrhea (Cáceres et al., 1990; Gupta et al., 1993), bronchial catarrh, dysentery, and wound healing, but it is also applied as an anticoagulant, hair tonic, antifungal, and insect repellent (Taddei & Rosas-Romero, 2000; Akbar et al., 2002). T. procumbens has been analyzed in several biological screenings to look for antibacterial, anticancer, antifertility, antifungal, antihelminthic, antiprotozoal (Bhakuni et al., 1969), antiviral (Taylor et al., 1996), and hepatoprotective activity (Pathak et al., 1991; Saraf & Dixit, 1991). T. procumbens is included in a list of plants most frequently used by the Guatemalan population to cure bacterial and protozoal diseases, and in a screening for antiprotozoal activity the dichloromethane extract of this plant was shown to be active against Trypanosoma cruzi trypomastigotes in vitro at a concentration of 1 mg/ml, while no in vivo toxicity was observed (Cáceres et al., 1998). Extracts of *T. procumbens* have been reported to contain hydrocarbons (Gadre & Gabhe, 1988a), saturated and unsaturated fatty acids (Gadre & Gabhe, 1988b), lipid constituents (Verma & Gupta, 1988), flavonoids (Yadava & Saurabh, 1998; Ali et al., 2001; Akbar et al., 2002), polysaccharides (Raju & Davidson, 1994), and a *bis*bithiophene (Ali & Jahangir, 2002), but up to now no reports on the isolation of metabolites with antiprotozoal activity from this plant have been published.

*C. dentata*, a plant known as *canilla de venado* ("deer's leg") in Mexico, had the second highest effect against *G. lamblia* (IC<sub>50</sub> = 7.59  $\mu$ g/ml). No biological activities for extracts of this species are reported in the literature, nor have chemical components been isolated from it to the present. In traditional medicine, an infusion prepared from leaves of *C. dentata* is used to treat dysentery (Zamora-Martínez & Nieto de Pascual-Pola, 1992).

Evaluation of the leaves of *D. carthagenensis* also showed in this bioassay a potent effect ( $IC_{50} =$ 11.53 µg/ml). *D. carthagenensis* is a small tree with yellow flowers known in Yucatan as *babalché* (Martínez, 1987). Searches in NAPRALERT and in several online databases showed no information on the chemistry of this plant.

B. crassifolia, which showed a significant antiprotozoal effect against G. lamblia (IC<sub>50</sub> = 15.55), is a tropical tree, 3-10 m high, widely distributed in various regions of Mexico and Central and South America, and popularly known as nance in Yucatan, where its round, yellow fruits are highly appreciated and commonly sold in local markets (Martínez-Vázquez et al., 1999). B. crassifolia is one of the most used medicinal plants to treat gastrointestinal disorders, including diarrhea, dysentery, stomach pain, intestinal parasites, and loss of appetite (Cáceres et al., 1990; Béjar & Malone, 1993). It has been described in various studies as a bactericide (Cáceres et al., 1998), and fungicide (Cáceres et al., 1991; 1993), and the ethanol extracts of its leaves have showed some trypanocidal activity (Berger et al., 1998). The genus Byrsonima comprises a small number of species, and not many have been revised chemically (Béjar & Malone, 1993). Reports on previous studies of B. crassifolia indicate the occurrence of benzenoids, glycolipids, proanthocyanidins, triterpenes, and steroids (Béjar et al., 1995; Geiss et al., 1995; Rastrelli et al., 1997).

The whole plant of *D. contrajerva*, when tested in the assay against *G. lamblia*, showed an effective activity ( $IC_{50} = 34.38 \,\mu\text{g/ml}$ ). *D. contrajerva* is an herbaceous plant popularly known as *contra hierba* in Yucatan. It is used as part of local traditional medicines in the south-eastern part of Mexico as a rattle-snake bite tonic and to treat diarrhea, dysentery, and stomach pain (Cáceres et al., 1990; Commerford, 1996; Tovar-Miranda et al., 1998). Chemically, *D. contrajerva* is a species known to be rich in furanocoumarins and flavonoids (Tovar-Miranda et al, 1998; Cáceres et al., 2001).

The results of this study demonstrated that only the methanol extracts of the five plants above described above have an effective growth inhibition of G. lamblia trophozoites in vitro, and they can be considered as good natural sources to isolate the metabolites responsible for the activity. Although G. sepium, J. spicigera, P. odorata, S. anthelmia, and T. semitriloba showed less notable activity, they can also be considered for subjection to a bioassay-directed fractionation, as plant extracts with  $IC_{50}$  values falling within that moderate range (46.41–  $117.41 \,\mu g/ml$ ) have been reported to yield active antiprotozoal metabolites (Calzada et al., 1998; Arrieta et al., 2001; Alanís et al., 2003). None of the 10 plants tested in this work had been previously assayed in vitro against G. lamblia trophozoites except for J. spicigera. In a previous study (Ponce-Macotela et al., 1994, 2001), an ethanol extract of leaves of J. spicigera showed a similar result to the one we obtained  $(IC_{50} = 125 \,\mu g/ml \, vs.)$  $117.43 \,\mu g/ml$ ).

## Acknowledgments

The authors are very grateful to the International Foundation for Science (Research Grant Agreement No. F/3278-1) for financial support to this work. We express thanks to the American Society of Pharmacognosy Foundation for a Research Starter Grant awarded to S.R. Peraza-Sanchez. We are also thankful to NAPRALERT database (University of Illinois at Chicago, Chicago, IL, USA) for providing the ethnobotanical, biological, and chemical information on the plant material used in this work.

#### References

- Akbar E, Malik A, Afza N, Abdul-Hai SM (2002): Flavone glycosides and bergenin derivatives from *Tridax procumbens. Heterocycles* 57: 733–739.
- Alanís AD, Calzada F, Cedillo-Rivera R, Meckes M (2003): Antiprotozoal activity of the constituents of *Rubus* coriifolius. Phytother Res 17: 681–682.
- Ali MS, Jahangir M (2002): A bis-bithiophene from Tridax procumbens L. (Asteraceae). Nat Prod Lett 16: 217–221.
- Ali M, Ravinder E, Ramachandram R (2001): A new flavonoid from the aerial parts of *Tridax procumbens*. *Fitoterapia* 72: 313–315.
- Arrieta J, Reyes B, Calzada F, Cedillo-Rivera R, Navarrete A (2001): Amoebicidal and giardicidal compounds from the leaves of *Zanthoxylum liebmannianun*. *Fitoterapia* 72: 295–297.
- Babb RR (1995): Giardiasis. Taming this pervasive parasitic infection. *Postgrad Med* 98: 155–158.

- Béjar E, Malone MH (1993): Pharmacological and chemical screening of *Byrsonima crassifolia*, a medicinal tree from Mexico. Part I. J Ethnopharmacol 39: 141–158.
- Béjar E, Amarquaye A, Che CT, Malone MH, Fong HHS (1995): Constituents of *Byrsonima crassifolia* and their spasmogenic activity. *Int J Pharmacog 33*: 25–32.
- Berger I, Barrientos AC, Cáceres A, Hernández M, Rastrelli L, Passreiter CM, Kubelka W (1998): Plants used in Guatemala for the treatment of protozoal infections.
  II. Activity of extracts and fractions of five Guatemalan plants against *Trypanosoma cruzi*. J Ethnopharmacol 62: 107–115.
- Berlin EA, Berlin B (1996): Medical Ethnobiology of the Highland Maya of Chiapas, Mexico. The Gastrointestinal Diseases. Princeton, NJ, Princeton University Press.
- Bhakuni DS, Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN (1969): Screening of Indian plants for biological activity. Part II. *Indian J Exp Biol* 7: 250–262.
- Cáceres A, Cano O, Samayoa B, Aguilar L (1990): Plants used in Guatemala for the treatment of gastrointestinal disorders. 1. Screening of 84 plants against enterobacteria. J Ethnopharmacol 30: 55–73.
- Cáceres A, López BR, Girón MA, Logemann H (1991): Plants used in Guatemala for the treatment of dermatophytic infections. 1. Screening for antimycotic activity of 44 plant extracts. J Ethnopharmacol 31: 263–276.
- Cáceres A, López B, Juárez X, Del Aguila J, García S (1993): Plants used in Guatemala for the treatment of dermatophytic infections. 2. Evaluation of antifungal activity of seven American plants. *J Ethnopharmacol* 40: 207–213.
- Cáceres A, López B, González S, Berger I, Tada I, Maki J (1998): Plants used in Guatemala for the treatment of protozoal infections. I. Screening of activity to bacteria, fungi and American trypanosomes of 13 native plants. *J Ethnopharmacol 62*: 195–202.
- Cáceres A, Rastrelli L, De Simone F, De Martino G, Saturnino C, Saturnino P, Aquino R (2001): Furanocoumarins from the aerial parts of *Dorstenia contrajerva*. *Fitoterapia 72*: 376–381.
- Calzada F, Alanís AD, Meckes M, Tapia-Contreras A, Cedillo-Rivera R (1998): *In vitro* susceptibility of *Entamoeba histolytica* and *Giardia lamblia* to some medicinal plants used by the people of Southern Mexico. *Phytother Res* 12: 70–72.
- Calzada F, Cerda-García Rojas CM, Meckes M, Cedillo-Rivera R, Bye R, Mata R (1999): Geranins A and B, new antiprotozoal A-type proanthocyanidins from *Geranium niveum. J Nat Prod* 62: 705–709.
- Cedillo-Rivera R (2001): Parasitosis Intestinales. In: Kumate J, ed., *Manual de Infectología Clínica*, 16th ed. Mexico City, Méndez Editores, pp. 201–209.
- Cedillo-Rivera R, Muñoz O (1992): In vitro susceptibility of Giardia lamblia to albendazol, mebendazol and

other chemotherapeutic agents. J Med Microbiol 37: 221–224.

- Cedillo-Rivera R, Ramírez A, Muñoz O (1992): A rapid colorimetric assay with the tetrazolium salt MTT and phenazine methosulfate (PMS) for viability of *Entamoeba histolytica*. *Arch Med Res 23*: 59–61.
- Cedillo-Rivera R, Enciso J, Ortega-Pierres G, Martínez-Palomo A (1991): Isolation and axenization of *Giardia lamblia* isolates obtained from symptomatic and asymptomatic patients in Mexico. *Arch Med Res 22*: 79–85.
- Cedillo-Rivera R, Darby JM, Enciso-Moreno JA, Ortega-Pierres G, Ey PL (2003): Genetic homogeneity of axenic isolates of *Giardia intestinalis* derived from acute and chronically infected individuals in Mexico. *Parasitol Res* 90: 119–123.
- Commerford SC (1996): Medicinal plants of two Mayan healers from San Andrés, Petén, Guatemala. *Econ Bot* 50: 327–336.
- Gadre A, Gabhe SY (1988a): Hydrocarbons of *Tridax pro*cumbens. Indian J Nat Prod 4: 15–16.
- Gadre A, Gabhe SY (1988b): Saturated and unsaturated fatty acids of *Tridax procumbens*. *Indian J Pharm Sci* 38: 149–155.
- Gardner TB, Hill DR (2001): Treatment of giardiasis. *Clin Microbiol Rev 14*: 114–128.
- Geiss F, Heinrich M, Hunkler D, Rimpler H (1995): Proanthocyanidins with (+)-epicatechin units from *Byrsonima crassifolia* bark. *Phytochemistry* 39: 635–643.
- Gupta S, Yadava JNS, Tandon JS (1993): Antisecretory activity (antidiarrhoeal) activity of Indian medicinal plants against *Escherichia coli* enterotoxin-induced secretion in rabbit and Guinea pig ileal loop models. *Int J Pharmacog 31*: 198–204.
- Jones JE (1991): Giardiasis. Prim Care 18: 43-52.
- Juckett G (1996): Intestinal protozoa. Am Fam Phys 53: 2507–2516.
- Martínez M (1987): *Catálogo de nombres vulgares y científicos de plantas mexicanas*. Mexico City, Fondo de Cultura Económica.
- Martínez-Vázquez M, González-Esquinca AR, Cazares-Luna L, Moreno-Gutiérrez MN, García-Argáez AN (1999): Antimicrobial activity of *Byrsonima crassifolia* (L.) H.B.K. *J Ethnopharmacol* 66: 79–82.
- McEvoy GK, ed. (1997): *AHFS Drug Information*. Bethesda, MD, American Society of Health-System Pharmacists.
- Morton JF (1981): Atlas of Medicinal Plants of Middle America. Bahamas to Yucatan. Springfield, IL, Charles C. Thomas Publisher.
- Orellana SL (1987): *Indian Medicine in Highland Guatemala*. Albuquerque, NM, University of New Mexico Press.
- Pathak AK, Saraf S, Dixit VK (1991): Hepatoprotective activity of *Tridax procumbens*. Part I. *Fitoterapia* 62: 307–313.
- Ponce-Macotela M, Navarro-Alegría I, Martínez-Gordillo MN, Alvaro-Chacón R (1994): Efecto antigiardiásico

*in vitro* de 14 extractos de plantas. *Rev Invest Clin 46*: 343–347.

- Ponce-Macotela M, Rufino-González Y, De la Mora-De la Mora JI, González-Maciel A, Reynoso-Robles R, Martínez-Gordillo MN (2001): Mortality and morphological changes in *Giardia duodenalis* induced by exposure to ethanolic extracts of *Justicia spicigera*. Proc West Pharmacol Soc 44: 151–152.
- Raju TS, Davidson EA (1994): Structural features of watersoluble novel polysaccharide components from the leaves of *Tridax procumbens* L. *Carbohydr Res* 258: 243–254.
- Rastrelli L, De Tommassi N, Berger I, Cáceres A, Saravia A, De Simone F (1997): Glycolipids from *Byrsonima* crassifolia. Phytochemistry 45: 647–650.
- Roys RL (1931): *The Ethno-Botany of the Maya*. Philadelphia, PA, Institute for the Study of Human Issues.
- Saraf S, Dixit VK (1991): Hepatoprotective activity of *Tridax procumbens*. Part II. *Fitoterapia* 62: 534–536.
- Secretaría de Salubridad y Asistencia (SSA) (2002): Sistema Único de Información para la Vigilancia Epidemiológica, Dirección General de Epidemiología. Mexico City, SSA.
- Taddei A, Rosas-Romero AJ (2000): Bioactivity studies of extracts from *Tridax procumbens*. *Phytomedicine* 7: 235–238.
- Taylor RSL, Hudson JB, Manandhar NP, Towers GHN (1996): Antiviral activities of medicinal plants of Southern Nepal. *J Ethnopharmacol* 53: 97–104.

- Tovar-Miranda R, Cortés-García R, Santos-Sánchez NF, Joseph-Nathan P (1998): Isolation, total synthesis, and relative stereochemistry of a dihydrofurocoumarinfrom *Dorstenia contrajerva*. J Nat Prod 61: 1216–1220.
- Upcroft P, Upcroft JA (2001): Drug targets and mechanisms of resistance in the anaerobic protozoa. *Clin Microbiol Rev 14*: 150–164.
- Verma RK, Gupta MM (1988): Lipid constituents of *Tridax* procumbens. Phytochemistry 27: 459–463.
- Walsh JA (1989): Estimating the burden of illness in the tropics. In: Warren KS, Mahmoud AF, eds., *Tropical and Geographic Medicine*. New York, McGraw-Hill, pp. 1073–1085.
- Warren KS (1988): The global impact of parasitic diseases. In: Englund PT, Sher A, eds., *The Biology of Parasitism: A Molecular and Immunological Approach*. New York, Alan R. Liss, pp. 3–12.
- Wright JM, Dunn LA, Upcroft P, Upcroft JA (2003): Efficacy of antigiardial drugs. *Expert Opin Drug Safety 2*: 529–541.
- Yadava RN, Saurabh K (1998): A new flavone glycoside: 5,7,4'-trihydroxy-6,3'-dimethoxy flavone 5-*O*-α-L-rhamnopyranoside from the leaves of *Tridax procumbens* L. *J Asian Nat Prod Res 1*: 147–152.
- Zamora-Martínez MC, Nieto de Pascual-Pola C (1992): Medicinal plants used in some rural populations of Oaxaca, Puebla, and Veracruz, Mexico. J Ethnopharmacol 35: 229–257.