



Journal of Enzyme Inhibition and Medicinal Chemistry

ISSN: 1475-6366 (Print) 1475-6374 (Online) Journal homepage: informahealthcare.com/journals/ienz20

Synthesis, spectroscopic, and antibacterial activity of tetraazamacrocyclic complexes of trivalent chromium, manganese, and iron

D.P. Singh, Vandna Malik, Ramesh Kumar & Jitender Singh

To cite this article: D.P. Singh, Vandna Malik, Ramesh Kumar & Jitender Singh (2009) Synthesis, spectroscopic, and antibacterial activity of tetraazamacrocyclic complexes of trivalent chromium, manganese, and iron, Journal of Enzyme Inhibition and Medicinal Chemistry, 24:5, 1201-1206, DOI: 10.1080/14756360902779383

To link to this article: https://doi.org/10.1080/14756360902779383



Published online: 23 Sep 2009.

|--|

Submit your article to this journal 🖸

Article views: 574



View related articles 🗹



Citing articles: 1 View citing articles 🗹

RESEARCH ARTICLE

Synthesis, spectroscopic, and antibacterial activity of tetraazamacrocyclic complexes of trivalent chromium, manganese, and iron

D.P. Singh¹, Vandna Malik¹, Ramesh Kumar¹, and Jitender Singh²

¹Department of Chemistry, National Institute of Technology, Kurukshetra, India, and ²D.M. Division, National Dairy Research Institute, Karnal, India

Abstract

A new series of macrocyclic complexes of type $[M(TML)X]X_2$, where M = Cr(III), Mn(III), or Fe(III), TML is tetradentate macrocyclic ligand, and $X = CI^-$, NO_3^- , CH_3COO^- for Cr(III), Fe(III) and $X = CH_3COO^-$ for Mn(III), has been synthesized by condensation of benzil and succinyldihydrazide in the presence of metal salt. The complexes have been so formulated due to the 1:2 electrolytic nature of these complexes as shown by conductivity measurements. The complexes have been characterized with the help of various physicochemical techniques such as elemental analysis, molar conductance, electronic and infrared spectral studies, and magnetic susceptibility. On the basis of these studies, a five-coordinate distorted square pyramidal geometry, in which two nitrogens and two carbonyl oxygen atoms are suitably placed for coordination toward the metal ion, has been proposed for all the complexes. The complexes have been tested for their *in vitro* antibacterial activity. Some of the complexes show remarkable antibacterial activities against some selected bacterial strains. The minimum inhibitory concentrations shown by these complexes have been compared with those shown by some standard antibiotics such as linezolid and cefaclor.

Keywords: Macrocyclic complexes; template; infrared; ligand; antibacterial

Abbreviations: MIC, minimum inhibitory concentration; MTCC, microbial type culture collection; MHA, Muller–Hinton agar; CFU, colony forming unit; BM, Bohr magneton; DMF, N,N-dimethylformamide; DMSO, dimethyl sulfoxide; BHI, brain heart infusion

Introduction

Transition metals play an important role in the development of new molecular materials which show magnetic properties and find applications in various fields. During the past few decades, the synthesis of macrocyclic complexes has been a fascinating area of research growing at a very fast pace, owing to their resemblance to naturally occurring macrocycles and analytical, industrial, and medical applications¹⁻⁵. Macrocyclic nickel complexes find use in DNA recognition and oxidation⁶ while macrocyclic copper complexes find use in DNA binding and cleavage⁷. Macrocyclic metal complexes of lanthanides, e.g. Gd³⁺ are used as magnetic resonance imaging (MRI) contrast agents^{8,9}. Macrocyclic metal chelating agents (DOTA) are useful for detecting tumor lesions¹⁰. The chemistry of macrocyclic complexes is also important due to their use as dyes and pigments¹¹ as well as nuclear magnetic resonance (NMR) shift reagents¹². Some macrocyclic complexes have been found to exhibit potential antibacterial activities¹³. Prompted by these, in the present article a new series of macrocyclic complexes of Cr(III), Fe(III), and Mn(III) obtained by template condensation reaction of succinyldihydrazide and benzil is reported. The complexes have been characterized with the help of infrared (IR) and magnetic susceptibilities, elemental analysis, and molar conductance. These complexes have also been tested for their *in vitro* antibacterial activities. Some complexes show remarkable antibacterial activity.

Address for Correspondence: D. P. Singh, Department of Chemistry, National Institute of Technology, Kurukshetra-136 119, India. Tel: +911744 233 512, +911744 238 241. Fax: +911744 238 050. E-mail: dpsinghchem@yahoo.co.in

⁽Received 10 October 2008; revised 23 December 2008; accepted 21 January 2009)

Experimental

Chemistry

Materials

Diethylsuccinate, hydrazine hydrate, and benzil were obtained from SRL Pvt. Ltd., Mumbai. All other chemicals used were of AnalR grade.

Analytical and physical measurements

Microanalyses of C, H, and N were carried out at the Sophisticated Analytical Instrument Facility, CDRI, Lucknow. The metal contents were determined by standard ethylenediaminetetraacetic acid (EDTA) methods. Electronic spectra (DMF) were recorded on a Cary 14 spectrophotometer. The magnetic susceptibility measurements were carried at IIT, Roorkee. The IR spectra were recorded on an infrared spectrophotometer in the range 4000–200 cm⁻¹. The conductivity was measured using a digital conductivity meter (HPG System, G-3001).

Isolation of complexes

All the reported complexes were prepared using the template method. To a stirring methanolic solution (\sim 50 cm³) of succinyldihydrazide (10 mmol) was added trivalent chromium, manganese, and iron salt (10 mmol) dissolved in a minimum quantity of methanol (20 cm³). The resulting solution was refluxed for 0.5 h. After that, benzil (10 mmol) dissolved in \sim 20 mL of methanol was added to the refluxing mixture which was refluxed again for 6–8 h. On overnight cooling, a dark colored precipitate formed, which was filtered, washed with methanol, acetone, and diethyl ether, and dried *in vacuo* (yield 45%). The complexes were found to be soluble in DMF and DMSO, but were insoluble in common organic solvents and water. They were found to be thermally stable up to \sim 240°C and then they decomposed.

The template syntheses of the complexes may be represented by the scheme shown in Figure 1, where M = Cr(III), Mn(III), Fe(III), and $X = Cl^-$, NO_3^- , CH_3COO^- for Cr(III) and Fe(III) and CH_3COO^- for Mn(III).

Pharmacology

In vitro antibacterial activity

Some of the synthesized macrocyclic complexes were tested for *in vitro* antibacterial activity against some bacterial strains using spot-on-lawn on Muller–Hinton agar by a reported method¹⁴. *Test pathogens* Four test pathogenic bacterial strains, viz. *Bacillus cereus* (MTCC 1272), *Salmonella typhi* (MTCC 733), *Escherichia coli* (MTCC 739), and *Staphylococcus aureus* (MTCC 1144), were considered for determination of the MIC (minimum inhibitory concentration) of selected complexes.

Culture conditions The test pathogens were subcultured aerobically using brain heart infusion agar (HiMedia, Mumbai, India) at $37^{\circ}C/24$ h. Working cultures were stored at 4°C in brain heart infusion (BHI) broth (HiMedia), while stock cultures were maintained at $-70^{\circ}C$ in BHI broth containing 15% (v/v) glycerol (Qualigens, Mumbai, India). Organism was grown overnight in 10 mL BHI broth, which was then centrifuged at 5000g for 10 min, and the pellet was suspended in 10 mL of phosphate buffered saline (PBS, pH 7.2). The optical density at 545 nm (OD-545) was adjusted to obtain 10⁸ CFU/mL, followed by plating serial dilution onto plate count agar (HiMedia).

Determination of minimum inhibitory concentration The minimum inhibitory concentration (MIC) is the lowest concentration of the antimicrobial agent that prevents the development of viable growth after overnight incubation. The antimicrobial activity of the compounds was evaluated using the spot-on-lawn method on Muller-Hinton agar (MHA; HiMedia). Soft agar was prepared by adding 0.75% agar in Muller-Hinton broth (HiMedia). The soft agar was inoculated with 1% of 108 CFU/mL of the test pathogen, and 10 mL was overlaid on MHA. From a ×1000 solution of compound (1 mg/ml of DMSO), $\times 1$, $\times 2$, $\times 4$, $\times 8$, $\times 16$, $\times 32$, $\times 64$, and $\times 128$ solutions were prepared. Dilutions of standard antibiotics (linezolid and cefaclor) were also prepared in the same manner. Five microliters of the appropriate dilution was spotted on the soft agar and incubated at 37°C for 24h. Zones of inhibition of the compounds were considered after subtraction of the inhibition zones of DMSO. A negative control (with no compound) was also observed.

Results and discussion

Chemistry

The analytical data show the formula of macrocyclic complexes as: $[M(C_{18}H_{16}N_4O_2)X]X_2$, where M = Cr(III), Mn(III), Fe(III) and $X = Cl^-$, NO_3^- , CH_3COO^- for Cr(III) and Fe(III) and $X = CH_3COO^-$ for Mn(III). The test for anions was positive before and after decomposing the complexes with



Figure 1. Synthesis of the complexes.

		Found (calculated) (%)					
Series no.	Complex	М	С	Н	Ν	Color	Mol. wt.
1	$[Cr(C_{18}H_{16}N_{4}O_{2})Cl]Cl_{2}$	10.88 (10.86)	45.21 (45.14)	3.29 (3.34)	11.79 (11.70)	Light green	478.5
2	$[Cr(C_{18}H_{16}N_4O_2)(NO_3)](NO_3)_2$	9.23 (9.31)	38.67 (38.70)	2.95 (2.86)	17.62 (17.56)	Light yellow	558
3	$[Cr(C_{18}H_{16}N_{4}O_{2})(OAc)](OAc)_{2}$	9.49 (9.47)	52.53 (52.45)	4.48 (4.55)	10.23 (10.20)	Green	549
4	$[Mn(C_{18}H_{16}N_4O_2)(OAc)](OAc)_2$	9.89 (9.96)	52.26 (52.17)	4.49 (4.52)	10.27 (10.14)	Off white	552
5	$[Fe(C_{18}H_{16}N_4O_2)Cl]Cl_2$	11.64 (11.55)	44.83 (44.76)	3.29 (3.31)	11.54 (11.60)	Dark green	482.5
6	$[Fe(C_{18}H_{16}N_4O_2)(NO_3)](NO_3)_2$	9.87 (9.96)	38.39 (38.43)	2.93 (2.84)	17.21 (17.43)	Dark red	562
7	$[Fe(C_{18}H_{16}N_4O_2)(OAc)](OAc)_2$	10.19 (10.12)	52.12 (52.07)	4.64 (4.52)	10.18 (10.12)	Yellow	553

Table 1. Analytical data for trivalent chromium, manganese, and iron complexes derived from succinyldihydrazide and benzil.

concentrated HNO_3 , indicating their presence inside as well as outside the coordination sphere. Conductivity measurements in DMSO indicated them to be electrolytic in nature $(140-150 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1})^{15}$. All compounds gave satisfactory elemental analysis results, as shown in Table 1.

IR spectra

In the infrared spectrum of succinyldihydrazide, a pair of bands corresponding to ν (NH₂) is present at ~3200 cm⁻¹ and ~3250 cm⁻¹, but this was absent in the IR spectra of all the complexes¹⁶. However, a single, broad, medium band at ~3350-3400 cm¹ was observed in the spectra of all the complexes, which may be assigned to ν (NH)^{17,18}. A strong peak at ~1665 cm⁻¹ in the IR spectrum of succinvldihydrazide is assigned to the >C=O group of the CONH moiety. This peak was shifted to a lower frequency $(\sim 1625 - 1640 \text{ cm}^{-1})$ in the spectra of all the complexes^{19,20}, suggesting coordination of the oxygen of the carbonyl group with the metal. Further, no strong absorption band was observed near 1700 cm⁻¹ in the IR spectra of all complexes, as observed in the spectrum of benzil. This indicates the absence of >C=O groups of the benzil moiety in the complexes. These facts confirm the condensation of carbonyl groups of the benzil and amino groups of succinyldihydrazide^{21,22}. The IR spectra of the complexes showed a new strong absorption band in the region ~1590–1610 cm⁻¹, which may be attributed to ν (C=N)^{23,24}. These results provide strong evidence for the formation of a macrocyclic frame²⁵. The lower value of ν (C=N) indicates coordination of the nitrogens of azomethine and the metal²⁶. The bands present at ~1350–1000 cm⁻¹ are assigned to ν (C–N) vibration. The bands presents at ~3040 cm⁻¹ may be assigned to ν (C-H) vibrations of the benzil moiety.

Far infrared spectra

The far infrared spectra showed bands in the region ~425–445 cm⁻¹ corresponding to ν (M–N) vibrations in all the complexes²⁷⁻²⁹. The presence of a band in all the complexes at the ~425–445 cm⁻¹ region originates from (M–N) azomethine vibration modes and gives an idea about coordination of azomethine nitrogens with the metal³⁰. The bands present at ~300–315 cm⁻¹ are due to ν (M–Cl)^{27,29} and bands present at ~220–250 cm⁻¹ in all nitrato complexes to ν (M–O)²⁷.

Magnetic measurements and electronic spectra

Chromium complexes Magnetic moments of chromium complexes were found in the range of 4.25–4.60 BM. The

electronic spectra of chromium complexes showed bands at ~9010–9260, 13,000–13,550, 17,460–18,300, 27,400–27,800, and 34,850 cm⁻¹. However, these spectral bands cannot be interpreted in terms of four or six coordinated environments around the metal atom. In turn, the spectra are comparable to those of five-coordinated Cr(III) complexes, whose structures have been confirmed with the help of X-ray measurements³¹. Thus, keeping in view the analytical data and electrolytic nature of these complexes, a five-coordinated square pyramidal geometry may be assigned to these complexes. Thus, assuming the symmetry C_{4V} for these complexes³², the various spectral bands may be assigned as: ⁴B₁ \rightarrow ⁴E^a, ⁴B₁ \rightarrow ⁴B₂, ⁴B₁ \rightarrow ⁴A₂, and ⁴B₁ \rightarrow ⁴E^b.

Manganese complex The magnetic moment of the manganese complex was found to be 4.80 BM. The electronic spectrum of the manganese complex showed three d–d bands which lay in the regions 12,200, 16,100, and 35,400 cm⁻¹. The higher energy band at 35,400 cm⁻¹ may be assigned to charge transfer transitions. The spectrum resembled those reported for five-coordinate square pyramidal manganese porphyrins³². This idea is further supported by the presence of the broad-ligand field band at 20,400 cm⁻¹, diagnostic of C_{4V} symmetry, and thus the various bands may be assigned as follows: ${}^{5}B_{1} \rightarrow {}^{5}A_{1}$, ${}^{5}B_{1} \rightarrow {}^{5}B_{2}$, and ${}^{5}B_{1} \rightarrow {}^{5}E$, respectively. The band assignment in single electron transition may be made as: $d_{2^{2}} \rightarrow d_{x^{2}-y^{2}}$, $d_{xy} \rightarrow d_{x^{2}-y^{2}}$, and d_{xy} , $d_{yz} \rightarrow d_{x^{2}-y^{2}}$, respectively, in order of increasing energy. However, the complexes do not have idealized C_{4V} symmetry.

Iron complexes The magnetic moments of the iron complexes lay in the range 5.80–5.92 BM. The electronic spectra of the trivalent iron complexes showed various bands, 9800–9900, 15,500–15,550, 27,600–27,700 cm⁻¹, and these bands do not suggest octahedral or tetrahedral geometry around the metal atom. The spectral bands are consistent with the range of those reported for five-coordinate square pyramidal iron(III) complexes³³. Assuming C_{4V} symmetry for these complexes, the various bands can be assigned as: $d_{xy} \rightarrow d_{xz}$, d_{yz} and $d_{xy} \rightarrow d_{z^2}$. Any attempt to make an accurate assignment is difficult due to interactions of the metal-ligand pi-bond systems lifting the degeneracy of the d_{xz} and d_{xy} pair.

Based on elemental analyses, conductivity and magnetic measurements, and electronic, IR and far IR spectral studies, the general structure shown in Figure 2 may be proposed for these complexes.

Biology

The synthesized macrocyclic complexes were tested for their *in vitro* antibacterial activity against four test bacteria, *Bacillus cereus* (MTCC 1272), *Salmonella typhi* (MTCC 733), *Escherichia coli* (MTCC 739), and *Staphylococcus aureus* (MTCC 1144). The MIC (minimum inhibitory concentration) shown by the complexes against these bacterial strains was compared with the MIC shown by standard antibiotics linezolid and cefaclor (Table 2 and Figure 3). Complex 1 showed



Figure 2. General structure of the complexes, where M = Cr(III), Mn(III), Fe(III), $X = Cl^-$, NO_3^- , CH_3COO^- for Cr(III) and Fe(III) and $X = CH_3COO^-$ for Mn(III).

a minimum inhibitory concentration of 32µg/mL against bacterial strain Salmonella typhi (MTCC 733), which is equal to the MIC shown by standard antibiotic linezolid against the same bacterial strain. Complex 3 registered a minimum inhibitory concentration of 4µg/mL against bacterial strain Staphylococcus aureus (MTCC 1144), which is equal to the MIC shown by standard antibiotic linezolid against the same bacterial strain. Further, it showed a minimum inhibitory concentration of 16 µg/mL against bacterial strain Escherichia coli (MTCC 739), which is equal to the MIC shown by standard antibiotic linezolid against the same bacterial strain. Complex 4 registered a minimum inhibitory concentration ranging from 16 to 64µg/mL. Complex 6 registered a minimum inhibitory concentration of 8 µg/mL against the bacterial strain Bacillus cereus (MTCC 1272), which is equal to the MIC shown by standard antibiotic cefaclor against the same bacterial strain. Complex 7 registered a minimum inhibitory concentration of 32 µg/mL against bacterial strain Salmonella typhi (MTCC 733), which is equal to the MIC shown by standard antibiotic linezolid against the same bacterial strain. Among the series under test for determination of the minimum inhibitory concentration, complex 3 was found to be the most potent.

Table 2. Minimum inhibitory concentration (MIC) shown by complexes against test bacteria using agar dilution assay.

		MIC (µg/mL)				
Series no.	Complex	a	b	С	d	
1	$[Cr(C_{18}H_{16}N_{4}O_{2})Cl]Cl_{2}$	16	64	64	32	
2	$[Cr(C_{18}H_{16}N_4O_2)(NO_3)](NO_3)_2$	128	64	128	128	
3	$[Cr(C_{18}H_{16}N_4O_2)(OAc)](OAc)_2$	32	4	16	64	
4	$[Mn(C_{18}H_{16}N_{4}O_{2})(OAc)](OAc)_{2}$	_	16	64	>128	
5	$[Fe(C_{18}H_{16}N_{4}O_{2})Cl]Cl_{2}$	_	128	>128	64	
6	$[Fe(C_{18}H_{16}N_4O_2)(NO_3)](NO_3)_2$	8	16	64	64	
7	$[Fe(C_{18}H_{16}N_4O_2)(OAc)](OAc)_2$	128	_	64	32	
	Cefaclor	8	2	8	16	
	Linezolid	4	4	16	32	

Note. —, no activity; a, Bacillus cereus (MTCC 1272); b, Staphylococcus aureus (MTCC 1144); c, Escherichia coli (MTCC 739); d, Salmonella typhi (MTCC 733); cefaclor and linezolid are standard antibiotics.



Figure 3. Comparison of minimum inhibitory concentration (MIC) of complexes with those of standard antibiotics up to concentration of 64 µg/mL. a, *Bacillus cereus* (MTCC 1272); b, *Staphylococcus aureus* (MTCC 1144); c, *Escherichia coli* (MTCC 739); d, *Salmonella typhi* (MTCC 733); cefaclor and linezolid are standard antibiotics.

However, complexes 2, 4, and 5 showed poor antibacterial activity or no activity against all bacterial strains among the whole series.

Conclusions

Based on elemental analyses, conductivity and magnetic measurements, and electronic, IR, and far IR spectral studies, the structure as shown in Figure 2 may be proposed for these complexes.

It has been suggested that chelation/coordination reduces the polarity of the metal ion mainly because of partial sharing of its positive charge with the donor group within the whole chelate ring system. This process of chelation thus increases the lipophilic nature of the central metal atom, which in turn favors its permeation through the lipid layer of the membrane, thus causing the metal complex to cross the bacterial membrane more effectively and hence increase the activity of the complex. Aside from this, many other factors such as solubility, dipole moment, and conductivity influenced by the metal ion may be possible reasons for the remarkable antibacterial activities of these complexes³⁴⁻³⁷. It also has been observed that some moieties, such as the azomethine linkage or heteroaromatic nucleus, introduced into such compounds exhibit extensive biological activities that may be responsible for the increase in hydrophobic character and liposolubility of the molecules in crossing the cell membrane of the microorganism, thus enhancing the biological utilization ratio and activity of complexes^{38,39}.

Acknowledgements

D. P. Singh thanks the University Grants Commission, New Delhi, for financial support in the form of a Major Research Project. Thanks are also due to the authorities of NIT, Kurukshetra, for providing necessary research facilities.

Declaration of interest: The authors report no conflicts of interest.

References

- Gloe K, ed. Macrocyclic Chemistry: Current Trends and Future Perspectives. Dordrecht: Springer, 2005.
- 2. Lindoy LF, ed. The Chemistry of Macrocyclic Ligand Complexes. Cambridge: Cambridge University Press, 1989.
- 3. Constable EC, ed. Coordination Chemistry of Macrocyclic Compounds. Oxford: Oxford University Press, 1999.
- Singh DP, Kumar R, Malik V, Tyagi P. Synthesis and characterization of complexes of Co(II), Ni(II), Cu(II), Zn(II), Cd(II) with macrocycle 3,4,11,12-tetraoxo-1,2,5,6,9,10,13,14-octaazacyclohexadeca-6,8,14,16tetraene and their biological screening. Transition Met Chem 2007;32:1051-5.
- Singh DP, Kumar R, Malik V, Tyagi P. Template synthesis, spectroscopic studies and biological activities of macrocyclic complexes derived from thiocarbohydrazide and glyoxal. J Enzyme Inhib Med Chem 2007;22:177-82.

- Muller JG, Chen X, Dadiz AC, Rokita SE, Burrows CJ. Macrocyclic nickel complexes in DNA recognition and oxidation. Pure Appl Chem 1993;65:545-50.
- Liu J, Lu TB, Deng H, Ji LN, Qu LH, Zhou H. Synthesis DNA-binding and cleavage studies of macrocyclic copper(II) complexes. Transition Met Chem 2003;28:116-21.
- 8. Paryzek WR, Krzyminiewska VP. Yttrium(III) complexes of pentadentate Schiff base macrocyclic ligands with N_3O_2 and N_5 set of donor atoms. Polyhedron 1995;14:2059–62.
- Watson AD, Rockladge SM. In: Higgins CB, ed. Magnetic Resonance Imaging of the Body. New York: Raven Press, 1992.
- Kosmas C, Snook D, Gooden CS, Courtenay-Luck NS, McCall MJ, Claude F, Meares CF, Epenetos AA. Development of humoral immune responses against a macrocyclic chelating agent (DOTA) in cancer patients receiving radioimmunoconjugates for imaging and therapy. Cancer Res 1992;52:904–11.
- 11. Seto J, Tamura S, Asai N, Kishii N, Kijima Y, Matsuzawa N. Macrocyclic functional dyes: applications to optical disk media, photochemical hole burning and non-linear optics. Pure Appl Chem 1996;68:1429–34.
- Dong W, Yang R, Yan L. Rare earth complexes of a new Schiff base macrocyclic ligand derived from 2,2'-(ethylenedioxy) bisbenzaldehyde and 2,2'-(ethylenedioxy)bisbenzyl amine. Indian J Chem 2001;40A:202-6.
- Singh DP, Kumar R, Tyagi P. Template synthesis, spectroscopic studies and biological screening of macrocyclic complexes derived from thiocarbohydrazide and benzil. Transition Met Chem 2006;31:970-3.
- 14. Singh DP, Kumar R, Singh J. Biologically active trivalent metal macrocyclic complexes derived from oxalyldihydrazide and benzil: synthesis and spectroscopic approach. Eur J Med Chem 2009 (in press).
- Kumar R, Singh R. Chromium(III) complexes with different chromospheres macrocyclic ligand: synthesis and spectroscopic studies. Turk J Chem 2006;30:77-87.
- Prasad RN, Mathur M, Upadhyay A. Synthesis and spectroscopic studies of Cr^{III}, Fe^{III} and Co^{II} complexes of hexaazamacrocycles. J Indian Chem Soc 2007;84:1202-4.
- 17. Khan TA, Rather MA, Jahan N, Varkey SP, Shakir M. Tetraoxotetraamide macrocyclic complexes. Transition Met Chem 1998;23:283–5.
- 18. Singh AK, Panwar A, Singh R, Beniwal S. New bis-macrocyclic complexes with transition metal ion. Transition Met Chem 2003;28:160–2.
- 19. Pavia DL, Lampman GM, Kriz GS.Introduction to Spectroscopy. New York: Harcourt College Publishers, 2001.
- Nishat N, Ud-din R, Haq MM, Siddiqi KS. Synthesis and characterization of new 13- and 14-membered macrocycles and their transition metal complexes. Transition Met Chem 2003;28:948–53.
- Prasad RN, Mathur M. Cr^{III}, Fe^{III}, Co^{II}, Ni^{II}, Cu^{II} and Zn^{II} complexes of 26- and 28-membered tetraazamacrocycles. J Indian Chem Soc 2006;83:1208-13.
- Zeng Q, Sun J, Gou S, Zhou K, Fang J, Chen H. Synthesis and spectroscopic studies of dinuclear copper(II) complexes with new pendant-armed macrocyclic ligands. Transition Met Chem 1998;23:371–3.
- 23. Singh AK, Singh R, Saxena P. Macrocyclic metal complexes: synthesis and characterization of 14- and 16-membered tetraaza macrocyclic complexes of transition metals. Transition Met Chem 2004;29:867-9.
- Gupta LK, Chandra S. Physicochemical and biological characterization of transition metal complexes with a nitrogen donor tetra-dentate novel macrocyclic ligand. Transition Met Chem 2006;31:368–73.
- Mohamed AK, Islam KS, Hasan SS, Shakir M. Metal ion directed synthesis of 14-16 membered tetraimine macrocyclic complexes. Transition Met Chem 1999;24:198–201.
- 26. Lodeiro C, Basitida R, Bertolo E, Macias A, Rodriguez R. Synthesis and characterization of four novel $N_x O_y$ -Schiff base macrocyclic ligands and their metal complexes. Transition Met Chem 2003;28:388-94.
- Shakir M, Islam KS, Mohamed AK, Shagufta M, Hasan SS. Macrocyclic complexes of transition metals with divalent polyaza units. Transition Met Chem 1999;24:577-80.
- Shakir M, Varkey SP. A new synthetic route for the preparation of a new series of 14-22-membered tetraoxomacrocyclic tetraamines and their transition metal complexes. Polyhedron 1995;14:1117-27.
- 29. Aqra FMAM. New macrocyclic complexes containing amide, imine and secondary amine functions. Transition Met Chem 1999;24:337-9.
- Chandra S, Kumar R. Synthesis and spectral studies on mononuclear complexes of chromium(III) and manganese(II) with 12-membered tetradentate N₂O₂, N₂S₂ and N₄ donor macrocyclic ligand. Transition Met Chem 2004;29:269-75.
- 31. Wood JS. Stereochemical electronic structural aspects of five coordination. Prog Inorg Chem 1972;16:227.

1206 *D.P. Singh et al.*

- Singh DP and Rana VB. Binuclear chromium (III), manganese (III), iron (III) and cobalt (III) complexes bridged by diaminopyridine. Polyhedron. 1995; 14: 2901–2906.
- Lever ABP. Inorganic Electronic Spectroscopy. Amsterdam: Elsevier, 1984.
- Chohan ZH, Supuran CT. Structure and biological properties of first row d-transition metal complexes with N-substituted sulfonamides. J Enzyme Inhib Med Chem 2008;23:240–51.
- Chohan ZH. Metal-based sulfonamides: their preparation, characterization and *in-vitro* antibacterial, antifungal & cytotoxic properties. X-ray structure of 4-(2-hydroxybenzylidene) amino benzenesulfonamide. J Enzyme Inhib Med Chem 2008;23:120–30.
- 36. Chohan ZH, Shad HA. Structural elucidation and biological significance of 2-hydroxy-1-naphthaldehyde derived sulfonamides and

their first row d-transition metal chelates. J Enzyme Inhib Med Chem 2008;23:369–79.

- Chohan ZH. Synthesis of organometallic-based biologically active compounds: *in vitro* antibacterial, antifungal and cytotoxic properties of some sulfonamide incorporated ferrocences. J Enzyme Inhib Med Chem 2009;24:169–75..
- Chohan ZH, Supuran CT, Hadda TB, Nasim F, Khan KM. Metal based isatin-derived sulfonamides: their synthesis, characterization, coordination behavior and biological activity. J Enzyme Inhib Med Chem 2008 Sep 30:1. [Epub ahead of print]
- Singh K, Singh DP, Barwa MS, Tyagi P, Mirza Y. Antibacterial Co(II), Ni(II), Cu(II) and Zn(II) complexes of Schiff bases derived from fluorobenzaldehyde and triazoles. J Enzyme Inhib Med Chem 2006;21:557-62.