



World Trade Center-related asthma: clinical care essentials

Denise Harrison & Joan Reibman

To cite this article: Denise Harrison & Joan Reibman (2023) World Trade Center-related asthma: clinical care essentials, Archives of Environmental & Occupational Health, 78:4, 206-211, DOI: [10.1080/19338244.2023.2185191](https://doi.org/10.1080/19338244.2023.2185191)

To link to this article: <https://doi.org/10.1080/19338244.2023.2185191>



Published online: 20 Mar 2023.



Submit your article to this journal [↗](#)



Article views: 3232



View related articles [↗](#)



View Crossmark data [↗](#)



BRIEF REPORT



World Trade Center-related asthma: clinical care essentials

Denise Harrison^a and Joan Reibman^{a,b}

^aDepartment of Medicine, New York University Grossman School of Medicine, New York, NY, USA; ^bDepartment of Environmental Medicine, New York University Grossman School of Medicine, New York, NY, USA

ABSTRACT

Asthma is defined as a heterogeneous disease with respiratory symptoms (wheeze, shortness of breath, chest tightness and cough) that vary over time and intensity, and variable expiratory airflow limitation. Environmental and occupational exposures contribute to its causation. WTC-related or aggravated asthma is considered a World Trade Center (WTC) Health Program certifiable disease. Criteria include defined exposures to the WTC dust and fumes, the presence of symptoms, or aggravated symptoms that are present within 5 years after the last potential for WTC dust/fume exposures (the last 9/11 exposures occurred on July 31, 2002), and a WTC-provider diagnosis of asthma. Asthma is the 3rd most common non-cancer certification among WTC responders and survivors. In this review we provide evidence-based information on the evaluation, diagnosis, and treatment of patients with WTC-related or aggravated asthma and include peer-reviewed research findings in WTC-exposed populations.

ARTICLE HISTORY

Received 11 January 2023
Accepted 13 February 2023

KEYWORDS

World Trade Center; environment; adults; air pollution; asthma; dust; workers; occupational lung disease; particulate matter

Introduction

Acute and chronic exposures to World Trade Center (WTC) dust and fumes contribute to the causation and aggravation of asthma in WTC responders and survivors (local workers, residents, students, those passing by on 9/11). In this review we provide evidence-based information on the evaluation, diagnosis, and treatment of patients with potential for WTC-related or aggravated asthma and include peer-reviewed research findings in WTC-exposed populations. This is one of a series of papers for the diagnosis and treatment of persons with WTC-related diseases.¹

Relevant WTC exposures

The ~ 1.2 million tons of dust generated by the collapse of the WTC towers had a unique composition with potential for massive acute and chronic exposure dose. Most studies of the dust composition are based on settled dust, as local monitors for airborne particles were destroyed. Over 90% of the settled dust was >2.5 µm aerodynamic diameter. Although less than 1% of the settled dust was in the fine size range (<2.5 µm), the quantity of the dust resulted in significant airborne concentrations of fine dust.^{2–5} Coarse

particles in the WTC dust are more likely to deposit in the upper airways, however the massive quantity of dust resulted in appreciable penetration in the conducting as well as smaller airways, alveoli and interstitium of the lungs.^{2,6–8} Importantly, the settled WTC dust was highly alkaline.² Exposure dose analyses for WTC responders and survivors are complex, due to differences in acute exposures (i.e., arrival times, dust cloud exposure), as well as chronic exposures (e.g., prolonged work efforts, incompletely cleaned workplaces, homes or schools, resuspended dust, and fumes).

Asthma as a WTC-certified condition

The biologic plausibility for the development of asthma and the clinical and epidemiologic studies in responders and survivors resulted in the inclusion of asthma or aggravated asthma as a condition certified for health coverage by the WTC Health Program.^{9–20} WTC-related asthma is the third most common non-cancer certified condition in the CDC-NIOSH WTCHP.²¹ To certify a patient for asthma and asthma treatment, a WTC-3 application must be submitted for a patient whose WTC exposure was substantially likely to have caused, aggravated, or contributed to the diagnosis. To be certified, the WTC Health

Program member must meet 9/11 exposure requirements²² and lower respiratory symptoms must have had an onset within 5 years after the last date of the 9/11 exposure (the last 9/11 exposures occurred on July 31, 2002).²³

Asthma guidelines for diagnosis and management

Major guidelines have been published for asthma and severe asthma. For this best practices article, the selected guidelines are the NIH Expert Panel report, published in 2007 and updated in 2020^{24,25} and the Global Initiative for Asthma (GINA) Guidelines for the Management of Asthma and The GINA Guidelines for the Diagnosis and Treatment of Difficult-to-Treat and Severe Asthma^{26,27} These guidelines are enhanced by a position paper on irritant asthma.²⁸

Diagnosing asthma in the WTC population

Asthma is defined as a heterogeneous disease with respiratory symptoms (wheeze, shortness of breath, chest tightness and cough) that vary over time and intensity, and variable expiratory airflow limitation. Environmental and occupational exposures, including those from irritants, are potential causes. Symptoms often predate an asthma diagnosis. Spirometry with assessment of bronchodilator reversibility is an initial step in an evaluation for asthma, however, bronchodilator reversibility may not always be detected, and spirometry may not fully characterize the respiratory symptoms. Bronchoprovocation, peak flow monitoring, clinical treatment trials, or exercise studies can also be considered. Fixed airflow obstruction consistent with chronic obstructive pulmonary disease may also occur in severe or longstanding asthma. Although not included in the guidelines referenced above, small airway abnormalities have been documented in symptomatic obstructive airway diseases using pathologic studies, imaging, and physiologic techniques, including respiratory oscillometry.^{29–31}

Asthma symptoms associated with WTC dust/fume exposure have been well-described in responder and pediatric and adult survivor populations.^{9–20,32} Most have spirometry values within normal predicted values;^{11,12,33–36} airflow obstruction on spirometry is less common, but includes some who fit the definition of COPD.^{37–39} Because of symptoms consistent with asthma with normal spirometry, additional studies have been performed, which show bronchial hyperresponsiveness^{10,12,20,34,40} and possible small airway

involvement.^{35,41–44} Asthma presentation in WTC patients can be described as follows:

New onset WTC-related asthma

Many responders and survivors developed new asthma symptoms within days, months and years after their acute or chronic exposures to the WTC dust fumes. As such, most do not fit the definition of Reactive Airways Dysfunction (RADs), which requires acute onset of asthma.⁴⁵ The variable intensity of WTC exposure, the often delayed onset of symptoms, as well as the persistence of symptoms, is consistent with a diagnosis of “non-acute” irritant induced asthma (IIA).^{28,46–49}

WTC aggravated asthma

Some responders and survivors had preexisting asthma or childhood asthma with symptoms that worsened within a short period after their exposure to the WTC dust/fumes.

Asthma mechanisms and phenotypes

Asthma guidelines describe the presence of chronic airway inflammation^{26,50} and the phenotype has important implications for a personalized approach to management. Airway type 2 (T2) immune responses are mediated by eosinophils, mast cells, Th2 cells, group 2 innate lymphoid cells type 2, and/or IgE-producing B cells. Non-allergic stimuli (viruses, irritants, pollutants, tobacco smoke) can also activate T2 pathways.⁵⁰ In contrast, low T2 asthma is a diagnosis of exclusion, often associated with obesity/metabolic syndrome or environmental exposures.⁵¹ Fixed airflow obstruction consistent with COPD may occur with either.

Heterogeneity of WTC-related or aggravated asthma may be due to differences in WTC or other exposures (tobacco use, other occupations), social determinants of health, co-morbidities, as well as genetic underpinnings. Perennial and/or seasonal or mold allergen sensitization was noted in 17–40% of responders.^{37,52,53} Elevated levels of IL-5 and IL-4 are reported in firefighters, and blood neutrophil or eosinophil counts are associated with the decline or reduced FEV₁.^{54,55} However, many WTC-exposed patients with asthma symptoms have neither elevated total IgE, blood eosinophils nor FeNO.^{37–39,54–56}

Treatment of WTC-related or aggravated asthma

Current goals of asthma management are to reduce symptoms and minimize risk.^{24–27} Guidelines

reinforce the need to reduce modifiable risks such as tobacco use, and to use non-pharmacologic approaches (asthma education, trigger recognition and avoidance, pulmonary rehabilitation, weight management) and pharmacologic therapies with a patient-provider partnership. Assessment of symptom control including impairment (symptom frequency and functional status) and risk (exacerbations, reduced lung function) is stressed. Assessment of asthma severity is recommended only after treatment and incorporates the level of medication needed to control the disease. A written asthma action plan is also encouraged. Pharmacologic approaches include the early introduction of inhaled corticosteroids (ICS) either with a short acting β_2 agonist or as needed ICS-formoterol, even for mild asthma. Failing control, step-up therapy with an ICS and an inhaled long acting β_2 agonist, including single agent ICS-formoterol as maintenance and reliever therapy, and eventually anti-muscarinic agents are suggested.²⁶ Controlling co-morbidities including chronic rhinosinusitis with/without nasal polyposis, gastroesophageal reflux disease, elevated BMI, psychiatric disease, and metabolic syndrome is of great importance.²⁶ Patients with moderate-severe uncontrolled asthma who fail to attain control with these interventions should be referred to specialists for further diagnostic evaluation, phenotyping and consideration of add-on biologic agents targeting IgE, IL-5, IL-4Ra or thymic stromal lymphopoietin.²⁷

Few studies have examined pharmacologic efficacy in the WTC affected populations. One study suggests that early initiation of treatment is important for the reduction of long term effects.⁵⁷ A small study in survivors with persistent uncontrolled symptoms despite prescribed treatment, failed to show an improvement in symptom control despite adherence to high dose ICS-LABA.⁵⁸ Elevated BMI and metabolic syndrome are associated with persistence and worsening of symptoms, making these potential targets for treatment.^{16–18,47,59–62}

Program coverage

Asthma diagnostic services are covered by the WTC Health Program. For diagnostic coverage, the patient must be enrolled in the WTC Health Program. Treatment coverage requires that the enrolled member's asthma be certified. Asthma treatment is fully covered by the WTC Health Program for responders, and for survivors the treatment benefit is coordinated with other health insurance. To receive certification, a WTC Health Program-affiliated Clinical Center of

Excellence (CCE) or the Nationwide Provider Network (NPN) must submit a WTC-3 form (<https://www.cdc.gov/wtc/pdfs/Appendix-WTC3.pdf>). Among other things, on the WTC-3 form, the CCE/NPN physician must attest that 9/11 exposures were substantially likely to have been a significant factor in aggravating, contributing to, or causing the enrolled WTC member's asthma. To receive asthma certification, the member must also satisfy 9/11 exposure (NIOSH 2015) and maximum time interval requirements (NIOSH 2017).

Conclusion

WTC-related or aggravated asthma is a common outcome of WTC exposures in responders and survivors. The disease is a heterogeneous disease with differing phenotypes and associated co-morbidities. These characteristics require personalized evaluation and approaches consistent with high quality asthma guidelines.

Disclosure statement

No potential conflict of interest was reported by the authors.

Disclaimer

The contents of this article are the sole responsibility of the authors and do not necessarily represent the official views of, nor an endorsement, by the National Institute for Occupational Safety and Health (NIOSH), the Centers for Disease Control and Prevention of the U.S. Department of Health and Human Services (CDC/HHS), or the U.S. Government.

Institutional review board (IRB) review

This activity did not involve human subjects and therefore did not require IRB review.

Funding

This work was funded in part by the National Institute for Occupational Safety and Health (NIOSH), Centers for Disease Control and Prevention of the U.S. Department of Health and Human Services (CDC/HHS) (CDC/NIOSH: 200-2017-93327 [JR], 200-2017-93427 [JR] and 200-2017-93430 [DH]).

References

1. Calvert GM, Anderson K, Cochran J, et al. The World Trade Center Health Program: an introduction to best practices. *Arch Environ Occup Health*. 2022;1–7. doi: [10.1080/19338244.2022.2156975.36533439](https://doi.org/10.1080/19338244.2022.2156975.36533439)

2. Lippmann M, Cohen MD, Chen LC. Health effects of World Trade Center (WTC) Dust: An unprecedented disaster's inadequate risk management. *Crit Rev Toxicol*. 2015;45(6):492–530. doi:10.3109/10408444.2015.1044601.
3. Liou PJ, Weisel CP, Millette JR, et al. Characterization of the dust/smoke aerosol that settled east of the World Trade Center (WTC) in lower Manhattan after the collapse of the WTC 11 September 2001. *Environ Health Perspect*. 2002;110(7):703–714. doi:10.1289/ehp.02110703.
4. McGee JK, Chen LC, Cohen MD, et al. Chemical analysis of World Trade Center fine particulate matter for use in toxicologic assessment. *Environ Health Perspect*. 2003;111(7):972–980. doi:10.1289/ehp.5930.
5. Yiin LM, Millette JR, Vette A, et al. Comparisons of the dust/smoke particulate that settled inside the surrounding buildings and outside on the streets of southern New York City after the collapse of the World Trade Center, September 11, 2001. *J Air Waste Manag Assoc*. 2004;54(5):515–528. doi:10.1080/10473289.2004.10470935.
6. Rom WN, Weiden M, Garcia R, et al. Acute eosinophilic pneumonia in a New York City firefighter exposed to World Trade Center dust. *Am J Respir Crit Care Med*. 2002;166(6):797–800. doi:10.1164/rccm.200206-576OC.
7. Fireman EM, Lerman Y, Ganor E, et al. Induced sputum assessment in New York City firefighters exposed to World Trade Center dust. *Environ Health Perspect*. 2004;112(15):1564–1569. doi:10.1289/ehp.7233.
8. Caplan-Shaw CE, Yee H, Rogers L, et al. Lung pathologic findings in a local residential and working community exposed to World Trade Center dust, gas, and fumes. *J Occup Environ Med*. 2011;53(9):981–991. doi:10.1097/JOM.0b013e31822fff60.
9. Centers for Disease C, Prevention. Self-reported increase in asthma severity after the September 11 attacks on the World Trade Center–Manhattan, New York, 2001. *MMWR Morb Mortal Wkly Rep*. 2002; 51(35):781–784.
10. Prezant DJ, Weiden M, Banauch GI, et al. Cough and bronchial responsiveness in firefighters at the World Trade Center site. *N Engl J Med*. 2002;347(11):806–815. doi:10.1056/NEJMoa021300.
11. Banauch GI, Alleyne D, Sanchez R, et al. Persistent hyperreactivity and reactive airway dysfunction in firefighters at the World Trade Center. *Am J Respir Crit Care Med*. 2003;168(1):54–62. doi:10.1164/rccm.200211-1329OC.
12. Reibman J, Lin S, Hwang SA, et al. The World Trade Center residents' respiratory health study: new-onset respiratory symptoms and pulmonary function. *Environ Health Perspect*. 2005;113(4):406–411. doi:10.1289/ehp.7375.
13. Lin S, Reibman J, Bowers JA, et al. Upper respiratory symptoms and other health effects among residents living near the World Trade Center site after September 11, 2001. *Am J Epidemiol*. 2005;162(6): 499–507. doi:10.1093/aje/kwi233.
14. Banauch GI, Hall C, Weiden M, et al. Pulmonary function after exposure to the World Trade Center collapse in the New York City Fire Department. *Am J Respir Crit Care Med*. 2006;174(3):312–319. doi:10.1164/rccm.200511-1736OC.
15. Wheeler K, McKelvey W, Thorpe L, et al. Asthma diagnosed after 11 September 2001 among rescue and recovery workers: findings from the World Trade Center Health Registry. *Environ Health Perspect*. 2007; 115(11):1584–1590. doi:10.1289/ehp.10248.
16. Farfel M, DiGrande L, Brackbill R, et al. An overview of 9/11 experiences and respiratory and mental health conditions among World Trade Center Health Registry enrollees. *J Urban Health*. 2008;85(6):880–909.
17. Brackbill RM, Hadler JL, DiGrande L, et al. Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. *JAMA*. 2009;302(5):502–516.
18. Friedman SM, Farfel MR, Maslow C, et al. Risk factors for and consequences of persistent lower respiratory symptoms among World Trade Center Health Registrants 10 years after the disaster. *Occup Environ Med*. 2016;73(10):676–684.
19. Kim H, Herbert R, Landrigan P, et al. Increased rates of asthma among World Trade Center disaster responders. *Am J Ind Med*. 2012;55(1):44–53.
20. Aldrich TK, Weakley J, Dhar S, et al. Bronchial reactivity and lung function after world trade center exposure. *Chest*. 2016;150(6):1333–1340.
21. <https://www.cdc.gov/wtc/ata glance.html#top10Conditions2022>. 2022
22. Centers for Disease Control and Prevention NIOSH. *Policy and Procedures for Certification of Physician Determinations for Aerodigestive and Cancer Health Conditions*. Washington, DC: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. 2015 <https://www.cdc.gov/wtc/pdfs/policies/WTCHPPPCertPhysDetFINAL20Feb2015-508.pdf>. 2015.
23. Centers for Disease Control and Prevention NIOSH. *Time Intervals for New-Onset Aerodigestive Disorders*. Washington DC: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. https://www.cdc.gov/wtc/pdfs/policies/WTCHPP_PP_Time_Intervals_New_Onset_AeroDig_Disorders_30_August_2017-508.pdf. 2017.
24. National Heart L, and Blood Institute. Guidelines for the Diagnosis and Management of Asthma (EPR-3). 2007.
25. National Heart L, and Blood Institute. Asthma Management Guidelines: Focused Updates 2020.
26. Asthma. Gf. Global Strategy for Asthma Management and Prevention, 2021. 2021.
27. Asthma Gf. Diagnosis and Management of Difficult-to-Treat and Severe Asthma. 2021.
28. Vandenplas O, Wiszniewska M, Raulf M, et al. EAACI position paper: irritant-induced asthma. *Allergy*. 2014;69(9):1141–1153.
29. Postma DS, Brightling C, Baldi S, et al. Exploring the relevance and extent of small airways dysfunction in asthma (ATLANTIS): baseline data from a prospective cohort study. *Lancet Respir Med*. 2019;7(5):402–416.

30. Usmani OS, Han MK, Kaminsky DA, et al. Seven pillars of small airways disease in asthma and COPD: Supporting opportunities for novel therapies. *Chest*. 2021;160(1):114–134.
31. Kraft M, Richardson M, Hallmark B, et al. The role of small airway dysfunction in asthma control and exacerbations: a longitudinal, observational analysis using data from the ATLANTIS study. *Lancet Respir Med*. 2022;10(7):661–668.
32. Antao VC, Pallos LL, Graham SL, et al. 9/11 residential exposures: The impact of world trade center dust on respiratory outcomes of lower Manhattan residents. *Int J Environ Res Public Health*. 2019;16(5):798.
33. Banauch GI, Brantly M, Izbicki G, et al. Accelerated spirometric decline in New York City firefighters with alpha(1)-antitrypsin deficiency. *Chest*. 2010;138(5):1116–1124.
34. Aldrich TK, Gustave J, Hall CB, et al. Lung function in rescue workers at the World Trade Center after 7 years. *N Engl J Med*. 2010;362(14):1263–1272.
35. Friedman SM, Maslow CB, Reibman J, et al. Case-control study of lung function in World Trade Center Health Registry area residents and workers. *Am J Respir Crit Care Med*. 2011;184(5):582–589.
36. Liu M, Qian M, Cheng Q, et al. Longitudinal spirometry among patients in a treatment program for community members with World Trade Center-related illness. *J Occup Environ Med*. 2012;54(10):1208–1213.
37. Singh A, Liu C, Putman B, et al. Predictors of asthma/COPD Overlap in FDNY firefighters with world trade center dust exposure: A longitudinal study. *Chest*. 2018;154(6):1301–1310.
38. Haghighi A, Cone JE, Li J, de la Hoz RE. Asthma-COPD overlap in World Trade Center Health Registry enrollees, 2015–2016. *J Asthma*. 2021;58(11):1415–1423.
39. Baba RY, Zhang Y, Shao Y, et al. COPD in smoking and non-smoking community members exposed to the world trade center dust and fumes. *Int J Environ Res Public Health*. 2022;19(7):4249.
40. Berger KI, Kalish S, Shao Y, et al. Isolated small airway reactivity during bronchoprovocation as a mechanism for respiratory symptoms in WTC dust-exposed community members. *Am J Ind Med*. 2016;59(9):767–776.
41. Skloot G, Goldman M, Fischler D, et al. Respiratory symptoms and physiologic assessment of ironworkers at the World Trade Center disaster site. *Chest*. 2004;125(4):1248–1255.
42. Oppenheimer BW, Goldring RM, Herberg ME, et al. Distal airway function in symptomatic subjects with normal spirometry following World Trade Center dust exposure. *Chest*. 2007;132(4):1275–1282.
43. Berger KI, Turetz M, Liu M, et al. Oscillometry complements spirometry in evaluation of subjects following toxic inhalation. *ERJ Open Res*. 2015;1(2):00043–2015.
44. Pradhan D, Xu N, Reibman J, et al. Bronchodilator response predicts longitudinal improvement in small airway function in world trade center dust exposed community members. *Int J Environ Res Public Health*. 2019;16(8):1421.
45. Brooks SM, Weiss MA, Bernstein IL. Reactive airways dysfunction syndrome (RADS). Persistent asthma syndrome after high level irritant exposures. *Chest*. 1985;88(3):376–384.
46. de la Hoz RE. Occupational lower airway disease in relation to World Trade Center inhalation exposure. *Curr Opin Allergy Clin Immunol*. 2011;11(2):97–102.
47. Wisnivesky JP, Teitelbaum SL, Todd AC, et al. Persistence of multiple illnesses in World Trade Center rescue and recovery workers: a cohort study. *Lancet*. 2011;378(9794):888–897.
48. Caplan-Shaw C, Kazeros A, Pradhan D, et al. Improvement in severe lower respiratory symptoms and small airway function in World Trade Center dust exposed community members. *Am J Ind Med*. 2016;59(9):777–787.
49. Cleven KL, Rosenzvit C, Nolan A, et al. Twenty-year reflection on the impact of world trade center exposure on pulmonary outcomes in fire department of the City of New York (FDNY) rescue and recovery workers. *Lung*. 2021;199(6):569–578.
50. Israel E, Reddel HK. Severe and difficult-to-treat asthma in adults. *N Engl J Med*. 2017;377(10):965–976.
51. Fitzpatrick AM, Chipps BE, Holguin F, Woodruff PG. T2-“Low” Asthma: Overview and management strategies. *J Allergy Clin Immunol Pract*. 2020;8(2):452–463.
52. de la Hoz RE, Shohet MR, Wisnivesky JP, Bienenfeld LA, Afilaka AA, Herbert R. Atopy and upper and lower airway disease among former World Trade Center workers and volunteers. *J Occup Environ Med*. 2009;51(9):992–995.
53. Rojano B, West E, Ferdermann E, et al. Allergen sensitization and asthma outcomes among world trade center rescue and recovery workers. *Int J Environ Res Public Health*. 2019;16(5):737.
54. Zeig-Owens R, Singh A, Aldrich TK, et al. Blood leukocyte concentrations, FEV1 decline, and airflow limitation. A 15-year longitudinal study of world trade center-exposed firefighters. *Ann Am Thorac Soc*. 2018;15(2):173–183.
55. Weiden MD, Singh A, Goldfarb DG, et al. Serum Th-2 cytokines and FEV1 decline in WTC-exposed firefighters: A 19-year longitudinal study. *Am J Ind Med*. 2021;64(10):845–852.
56. Kazeros A, Maa MT, Patrawalla P, et al. Elevated peripheral eosinophils are associated with new-onset and persistent wheeze and airflow obstruction in world trade center-exposed individuals. *J Asthma*. 2013;50(1):25–32.
57. Goldfarb DG, Putman B, Lahousse L, et al. Lung function decline before and after treatment of World Trade Center associated obstructive airways disease with inhaled corticosteroids and long-acting beta agonists. *Am J Ind Med*. 2021;64(10):853–860.
58. Reibman J, Caplan-Shaw C, Wu Y, et al. Characterization of persistent uncontrolled asthma symptoms in community members exposed to world trade center dust and fumes. *Int J Environ Res Public Health*. 2020;17(18):6645.
59. Friedman SM, Farfel MR, Maslow CB, Cone JE, Brackbill RM, Stellman SD. Comorbid persistent lower respiratory symptoms and posttraumatic stress

- disorder 5-6 years post-9/11 in responders enrolled in the World Trade Center Health Registry. *Am J Ind Med.* 2013;56(11):1251–1261.
60. Moline JM, McLaughlin MA, Sawit ST, et al. The prevalence of metabolic syndrome among law enforcement officers who responded to the 9/11 World Trade Center attacks. *Am J Ind Med.* 2016; 59(9):752–760.
61. Kwon S, Crowley G, Mikhail M, et al. Metabolic syndrome biomarkers of world trade center airway hyperreactivity: A 16-year prospective cohort study. *Int J Environ Res Public Health.* 2019;16(9):1486.
62. Brite J, Friedman S, de la Hoz RE, Reibman J, Cone J. Mental health, long-term medication adherence, and the control of asthma symptoms among persons exposed to the WTC 9/11 disaster. *J Asthma.* 2019:1–10.