



Gut Microbes



ISSN: (Print) (Online) Journal homepage: informahealthcare.com/journals/kgmi20

Gut non-bacterial microbiota contributing to alcohol-associated liver disease

Wenkang Gao, Yixin Zhu, Jin Ye & Huikuan Chu

To cite this article: Wenkang Gao, Yixin Zhu, Jin Ye & Huikuan Chu (2021) Gut non-bacterial microbiota contributing to alcohol-associated liver disease, Gut Microbes, 13:1, 1984122, DOI: 10.1080/19490976.2021.1984122

To link to this article: https://doi.org/10.1080/19490976.2021.1984122

9	© 2021 The Author(s). Published with license by Taylor & Francis Group, LLC.
+	View supplementary material $oldsymbol{\mathcal{C}}$
	Published online: 28 Oct 2021.
	Submit your article to this journal $oldsymbol{\mathcal{C}}$
hh	Article views: 3551
Q ¹	View related articles 🗗
CrossMark	View Crossmark data ☑
2	Citing articles: 9 View citing articles 🗹



REVIEW 3 OPEN A



Gut non-bacterial microbiota contributing to alcohol-associated liver disease

Wenkang Gao^a, Yixin Zhu^b, Jin Ye 10a, and Huikuan Chu 10a

^aDivision of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; ^bDepartment of Medicine, University of California San Diego, La Jolla, California, USA

ABSTRACT

Intestinal microbiota, dominated by bacteria, plays an important role in the occurrence and the development of alcohol-associated liver disease (ALD), which is one of the most common liver diseases around the world. With sufficient studies focusing on the gut bacterial community, chronic alcohol consumption is now known as a key factor that alters the composition of gut bacterial community, increases intestinal permeability, causes intestinal dysfunction, induces bacterial translocation, and exacerbates the process of ALD via gut-liver axis. However, gut non-bacterial communities including fungi, viruses, and archaea, which may also participate in the disease, has received little attention relative to the gut bacterial community. This paper will systematically collect the latest literatures reporting non-bacterial communities in mammalian health and disease, and review their mechanisms in promoting the development of ALD including CLEC7A pathway, Candidalysin (a peptide toxin secreted by *Candida albicans*), metabolites, and other chemical substances secreted or regulated by gut commensal mycobiome, virome, and archaeome, hoping to bring novel insights on our current knowledge of ALD.

inflammation and fibrosis Cirrhosis Cirrhosis Cirrhosis CIECTA microbiota and its products captilated ISC hepatocyte injury structure and Getth microbiota and its products

ARTICLE HISTORY

Received 20 July 2021 Revised 9 September 2021 Accepted 14 September 2021

KEYWORDS

Alcohol-associated liver disease; gut-liver axis; gut barrier; liver sinusoid endothelial cells; mycobiome; CLEC7A; virome; archaeome; immune response

CONTACT Huikuan Chu 2012XH0827@hust.edu.cn Division of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1277 Jiefang Avenue, Wuhan, 430022, China

supplemental data for this article can be accessed on the publisher's website.

Introduction

Alcohol-associated liver disease (ALD) is a common disease caused by alcohol use disorder (AUD), ranging from asymptomatic liver steatosis to alcoholassociated hepatitis (AH), cirrhosis, and potentially hepatocellular carcinoma (HCC). ALD is the leading indication for liver transplantation in the United States.^{1,2} Globally, about 2 million people die of liver diseases each year. In Western Europe, onethird of liver cirrhosis can be attributed to alcohol.³ Moreover, it has been estimated that 60-80% of liver-related deaths can be attributed to alcohol consumption.⁴ Currently, the pathogenetic mechanisms have not been fully elucidated, but they might be related to oxidative stress, acetaldehyde-induced toxicity, cytokine, and chemokine-induced inflammation.⁵ There is no effective therapeutic method for ALD till now except for liver transplantation. A great number of studies have reported that gut microbiota has an intimate relationship with ALD, which provides broader insights and opportunities for understanding and treating this disease. 6-14 However, most studies put emphasis upon the interactions between gut bacteria and ALD, while the roles of non-bacterial communities remain unclear. Hence, this review mainly focuses on addressing the relationships between ALD and the non-bacterial communities (fungi, viruses, and archaea) in the gut, and the possible pathogenetic mechanisms mediated by those microorganisms.

The gut-liver axis

The term gut-liver axis was first described by *Volta* et al in 1987, which pointed out the interaction between intestine and liver. During the embryonic stage, both the liver and intestine originate from the same foregut, and the two organs are anatomically connected by the portal vein system after reaching their maturity. About 70 ~ 80% of the blood from the portal vein flows to the liver, transmitting a variety of signals generated by dietary, genetic, and environmental factors. (Figure 1) As a virtual metabolic organ, the gut-liver axis achieves close-knit functional collaboration and forms a sophisticated network structure through substance metabolism, immune regulation, and interaction with the neuroendocrine system.

Normally, the healthy intestinal barrier is composed of the mucus barrier, the intestinal epithelial barrier, and the gut vascular barrier (GVB), 18-21 which together protects the body against external insults (Figure 1(a)). In the colon, the mucus layer usually consists of a microbiota-colonized outer layer and an almost sterile inner layer. These inner and outer layers work together to provide the needed nutrients for microorganisms survival unique niches for microorganisms inhabitation. 22,23 Below the mucus barrier is the epithelial barrier, which is a single layer formed by the epithelial cells including enterocytes, goblet cells, tuft cells, and enterochromaffin cells.²⁴ The neighboring epithelial cells also remain in close contact through tight junctions. 25-27 Epithelial barrier's main functions are regulating the interactions between gut microbiota and the host immune system, and selectively absorbing water, electrolytes, and nutrients. 28,29 The GVB identified in 2015 is comprised of vascular endothelial cells, pericytes, and glial cells, which prevents the invasion of microbiota. 30,31 As the deepest protective layer and the last defense line of the gut barrier, once the GVB is injured, the intestinal pathogens will go into the blood stream and reach other organs (Figure 1(b)). Hence, in order to fight against exogenous substances in the gut lumen, these three layers summarized above constitute the body's first line of defense.

Once the intestinal barrier collapses, the liver becomes the first organ to encounter intestinal products, which makes it susceptible to pathological changes.^{32,33} Liver sinusoidal endothelial cells (LSECs), a major member of the hepatic barrier, provide a second line of defense against gut-derived antigens and inflammatory factors.³⁴ LSECs are highly fenestrated cells (Figure 1(c)). The diameters of transcellular pores on LSECs are 50 ~ 200 nm and these LSECs usually act as a dynamic filter under physiological conditions. 35,36 For example, small molecules, such as chylomicron remnants, plasma proteins, and lipoproteins, can cross fenestrae and reach the space of Disse for uptake and utilization by hepatocytes and HSCs.³⁷ Furthermore, LSECs can remove recycled waste products and toxicants, including viruses,^{38,39} bacteriophages,⁴⁰ lipopolysaccharides (LPS),⁴¹ and immune complexes.⁴² LSECs can also




Figure 1. The double line of defense in the gut-liver axis. a. The first line of defense is the gut barrier, which is composed of the mucus barrier, the intestinal epithelial barrier, and the GVB. The mucus layer consists of a microbiota-colonized outer layer and an almost sterile inner layer. The epithelial barrier below the mucus layer is formed by epithelial cells. The deepest protective layer is the GVB. Under physiological circumstances, the healthy intestine barrier allows the absorption of nutrients while prevents the penetration of most toxic and harmful substances. b. Once it is damaged, intestinal microorganisms and their metabolites will move into the liver through the portal vein. c. LSECs are highly fenestrated cells, which can remove recycled waste products by endocytosis and degradation to protect liver against microorganisms and toxicants. They can also induce hepatic immune tolerance and keep the quiescence of HSCs and Kupffer cells. d. Injured LSECs become capillarized and the main characteristics of capillarization are formation of basement membrane, reduction of fenestrae and increased expression of pro-inflammatory mediators. They recruit immune cells and activate HSCs and Kupffer cells to cause liver inflammation. GVB: gut vascular barrier; LSECs: liver sinusoid endothelial cells. HSCs: hepatic stellate cells.

maintain the quiescent state of hepatic stellate cells, induce hepatic immune tolerance and exert an antiinflammatory effect. 43,44 Due to the existence of the second line of defense, the sole injury of the intestinal barrier is unlikely to lead to significant liver damage, which has been confirmed experimentally in the previous studies. 32,33

However, when LSECs are impaired by intestinal pathogenic factors, all the above functions are repressed. LSECs become capillarized⁴⁵ and their phenotype is transformed into a pro-inflammatory pattern, with the formation of the basement membrane, a significant reduction of fenestrae, and increased expression of pro-inflammatory mediators such as tumor necrosis factor α (TNF- α), macrophage inflammatory protein 1 alpha (MIP-1α), monocyte chemoattractant protein-1 (MCP-1), and chemokine (C-C motif) ligand 5 (CCL5).⁴⁶ At the same time, LSECs recruit immune cells and activate HSCs and Kupffer cells to induce liver inflammation.⁴⁷ (Figure 1(d)). Inflammatory

factors and metabolites secreted by the injured liver will further aggravate dysfunction of the intestinal barrier, creating a vicious circle eventually. In the future, understanding the mechanism of barrier damage and repair, finding damage markers that are easily detectable, developing drugs to repair the barrier are important steps for the treatment of liver and intestine-related diseases, and achieving the transformation from basic research to clinical implementations.

A complete gut-liver axis relies on not only the intact intestinal barrier and normal liver function but also the healthy gut microbiota. Gut microbiota is composed of 500 to 1000 different species belonging to more than 70 genera. 48,49 The number of bacteria residing in the human body is close to the number of human cells. 50,51 Not merely can they participate in digestion, absorption and metabolism of food but can also influence the intestinal structure and immune function both directly and indirectly. Therefore, clarifying the changes and

roles of gut microbiota in ALD is of great importance for us to understand its pathogenesis and to develop effective treatment strategies. Whereas an immense number of reviews have covered gut bacteria in detail, 52-56 thus we center our discussion on the gut mycobiome, virome, and archaeome in human and animal models.

Gut mycobiome

Fungi are ubiquitous in the natural environment and are important parts of the earth's ecosystem.⁵⁷ The fungal community is considered as a fundamental component of the human microbiota, coexisting and interacting with other microorganisms in the gut.⁵⁸ However, they are usually neglected in studies owing to their relatively low abundance in the human body. For instance, there have been almost 100 times more research reports on microbiota than on mycobiota from 2008 to 2018.⁵⁹ With the increase of related research data and the development of methodology, scientists begin to realize its indispensable role in metabolism, immune regulation, and pathophysiology of the host.

Composition and colonization

Fungi occupy a relatively small proportion of the human microbiota. Fungi comprise approximately 0.03% of the fecal microbiome. However, this does not mean that it is meaningless to study fungi. Conversely, some scientists contended that previous efforts might underestimate the proportion of fungi in gut microbiota owing to limited annotated reference sequences. Besides, fungal cells are generally larger than bacterial cells, indicating that fungi contain much more biomass and metabolites, which cannot be reflected by simple genome-counting. Thus, more data and further studies are required for accurately estimating the amount of living gut fungi and completing the annotated fungal genome.

The gut fungi in adults are mainly composed of three phyla, *Ascomycetes*, *Basidiomycetes*, and *Zygomycetes*, ⁶³ which are regulated by many factors such as environment, ⁶⁴ diet, ^{65–67} host immunity, ⁶⁸ etc. Both human and animal studies have confirmed that the composition of the intestinal mycobiome is dynamic over time and much more mutable than the component of gut bacteria. ^{69,70} A previous study indicated that intestinal fungi




Figure 2. The mechanisms of ALD induced by the gut mycobiota. **a.** Ethanol abuse leads to disordered gut mycobiota. *Epicoccum, Galactomyces*, and *Debaryomyces* decrease, while *Candida* increases significantly in the intestine. 1,3-β-glucan, a main component of fungal cell walls, and Candidalysin, a fungal metabolite, translocate into the systemic circulation through the injured intestinal barrier and reach the liver first. **b.** 1,3-β-glucan binds to dectin-1 of Kupffer cells. Src kinases phosphorylate tyrosine residue of dectin-1 and recruit Syk, which can activate the CARD9/BCL10/MALT1 complex. Then NF-κB will be activated by the complex and produce pro-IL-1β. Syk also promotes the secretion of caspase-1 through the activated NLRP3/ASC/pro-caspase-1 complex. Caspase-1 then cleaves pro-IL-1β into mature IL-1β, mediating liver inflammation and damage. Syk, spleen tyrosine kinase; CARD9, caspase recruitment domain 9; BCL10, B-cell lymphoma 10; MALT1, mucosa-associated lymphoid tissue lymphoma-translocation gene 1; NF-κB, nuclear factor kappa B; NLRP3, NOD-like receptor family pyrin domain containing 3; ASC, apoptosis-associated speck-like protein containing a CARD.

have colonized in human 10 days after birth. 71 But recently, fungi were found in the first-pass meconium, suggesting that colonization of gut fungi may exist earlier than expected. 72 In the gut of 10-day to 3-month-old infants, the fungal species that had the highest abundance are Debaryomyces hansenii and Rhodotorula mucilaginosa, while S. cerevisiae becomes the most abundant fungal species in the gut of babies 1-2 years after birth.⁷¹ From that timepoint and on, the gut fungi develops and gets closer and closer to the intestinal fungal components of the adult. Therefore, given the variability of fungal components, some scholars advocate that fungi do not colonize the gastrointestinal tract in healthy adults, and they may originate from the fungi already present in the mouth or diet, and persistent fungal colonization may be a symptom of disease.⁶⁷ In their study, when a healthy adult volunteer increased the frequency of cleaning teeth, the abundance of C. albicans in stool was lowered 10-fold to 100-fold, suggesting that the oral cavity may be the primary source of C. albicans detected in the stool of healthy people.⁶⁷ Indeed, existing research rarely consider the influence of oral and dietary fungi, especially in the studies of exploring changes in the gut fungi compositions. It is difficult to determine whether differences in stool fungi are due to increased transient fungi or due to real intestinal colonization. Overall, the colonization and composition of intestinal fungi are still in their infancy. In the future, further studies with a larger sample size should be done in this field, and sequencing methods with higher sensitivity and accuracy need to be invented to detect intestinal fungi at various time points of embryonic development.

Intestinal fungi and alcohol-associated liver disease

The studies on the correlation between gut mycobiota and ALD are limited and mostly focus on the changes in fungal composition and exploration of potential pathogenic mechanisms. Previous studies reported a decrease in the diversity of mycobiota in the feces of alcohol consumers, with a significant overgrowth of Candida, and a decrease in Epicoccum, unclassified fungi, Galactomyces, and Debaryomyces. 73,74 Similarly, recent research reported mycobiota dysbiosis with an overgrowth

of Candida as well. 75-77 Although there were differences in the severity of alcohol-related liver disease, there were no significant differences in the intestinal mycobiota among patients with nonprogressive alcohol-associated liver disease, alcohol-associated hepatitis, and alcoholic cirrhosis.⁷³

As for the role of gut fungi in the pathogenesis of ALD, there remain two main pathways at present: one is dependent on the C-type lectin domain family 7 member A (CLEC7A) pathway of Kupffer cells in liver and the other is independent of that pathway (Figure 2(a)).

CLEC7A (also called dectin-1) is a pattern recognition receptor primarily located on the surface of Kupffer cells and macrophages in the liver.⁷³ CLEC7A is capable of recognizing 1,3-β-glucan,⁷⁸ which is widely found in fungal cell walls, and plays an important role in resisting fungi and fungal products.(Figure 2(b))⁷⁹ Upon ligand binding, tyrosine residue of dectin-1 is phosphorylated by Src kinases, which creates a docking site for spleen tyrosine kinase (Syk), then Syk activates the caspase recruitment domain 9 (CARD9)/B-cell lymphoma 10 (BCL10)/mucosa-associated lymphoid tissue lymphoma-translocation gene 1 (MALT1) complex, which leads to the activation of nuclear factor kappa B (NF-κB) and promotes cytokines expression including pro-interleukin-1β (pro-IL-1β).^{80,81} Furthermore, Syk can activate NOD-like receptor family pyrin domain containing 3 (NLRP3) inflammasome, which usually forms a complex with apoptosis-associated speck-like protein containing a CARD (ASC) and pro-caspase-1. ASC can activate pro-caspase-1 into caspase-1, which will cleave pro-IL-1β to produce mature IL-1β and lead to liver damage by activating downstream cascade.^{82,83} An animal study reported that the development of ALD required IL-1β which was activated via a caspase-1-dependent process, and that IL-1β signaling was essential for alcohol-induced inflammation, steatosis, liver damage, and fibrosis.84 A subsequent study⁷³ also revealed that chronic alcohol intake could increase the number of fungi in mice intestine, and that overgrowth of fungi could produce more fungal products such as 1,3-βglucan, which can be translocated into the liver through already dysfunctional enteric tight junctions, bind to the Dectin-1 of Kupffer cells, and finally lead to the secretion of IL-1β to promote

liver inflammation. However, one study indicated that water-insoluble polysaccharides (WIP), a 1,3- β -glucan from *Wolfporia cocos*, alleviated the hepatic inflammatory injury and steatosis effectively in mice with alcoholic hepatic steatosis (AHS), which suggested the protective effect of 1,3- β -glucan in liver diseases but the mechanism needs to be further studied.

Another fungus-related mechanism is mainly associated with its metabolites. As an opportunistic pathogen in the intestine, Candida albicans has attracted extensive attention because of its overgrowth in ALD patients.⁷³ Recent studies have proven that Candida albicans could secrete a type of peptide toxin called Candidalysin, which directly damages epithelial membranes and activates host immune responses.86 One previous study found an interesting phenomenon that the number of Candidalysin positive C. albicans increased in patients with AH and the expression of the extent of cell elongation 1 (ECE1) gene encoding Candidalysin also increased significantly. 87 In addition, clinical studies suggested that Candidalysin was positively associated with the severity and mortality of AH.87 With regard to the underlying pathogenic mechanisms of Candidalysin, however, it does not affect the intestinal permeability nor interacts with the glucan receptor CLEC7A, but it increases the expression of IL-1\beta, CXCL1, and CXCL2 in mice liver, which contributes to recruit immune cells and induce hepatocyte death. In brief, the specific molecular mechanisms need further research.

A recent research also found the contribution of fungi-induced metabolite – prostaglandin E₂ (PGE₂) to the development of ALD.⁸⁵ They observed an overwhelming increase of the commensal fungus *Meyerozyma guilliermondii* and hepatic PGE₂ in mice with alcoholic hepatic steatosis (AHS). By introducing antifungal agents, the level of liver PGE₂ was significantly reduced. Interestingly, it is still unclear whether PGE₂ is produced directly by the gut fungi or indirectly by the fungi-stimulated liver. To sum up, the concrete mechanisms of gut fungi-induced PGE₂ in the development of ALD awaits further study.

In addition, the host immune response plays a key role in the process of ALD as well. A recent study⁷⁴ revealed a novel finding that the level of

serum anti-Saccharomyces cerevisiae antibodies (ASCA) was significantly higher in patients with AH compared with alcohol use disorder and non-alcoholic controls. As a marker of the host immune response to fungus and fungal products, ASCA was positively associated with mortality in AH patients.

In summary, it is not difficult to find that a few potential noninvasive indicators, such as the level of serum ASCA, serum Candidalysin, serum 1,3- β -glucan, and the expression of ECE1 in stool, can help us determine the progress or prognosis of ALD. In addition, inhibiting the increase of alcohol-related gut fungi, especially some specific antifungal drugs against harmful mycobiome, may be a promising therapeutic method for ALD, but its clinical application has yet to be confirmed in large-scale cohort studies. Also, further compelling investigations should explore the causal relationship between the specific fungus strain isolated from the gut and the pathogenesis of ALD.

Probiotic fungi

In 2014, the International Scientific Association of Probiotics and Prebiotics (ISAPP) defined probiotics as "live microorganisms that, when administered in adequate amounts, confer a health benefit on the host". 88 There are many types of probiotics, such as Bifidobacterium bifidus, Lactobacillus, and Saccharomyces boulardii, of which S. boulardii is probably the most commonly used probiotic fungi. Animal studies have confirmed that S. boulardii could change gut microbiota and attenuate liver injury, inflammation, 89-91 fibrosis, 92 potentially indicating its protective and therapeutic role in liver diseases. A recent study revealed that most strains of Saccharomyces and non-Saccharomyces yeasts evaluated in their work are safe microorganisms, and those could be regarded as a valid alternative to the widely available probiotic yeast S. boulardii. 93 However, some side effects of S. boulardii, mainly presented as fungemia, were reported in clinical application, which are usually found in the elderly, immunosuppressed, and patients with broad-spectrum antibiotics. 94-97 Furthermore, macrofungi with large sporocarps or fruiting bodies, also known as mushrooms, particularly showed their medicinal properties.⁹⁸ Some types of macrofungi including



Agaricus bisporus 99-101 and Pleurotus ostreatus 102 also possess hepato-protective activity. In conclusion, supplementing probiotic fungi to prevent or treat liver diseases can be expected in the future.

Gut virome

Composition and distribution

Viruses have their own important roles in the intestine. As the most numerous biological entities in nature, 103,104 viruses are composed of endogenous retroviruses, eukaryotic viruses, and bacteriophages. Most viruses in the human body reside in the gastrointestinal tract.

The gut viruses, even called virus fingerprints, are unique to each person and mainly consist of bacteriophages. 105 However, identifying viral sequences in large and mixed gut microbiota is extremely challenging because viruses lack a universal viral marker. 106 Thus, an overwhelming number of gut bacteriophages remain uncultured and unclassified till now, and their specific hosts and infection strategies have not been clarified either. Scientists also named gut viruses "the dark matter" in the intestine, which showed that relevant research on them were far from enough. 107-109

Fortunately, the Gut Virome Database (GVD)¹¹⁰ and the Gut Phage Database¹¹¹ have been established and are continuously updating, which will advance research of gut virome considerably. According to the GVD, 110 97.7% of viral populations are phages, 2.1% are eukaryotic viruses, and 0.1% are archaeal viruses. About 10¹⁵ bacteriophages settle in a healthy human intestine, which is 10 times the number of symbiotic bacteria. 112 The main phages come from the order Caudovirales including Siphoviridae, Myoviridae, and Podoviridae. 113,114 Some phages are highly specific to certain bacterial strains, while others have broader ranges of host cells. 115 After infection of host by phages, highly virulent phages often cause cell lysis (lytic cycle) while mild phages either lyse the host cell or keep the host cell alive and reproduce normally (lysogenic cycle). 116,117

In summary, large-scale efforts are still required urgently in order to culture and catalog gut phages, which is much like the process of building collections and genome databases of bacterial strains in the human microbiome. 118

Is it pathogens?

Phages are regarded as potential human pathogens. 119 This was probably first suggested with the genome of lambdoid phages that could encode Shiga toxins (Stx), 120 which poses a severe and lethal threat to human health. Moreover, animal models revealed that phages may act as pathogens by affecting the gut bacterial communities and increasing the gut permeability. 121 Besides, An increasing number of evidence indicates that human diseases, such as inflammatory bowel disease, 109,122 diabetes, 123 acquired immune deficiency syndrome, 124 and colorectal cancer, 125 are closely related to the gut viruses. Nevertheless, it is difficult to determine whether the changes in the gut virome are a cause or a result of those diseases. And the boundary between 'normal' and pathogenic virome is blurred because the same virus can be either a symbiont or a pathogen depending on the conditions such as the health status and development stage of the host. On the whole, the current knowledge of symbiotic viruses lags far behind that of pathogenic viruses. 126

Gut virome and alcohol-associated liver disease

The exact role of gut virome in the etiology of ALD is still unclear, as not only the complex pathogenesis of ALD but also our fragmented understanding of the gut viruses and limited relevant studies. In 2020, scientists systematically described an intestinal virome signature in AH patients for the first time. They observed an increased viral diversity in the stools of patients with ALD. In AH patients, Escherichia-, Enterobacteria-, and Enterococcus phages were overrepresented and mammalian viruses such as Parvoviridae and Herpesviridae significantly increased. Staphylococcus phages and Herpesviridae were associated with severity and mortality of ALD. 127128 Above findings were consistent with previous research. In fecal samples, most detected Herpesviridae such as herpesvirus-6128,129 and herpesvirus-8^{130,131} are associated with the severity of ALD, and they could be classified into Epstein-Barr virus (EBV). EBV infection often causes liver inflammation, but the pathogenesis remains unknown. 129 A retrospective study of patients with liver cirrhosis found that EBV-positive patients had

higher Child-Pugh scores, more severe liver injury and a higher rate of chronic acute liver failure, ¹³² which suggested the reactivation of EBV might contribute to the development of alcoholic hepatitis.

Gut virome may regulate the pathogenesis of ALD through multiple interactions with symbiotic bacteria. ¹³³ Phages, as the main component of the gut virome, are able to mediate bacterial cells lysis and regulate the abundance of bacteria. A recent study confirmed that the interaction between bacteria and phages in patients with cirrhosis and hepatic encephalopathy centered on *Streptococcus* species. ¹³⁴ Moreover, gut bacteria have access to additional genomes (such as antibiotic-resistance genes or bacterial virulence factors) transferred by phages, which will adjust bacterial virulence and adaptability and modify the composition of intestine microbiota. ¹³⁵

Furthermore, gut virome can interact with the host immune system directly. The intestinal immune system is mainly comprised of intestinal intraepithelial lymphocytes (IELs), lamina propria lymphocytes (LPLs), and Peyer's patches (PPs), ¹³⁶ which could be important in the pathogenesis of ALD. Among them, commensal viruses can be recognized by retinoic acid-inducible gene 1 (RIG-1) signaling in antigenpresenting cells (APCs), and RIG-1 signaling can promote IL-15 production, prevent inflammation and tissue damage as well as maintain the homeostasis of IELs. 137 Moreover, scientists found that the immune cells including CD4+ and CD8+ T cells increased in the germ-free mice treated with bacteriophages, and the increased abundance of bacteriophages can exacerbate intestinal colitis through tolllike receptor 9 (TLR9) and interferon-gamma (IFNy), indicating that phages can alter mucosal immunity to impact mammalian health. 138 But, the mechanism of the phage recognition by the mucosal immune system needs more studies in animal models. Moreover, some studies on pathogenic enteroviruses, such as norovirus and rotavirus, have revealed several signaling pathways of intestinal recognition of viral nucleic acids. 139,140 Thus, the gut virome either commensal or pathogenic viruses is a hot topic in host immune response.

To sum up, gut virome may promote the progression of ALD through interventions with the symbiotic bacteria and immune system¹⁴¹. However, it is still unclear for patients with ALD what specific impacts of increased gut viruses are on the other

microorganisms or the host body. We do not know yet whether liver inflammation or damage is caused by the virus itself, or its metabolites, or changes of other microbes induced by the altered gut virome. The studies of gut virome are just starting and more studies are needed to help us understand the correlations between the gut virus, the health of the host and liver diseases.

Phage therapy of ALD

Recent animal studies creatively revealed the importance of phages in the treatment of ALD. According to a study in 2019, 142 Enterococcus faecalis was significantly increased in fecal samples and the cytolysin-positive (cytolytic) E. faecalis correlated with the severity and mortality of AH patients. Then, researchers screened out bacteriophages that can lyse the cytolytic E. faecalis to treat mice transplanted with the gut microbiota of AH patients, which significantly reduced the levels of cytolysin in mice liver and attenuated alcoholinduced liver inflammation.

Editing intestinal microbes to treat or improve diseases related to microbial dysbiosis seems to be a quite promising direction. After all, antibiotics are not suitable for precise editing due to their broad spectrum, instead, some bacteriophages are highly specific to bacteria, suggesting a unique advantage of manipulating microbiota. 143,144 Certainly, the clinical application of phage therapy still needs the support of large-scale human data. In addition, the limitation of phage therapy may be due to its high specificity. Different patients need to deploy different phages in order to better efficacy, so its mass production-like antibiotics may be unrealistic in some extent. In the future, it is necessary to construct a huge phage bank, which will efficiently help us screen out concrete phages and make a personalized treatment within a short period of time.

Gut archaeome

Composition and distribution

Archaea were originally discovered and isolated from ecosystems with extreme conditions, including environments with high temperature, strong acid or base, and high ion concentration. However, with the continuous advancing of detection techniques, archaea had also been found in some mild environments such as the ocean ecosystems. 145-149 Archaea are similar to bacteria in terms of shape, size, and genetic information expression including DNA replication, RNA transcription, and protein synthesis. Beside these similarities, there are also some obvious differences between archaea and bacteria. For instance, the walls peptidoglycans, 150 and their cell membranes are composed of L-glycerol-ether/isoprenoid lipids, which are more stable and rigid than bacterial.¹⁵¹ Moreover, due to its special metabolic patterns, archaea can use sunlight, inorganic or organic substances as energy sources. 152

The most abundant archaea in the human gastrointestinal tract are Methanobacteriales Methanomassiliicoccales. 153 The former, mainly conof Methanobrevibacter smithii sisting Methanosphaera stadtmanae, 154 is identified as a keystone species that has a noticeable impact on the composition and function of other gut microbiome. 155,156 The species of archaea are usually affected by various factors such as diet, environment, age, genetics, etc., and they are also related to nonarchaea members of the host microbiota. Interestingly, the abundance of M. smithii was recently found to be stable over time, even after major changes in diet. 157 In addition, the archaea present highly specific adaptability in the gastrointestinal tract, and they often evolve unique features such as modifications of cell surface (adhesin-like proteins (ALPs), glycans, bile salt hydrolases, and biofilm formation), which are not found in wide archaea, to escape the host-defense mechanism. 158

Generally speaking, as an important part of the gastrointestinal microbiota of humans and animals, the role of archaea may be far more underestimated owing to the methodological shortcomings. 159 For instance, most of the so-called "universal" 16S rRNA primers fail to picture the diversity of archaeal signatures and thus are unable to detect certain archaeal lineages in specific sample types. 160161 The archaea research is still in its infancy and the academics of microbiology lack relevant knowledge as well.

Archaea and human diseases

Whether archaea are pathogens or not has always been a controversial topic. 158 Some studies verified the change of archaeome in certain human diseases. For example, M. smithii was found in urinary tract infections. 161 M. oralis was reported in periodontitis, and its quantity increased as the illness severity increased, but it was no longer present after healing, which highlighted the close association of *M. oralis* with the inflamed site. 162 The abundance of M. stadtmanae increased three-fold in inflammatory bowel disease (IBD) patients when compared to healthy individuals, indicating this archaeon might be involved in pathologic process within the human gut. 163

Metabolites produced by archaea such as methane can also affect the health status of the host. In dog models, methane can weaken gastrointestinal motility. 164 In a population with irritable bowel syndrome (IBS), the subjects with the constipation-dominant disease (IBS-C) were shown to have a higher proportion of methane producers than individuals with the diarrhea-dominant disease (IBS-D).165

Furthermore, people have also found the interaction between archaea and the human immune system. When stimulating the immune system with M. stadtmanae, 154 high levels of proinflammatory cytokines including interleukin and interferon were released. 166,167 Only recently was it demonstrated that RNA from M. stadtmanae was a potent immune stimulator, and toll-like receptor 7 (TLR7) and TLR8 were identified as the involved pattern recognition receptors. Moreover, this molecular interaction induced TLR8-dependent NLRP3 inflammasome activation. 168 The proinflammatory ability of archaea also varies from species to species. For example, M. stadtmanae is capable of inducing a stronger immune response than both M. smithii and M. luminyensis. 166,169 There are some speculations attempting to explain the mild response induced by M. smithii stating that M. smithii has the capacity to produce glycans which is similar to those found in the gut, 170 and thus may help M. smithii escape from the host immune system. In conclusion, understanding the molecular

mechanisms of how archaea induce inflammation in the body is thus an important step in uncovering how such diseases might develop.

The impact of ALD on the composition of archaea is rarely reported. According to a recent research, the proportion of archaea among the healthy control group, the alcohol use disorder group, and the alcoholic hepatitis group was 0.6%. 0.3%, and 0.0% respectively, which indicated that the abundance of archaea was reduced as the disease progresses. However, it is not clear which archaea have modified, nor is it clear whether these changes are a result of or a cause for the disease. But considering the proinflammatory ability, it is worthwhile to study the interspecies communication between archaea and other gut microbes, the potential role of archaeal metabolites, and the relationship of archaea and chronic liver diseases.

Conclusion

Although we have generally devoted significant efforts on studying the gut bacteriota, it is worth noting that non-bacterial communities also have potential functions. All these microbes will interact with the host and also with each other, in combination or alone, to influence the health and disease of the host. Eventually, we will unveil the role of non-bacterial microbiota in human with the remarkable development of technologies for the detection of fungi, viruses, and archaea. Furthermore, human microbiota investigations have now reached a critical inflection point. In the future, we need to move rapidly from mere description or correlation to causation, and ultimately to translation. ¹⁷²

To accomplish such a great mission, much work remains ahead of us. The first is the improvement of methodology. At present, there are three kinds of methods commonly used in studying intestinal microbiota: culture-dependent methods, culture-independent methods, and high-throughput sequencing (HTS). There are also several excellent literatures evaluating these methods in gut mycobiome, ¹⁷³ virome, ¹⁷⁴ and archaeome. ¹⁶⁰ Nevertheless, most research tools still have some defects. Culture methods are time-consuming and also susceptible to environmental factors. Scientists

might misestimate the abundance of certain gut microorganisms owing to the incompleteness of annotated gene databases or improper primers. In addition, the amount of relic DNA in human gut remains unknown currently, and a few fungi DNA can even be detected after years of organism decomposition. Thus, we need to be more prudent in face of the results of DNAbased sequencing. The study of gut nonbacterial microbiota will be more efficient through establishing a uniform, accurate, and highly operable detection system for nonbacteriota. And ensuring that the microbiome from different regions, races, and species can be compared or referenced is highly expected in future work.

Secondly, it is necessary to expand the genomics, transcriptomics, and metabolomics data of fungi, viruses, and archaea in the existing database, so that future research can accurately and quickly identify the corresponding microbiota in multiple dimensions. Moreover, the study of microbiome cannot simply stop at the macroscopic appearance of the gut community composition changes, it should be conducted from bacterial species to specific strains, from cells to subcellular organs, from metabolites to molecular regulation. Study in such detail will provide a fresh and fundamental perspective for manipulating and editing the gut microbiota logically and efficiently in the future. Finally, the commensal microbiota in the intestine is an emerging and promising field that has aroused great interest from scientists, physicians and patients. With a deeper understanding and transformation of the gut-liver axis, the investments may be rewarded and the management of ALD patients will also be improved.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This project was supported by National Natural Science Foundation of China (grant number 81770582 to JY and 82000561 to HKC).

ORCID

Jin Ye (b) http://orcid.org/0000-0002-1521-1746 Huikuan Chu http://orcid.org/0000-0001-9177-2804

Authors' contributions

WKG contributed to the conception and drafted the manuscript. YXZ polished the language and grammar and revised the manuscript. JY checked and improved the manuscript. HKC contributed to the conception and critically revised the manuscript. All authors read and approved the final version of the manuscript.

References

- 1. Cholankeril G, Wong RJ, Hu M, Perumpail RB, Yoo ER, Puri P, Younossi ZM, Harrison SA, Ahmed A. 2017. Liver transplantation for nonalcoholic steatohepatitis in the US: temporal trends and outcomes. Dig Dis Sci. 62 doi:10.1007/s10620-017-4684-x. (10):2915-2922. PMID: 28744836
- 2. Lee BP, Vittinghoff E, Dodge JL, Cullaro G, Terrault NA. 2019. National trends and long-term outcomes of liver transplant for alcohol-Associated liver disease in the United States. JAMA Intern Med. 179 (3):340-348.doi:10.1001/jamainternmed.2018.6536. PMID: 30667468
- 3. Mokdad AA, Lopez AD, Shahraz S, Lozano R, Mokdad AH, Stanaway J, Murray CJ, Naghavi M. 2014. Liver cirrhosis mortality in 187 countries between 1980 and 2010: a systematic analysis. BMC Med. 12 doi:10.1186/s12916-014-0145-y. (1):145.25242656
- 4. Sheron N. 2016. Alcohol and liver disease in Europe-Simple measures have the potential to prevent tens of thousands of premature deaths. J Hepatol. 64 (4):957–967. doi:10.1016/j.jhep.2015.11.006. PMID: 26592352
- 5. Seitz HK, Bataller R, Cortez-Pinto H, Gao B, Gual A, Lackner C, Mathurin P, Mueller S, Szabo G, Tsukamoto H, et al. 2018. Alcoholic liver disease. Nat Rev Dis Primers. 4(1):16. PMID: 30115921. doi:10.1038/ s41572-018-0014-7.
- 6. Rivera Ca, Bradford BU, Seabra V, Thurman RG. 1998. Role of endotoxin in the hypermetabolic state after acute ethanol exposure. Am J Physiol. 275:G1252-8. PMID: 9843760. doi:10.1152/ajpgi.1998.275.6.G1252.
- 7. Yan AW, Fouts DE, Brandl J, Starkel P, Torralba M, Schott E, Tsukamoto H, E. Nelson K, A. Brenner D, Schnabl B, et al. 2011. Enteric dysbiosis associated with a mouse model of alcoholic liver disease. Hepatology. 53 (1):96-105. PMID: 21254165. doi:10.1002/hep.24018.
- 8. Bajaj JS, Heuman DM, Hylemon PB, Sanyal AJ, White MB, Monteith P, Noble NA, Unser AB, Daita K, Fisher AR, et al. 2014. Altered profile of human gut

- microbiome is associated with cirrhosis and its complications. J Hepatol. 60(5):940-947. PMID: 24374295. doi:10.1016/j.jhep.2013.12.019.
- 9. Leclercq S, Matamoros S, Cani PD, Neyrinck AM, Jamar F, Starkel P, Windey K, Tremaroli V, Bäckhed F, Verbeke K, et al. 2014. Intestinal permeability, gut-bacterial dysbiosis, and behavioral markers of alcohol-dependence severity. Proc Natl Acad Sci U S A. 111(42):E4485-93. PMID: 25288760. doi:10.1073/ pnas.1415174111.
- 10. Chen P, Miyamoto Y, Mazagova M, Lee KC, Eckmann L, Schnabl B. 2015. Microbiota protects mice against acute alcohol-induced liver injury. Alcohol Clin Exp Res. 39 (12):2313-2323. doi:10.1111/acer.12900. PMID: 26556636
- 11. Llopis M, Cassard AM, Wrzosek L, Boschat L, Bruneau A, Ferrere G, Puchois V, Martin JC, Lepage P, Le Roy T, et al. 2016. Intestinal microbiota contributes to individual susceptibility to alcoholic liver disease. Gut.65(5):830-839. PMID: 26642859. doi:10.1136/gutjnl-2015-310585.
- 12. Wang L, Fouts DE, Starkel P, Hartmann P, Chen P, Llorente C, DePew J, Moncera K, Ho S, Brenner D, et al. 2016. Intestinal REG3 lectins protect against alcoholic steatohepatitis by reducing Mucosa-Associated microbiota and preventing bacterial translocation. Cell Host Microbe. 19(2):227-239. PMID: 26867181. doi:10.1016/j.chom.2016.01.003.
- 13. Llorente C, Jepsen P, Inamine T, Wang L, Bluemel S, Wang HJ, Loomba R, Bajaj JS, Schubert ML, Sikaroodi M, et al. 2017. Gastric acid suppression promotes alcoholic liver disease by inducing overgrowth of intestinal enterococcus. Nat Commun. 8(1):837. PMID: 29038503. doi:10.1038/s41467-017-00796-x.
- 14. Smirnova E, Puri P, Muthiah MD, Daitya K, Brown R, Chalasani N, Liangpunsakul S, Shah VH, Gelow K, Siddiqui MS, et al. 2020. Fecal microbiome distinguishes alcohol consumption from alcoholic hepatitis but does not discriminate disease severity. Hepatology. 72(1):271-286. PMID: 32056227. doi:10.1002/hep.31178.
- 15. Volta U, Bonazzi C, Bianchi FB, Baldoni AM, Zoli M, Pisi E. 1987. IgA antibodies to dietary antigens in liver cirrhosis. Ric Clin Lab. 17:235-242. PMID: 3671996. doi:10.1007/BF02912537.
- 16. Abdel-Misih SR, Bloomston M. 2010. Liver anatomy. Surg Clin North Am. 90(4):643-653. doi:10.1016/j. suc.2010.04.017. PMID: 20637938
- 17. Ger R. 1989. Surgical anatomy of the liver. Surg Clin North Am. 69(2):179-192. doi:10.1016/s0039-6109(16) 44780-8. PMID: 2928899
- 18. Albillos A, de Gottardi A, Rescigno M. 2020. The gut-liver axis in liver disease: pathophysiological basis for therapy. J Hepatol. 72(3):558-577. doi:10.1016/j. jhep.2019.10.003. PMID: 31622696
- 19. Brescia P, Rescigno M. 2021. The gut vascular barrier: a new player in the gut-liver-brain axis. Trends Mol 27(9):844-855. doi:10.1016/j. Med. molmed.2021.06.007. PMID: 34229973

- Ghosh S, Whitley CS, Haribabu B, Jala VR. 2021.
 Regulation of intestinal barrier function by microbial metabolites. Cell Mol Gastroenterol Hepatol. 11 (5):1463–1482. doi:10.1016/j.jcmgh.2021.02.007.
 PMID: 33610769
- 21. Chopyk DM, Grakoui A. 2020. Contribution of the Intestinal microbiome and gut barrier to hepatic disorders. Gastroenterology. 159(3):849–863. doi:10.1053/j.gastro.2020.04.077. PMID: 32569766
- 22. Ermund A, Schutte A, Johansson ME, Gustafsson JK, Hansson GC. 2013. Studies of mucus in mouse stomach, small intestine, and colon. I. Gastrointestinal mucus layers have different properties depending on location as well as over the Peyer's patches. Am J Physiol Gastrointest Liver Physiol. 305(5):G341–7. doi:10.1152/ajpgi.00046.2013. PMID: 23832518
- 23. Li H, Limenitakis JP, Fuhrer T, Geuking MB, Lawson MA, Wyss M, Brugiroux S, Keller I, Macpherson JA, Rupp S, et al. 2015. The outer mucus layer hosts a distinct intestinal microbial niche. Nat Commun. 6(1):8292. PMID: 26392213. doi:10.1038/ncomms9292.
- 24. Kurashima Y, Kiyono H. 2017. Mucosal ecological network of epithelium and immune cells for gut homeostasis and tissue healing. Annu Rev Immunol. 35 (1):119–147. doi:10.1146/annurev-immunol-051116-052424. PMID: 28125357
- 25. Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA. 1986. Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. J Cell Biol. 103(3):755–766. doi:10.1083/jcb.103.3.755. PMID: 3528172
- Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S, Tsukita S. 1993. Occludin: a novel integral membrane protein localizing at tight junctions. J Cell Biol. 123(6):1777–1788. PMID: 8276896. doi:10.1083/ jcb.123.6.1777.
- 27. Furuse M, Fujita K, Hiiragi T, Fujimoto K, Tsukita S. 1998. Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. J Cell Biol. 141(7):1539–1550. doi:10.1083/jcb.141.7.1539. PMID: 9647647
- 28. Odenwald MA, Turner JR. 2017. The intestinal epithelial barrier: a therapeutic target? Nat Rev Gastroenterol Hepatol. 14(1):9–21. doi:10.1038/nrgastro.2016.169. PMID: 27848962
- Rescigno M. 2011. The intestinal epithelial barrier in the control of homeostasis and immunity. Trends Immunol. 32(6):256–264. doi:10.1016/j.it.2011.04.003. PMID: 21565554
- 30. Spadoni I, Zagato E, Bertocchi A, Paolinelli R, Hot E, Di Sabatino A, Caprioli F, Bottiglieri L, Oldani A, Viale G, et al. 2015. A gut-vascular barrier controls the systemic dissemination of bacteria. Science. 350(6262):830–834. PMID: 26564856. doi:10.1126/science.aad0135.

- 31. Bouziat R, Jabri B. 2015. IMMUNOLOGY. Breaching the gut-vascular barrier. Science. 350(6262):742–743. doi:10.1126/science.aad6768. PMID: 26564835
- Gabele E, Dostert K, Hofmann C, Wiest R, Scholmerich J, Hellerbrand C, Obermeier F. 2011. DSS induced colitis increases portal LPS levels and enhances hepatic inflammation and fibrogenesis in experimental NASH. J Hepatol. 55(6):1391–1399. doi:10.1016/j. jhep.2011.02.035. PMID: 21703208
- El Kasmi KC, Anderson AL, Devereaux MW, Fillon SA, Harris JK, Lovell MA, Finegold MJ, Sokol RJ. 2012. Toll-like receptor 4-dependent Kupffer cell activation and liver injury in a novel mouse model of parenteral nutrition and intestinal injury. Hepatology. 55 (5):1518–1528. doi:10.1002/hep.25500. PMID: 22120983
- 34. Wang Y, Liu Y. 2021. Gut-liver-axis: barrier function of liver sinusoidal endothelial cell (LSEC). J Gastroenterol Hepatol. PMID: 33811372 10.1111/jgh.15512
- 35. Wisse E. 1970. An electron microscopic study of the fenestrated endothelial lining of rat liver sinusoids. J Ultrastruct Res. 31(1–2):125–150. doi:10.1016/s0022-5320(70)90150-4. PMID: 5442603
- 36. Monkemoller V, Oie C, Hubner W, Huser T, McCourt P. 2015. Multimodal super-resolution optical microscopy visualizes the close connection between membrane and the cytoskeleton in liver sinusoidal endothelial cell fenestrations. Sci Rep. 5(1):16279. doi:10.1038/srep16279. PMID: 26549018
- Poisson J, Lemoinne S, Boulanger C, Durand F, Moreau R, Valla D, Rautou P-E. 2017. Liver sinusoidal endothelial cells: physiology and role in liver diseases.
 J Hepatol. 66(1):212–227. doi:10.1016/j. jhep.2016.07.009. PMID: 27423426
- 38. Ganesan LP, Mohanty S, Kim J, Clark KR, Robinson JM, Anderson CL, Desrosiers RC. 2011. Rapid and efficient clearance of blood-borne virus by liver sinusoidal endothelium. PLoS Pathog. 7(9):e1002281. doi:10.1371/journal.ppat.1002281. PMID: 21980295
- Simon-Santamaria J, Rinaldo CH, Kardas P, Li R, Malovic I, Elvevold K,McCourt P, Smedsrod B, Hirsch HH, Sorensen KK. 2014. Efficient uptake of blood-borne BK and JC polyomavirus-like particles in endothelial cells of liver sinusoids and renal vasa recta. PLoS One. 9(11):e111762. PMID: 25375646. doi:10.1371/journal.pone.0111762.
- Oie CI, Wolfson DL, Yasunori T, Dumitriu G, Sorensen KK, McCourt PA, Ahluwalia BS, Smedsrod B. 2020. Liver sinusoidal endothelial cells contribute to the uptake and degradation of entero bacterial viruses. Sci Rep. 10(1):898. PMID: 31965000. doi:10.1038/s41598-020-57652-0.
- 41. Yao Z, Mates JM, Cheplowitz AM, Hammer LP, Maiseyeu A, Phillips GS, Wewers MD, Rajaram MVS, Robinson JM, Anderson CL, et al. 2016. Blood-Borne lipopolysaccharide is rapidly eliminated by liver

- sinusoidal endothelial cells via high-density lipoprotein. J Immunol. 197(6):2390-2399. PMID: 27534554. doi:10.4049/jimmunol.1600702.
- 42. Ganesan LP, Kim J, Wu Y, Mohanty S, Phillips GS, Birmingham DJ, Robinson JM, Anderson CL. 2012. FcgammaRIIb on liver sinusoidal endothelium clears immune complexes. J Immunol. (10):4981-4988. PMID: 23053513. doi:10.4049/ jimmunol.1202017.
- 43. Deleve LD, Wang X, Guo Y. 2008. Sinusoidal endothelial cells prevent rat stellate cell activation and promote reversion to quiescence. Hepatology. 48(3):920-930. doi:10.1002/hep.22351. PMID: 18613151
- 44. Xie G, Wang X, Wang L, Wang L, Atkinson RD, Kanel GC, Gaarde WA, DeLeve LD. 2012. Role of differentiation of liver sinusoidal endothelial cells in progression and regression of hepatic fibrosis in rats. Gastroenterology. 142(4):918-27e6. doi:10.1053/j.gastro.2011.12.017. PMID: 22178212
- 45. Schaffner F, Poper H. 1963. Capillarization of hepatic sinusoids in man. Gastroenterology. 44(3):239-242. PMID: 13976646. doi:10.1016/S0016-5085(63)80130-4.
- 46. Connolly MK, Bedrosian AS, Malhotra A, Henning JR, Ibrahim J, Vera V, Cieza-Rubio NE, Hassan BU, Pachter HL, Cohen S, et al. 2010. In hepatic fibrosis, liver sinusoidal endothelial cells acquire enhanced immunogenicity. J Immunol. 185(4):920-930. PMID: 20639479. doi:10.1002/hep.22351.
- 47. Miyao M, Kotani H, Ishida T, Kawai C, Manabe S, Abiru H, Tamaki K. 2015. Pivotal role of liver sinusoidal endothelial cells in NAFLD/NASH progression. Lab Invest. 95(10):1130-1144. doi:10.1038/labinvest.2015.95. PMID: 26214582
- 48. Human Microbiome PC. 2012. Structure, function and diversity of the healthy human microbiome. Nature. 486:207-214. PMID: 22699609. doi:10.1038/nature11234.
- 49. Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, Pollard KS, Sakharova E, Parks DH, Hugenholtz P, et al. 2021. A unified catalog of 204,938 reference genomes from the human gut microbiome. Nat Biotechnol. 39(1):105-114. PMID: 32690973. doi:10.1038/s41587-020-0603-3.
- 50. Sender R, Fuchs S, Milo R. 2016. Are we really vastly outnumbered? revisiting the ratio of bacterial to host cells in humans. Cell. 164(3):337-340. doi:10.1016/j. cell.2016.01.013. PMID: 26824647
- 51. Sender R, Fuchs S, Milo R. 2016. Revised estimates for the number of human and bacteria cells in the body. **PLoS** Biol. 14(8):e1002533. doi:10.1371/journal. pbio.1002533. PMID: 27541692
- 52. Bajaj JS. 2019. Alcohol, liver disease and the gut microbiota. Nat Rev Gastroenterol Hepatol. 16(4):235-246. doi:10.1038/ s41575-018-0099-1. PMID: 30643227
- 53. Lang S, Schnabl B. 2020. Microbiota and fatty liver disease-the known, the unknown, and the future. Cell Host Microbe. 28(2):233-244. doi:10.1016/j. chom.2020.07.007. PMID: 32791115

- 54. Szabo G. 2015. Gut-liver axis in alcoholic liver disease. Gastroenterology. 148(1):30-36. doi:10.1053/j.gastro.2014.10.042. PMID: 25447847
- 55. Betrapally NS, Gillevet PM, Bajaj JS. 2016. Changes in the intestinal microbiome and alcoholic and nonalcoholic liver diseases: causes or effects? Gastroenterology. 150(8):1745-55e3. doi:10.1053/j.gastro.2016.02.073. PMID: 26948887
- 56. Adolph TE, Grander C, Moschen AR, Tilg H. 2018. Liver-microbiome axis in health and disease. Trends Immunol. 39(9):712-723. doi:10.1016/j.it.2018.05.002. PMID: 29843959
- 57. Peay KG, Kennedy PG, Talbot JM. 2016. Dimensions of biodiversity in the earth mycobiome. Nat Rev Microbiol. 14(7):434-447. doi:10.1038/nrmicro.2016.59. PMID: 27296482
- 58. Seed PC. 2014. The human mycobiome. Cold Spring Harb Perspect Med. 5(5):a019810. doi:10.1101/cshperspect.a019810. PMID: 25384764
- 59. Forbes JD, Bernstein CN, Tremlett H, Van Domselaar G, Knox NC. 2018. A fungal world: could the gut mycobiome be involved in neurological disease? Front Microbiol. 9:3249. PMID: 30687254. doi:10.3389/ fmicb.2018.03249.
- 60. Ott SJ, Kuhbacher T, Musfeldt M, Rosenstiel P, Hellmig S, Rehman A, Drews O, Weichert W, Timmis KN, Schreiber S, et al. 2008. Fungi and inflammatory bowel diseases: alterations of composition and diversity. Scand J Gastroenterol. 43(7):831–841. PMID: 18584522. doi:10.1080/00365520801935434.
- 61. Hillman ET, Lu H, Yao T, Nakatsu CH. 2017. Microbial ecology along the gastrointestinal tract. Microbes Environ. 32(4):300-313. doi:10.1264/jsme2.ME17017. PMID: 29129876
- 62. Underhill DM, Iliev ID. 2014. The mycobiota: interactions between commensal fungi and the host immune system. Nat Rev Immunol. 14(6):405-416. doi:10.1038/ nri3684. PMID: 24854590
- 63. Hoffmann C, Dollive S, Grunberg S, Chen J, Li H, Wu GD, Lewis JD, Bushman FD. 2013. Archaea and fungi of the human gut microbiome: correlations with diet and bacterial residents. PLoS One. 8(6):e66019. doi:10.1371/journal.pone.0066019. PMID: 23799070
- 64. Yeung F, Chen YH, Lin JD, Leung JM, McCauley C, Devlin JC, Hansen C, Cronkite A, Stephens Z, Drake-Dunn C, et al. 2020. Altered immunity of laboratory mice in the natural environment is associated with fungal colonization. Cell Host Microbe. 27(5):809-22e6. PMID: 32209432. doi:10.1016/j.chom.2020.02.015.
- 65. Ukhanova M, Wang X, Baer DJ, Novotny JA, Fredborg M, Mai V. 2014. Effects of almond and pistachio consumption on gut microbiota composition in a randomised cross-over human feeding study. Br J Nutr. 111(12):2146-2152. doi:10.1017/S0007114514000385. PMID: 24642201
- 66. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, et al. 2014. Diet rapidly

- - reproducibly alters the human gut microbiome. Nature. 505(7484):559-563. PMID: 24336217. doi:10.1038/ nature12820.
- 67. Auchtung TA, Fofanova TY, Stewart CJ, Nash AK, Wong MC, Gesell JR, Auchtung JM, Ajami NJ, Petrosino JF. 2018. Investigating colonization of the healthy adult gastrointestinal tract by fungi. mSphere. 3(2). PMID: 29600282. doi:10.1128/mSphere.00092-18.
- 68. Iliev ID, Funari VA, Taylor KD, Nguyen Q, Reyes CN, Strom SP, Brown J, Becker CA, Fleshner PR, Dubinsky M, et al. 2012. Interactions between commensal fungi and the C-type lectin receptor Dectin-1 influence colitis. Science. 336(6086):1314-1317. PMID: 22674328. doi:10.1126/science.1221789.
- 69. Hallen-Adams HE, Kachman SD, Kim J, Legge RM, Martinez I. 2015. Fungi inhabiting the healthy human gastrointestinal tract: a diverse and dynamic community. Fungal Ecol. 15:9-17. PMID: WOS: 000356109500002. doi:10.1016/j.funeco.2015.01.006.
- 70. Dollive S, Chen YY, Grunberg S, Bittinger K, Hoffmann C, Vandivier L, Cuff C, Lewis JD, Wu GD, Bushman FD, et al. 2013. Fungi of the murine gut: episodic variation and proliferation during antibiotic treatment. PLoS One. 8(8):e71806. PMID: 23977147. doi:10.1371/journal.pone.0071806.
- 71. Schei K, Avershina E, Oien T, Rudi K, Follestad T, Salamati S, Ødegård RA. 2017. Early gut mycobiota and mother-offspring transfer. Microbiome. 5(1):107. doi:10.1186/s40168-017-0319-x. PMID: 28837002
- 72. Willis KA, Purvis JH, Myers ED, Aziz MM, Karabayir I, Gomes CK, Peters BM, Akbilgic O, Talati AJ, Pierre JF, et al. 2019. Fungi form interkingdom microbial communities in the primordial human gut that develop with gestational age. FASEB J. 33(11):12825-12837. PMID: 31480903. doi:10.1096/fj.201901436RR.
- 73. Yang AM, Inamine T, Hochrath K, Chen P, Wang L, Llorente C,Bluemel S, Hartmann P, Xu J, Koyama Y, et al. 2017. Intestinal fungi contribute to development of alcoholic liver disease. J Clin Invest. 127(7):2829-2841. PMID: 28530644. doi:10.1172/JCI90562.
- 74. Lang S, Duan Y, Liu J, Torralba MG, Kuelbs C, Ventura-Cots M, Abraldes JG, Bosques-Padilla F, Verna EC, Brown RS, et al. 2020. Intestinal fungal dysbiosis and systemic immune response to fungi in patients with alcoholic hepatitis. Hepatology. 71(2):522-538. PMID: 31228214. doi:10.1002/hep.30832.
- 75. Bajaj JS, Liu EJ, Kheradman R, Fagan A, Heuman DM, White M, Gavis EA, Hylemon P, Sikaroodi M, Gillevet PM, et al. 2018. Fungal dysbiosis in cirrhosis. Gut. 67 (6):1146-1154. PMID: 28578302. doi:10.1136/gutjnl-2016-313170.
- 76. Hartmann P, Lang S, Zeng S, Duan Y, Zhang X, Wang Y, Bondareva M, Kruglov A, Fouts DE, Stärkel P, et al. 2021. Dynamic changes of the fungal microbiome in alcohol use disorder. Front Physiol. 12:699253. PMID: 34349667. doi:10.3389/fphys.2021.699253.

- 77. Wu J, Wu D, Ma K, Wang T, Shi G, Shao J, Wang C, Yan G. 2020. Paeonol ameliorates murine alcohol liver disease via mycobiota-mediated Dectin-1/IL-1beta signaling pathway. J Leukoc Biol. 108(1):199-214. doi:10.1002/JLB.3MA0120-325RR. PMID: 32129526
- 78. Brown GD, Gordon S. 2001. Immune recognition. A new receptor for beta-glucans. Nature. 413 (6851):36-37. doi:10.1038/35092620. PMID: 11544516
- 79. Kimberg M, Brown GD. 2008. Dectin-1 and its role in antifungal immunity. Med Mycol. 46(7):631-636. doi:10.1080/13693780802140907. PMID: 18608924
- 80. Rogers NC, Slack EC, Edwards AD, Nolte MA, Schulz O, Schweighoffer E, Williams DL, Gordon S, Tybulewicz VL, Brown GD, et al. 2005. Syk-dependent cytokine induction by Dectin-1 reveals a novel pattern recognition pathway for C type lectins. Immunity. 22(4):507-517. PMID: 15845454. doi:10.1016/j.immuni.2005.03.004.
- 81. Gross O, Gewies A, Finger K, Schafer M, Sparwasser T, Peschel C, Förster I, Ruland J. 2006. Card9 controls a non-TLR signalling pathway for innate anti-fungal immunity. Nature. 442(7103):651-656. doi:10.1038/ nature04926. PMID: 16862125
- 82. Kankkunen P, Teirila L, Rintahaka J, Alenius H, Wolff H, Matikainen S. 2010. (1,3)-beta-glucans activate both dectin-1 and NLRP3 inflammasome in human macrophages. Immunol. J 184(11):6335-6342. doi:10.4049/jimmunol.0903019. PMID: 20421639
- 83. Gross O, Poeck H, Bscheider M, Dostert C, Hannesschlager N, Endres S, Hartmann G, Tardivel A, Schweighoffer E, Tybulewicz V, et al. 2009. Syk kinase signalling couples to the Nlrp3 inflammasome for anti-fungal host defence. Nature. 459(7245):433-436. PMID: 19339971. doi:10.1038/nature07965.
- 84. Petrasek J, Bala S, Csak T, Lippai D, Kodys K, Menashy V, Barrieau M, Min S-Y, Kurt-Jones EA, Szabo G, et al. 2012. IL-1 receptor antagonist ameliorates inflammasome-dependent alcoholic steatohepatitis in mice. J Clin Invest. 122(10):3476-3489. PMID: 22945633. doi:10.1172/JCI60777.
- 85. Sun S, Wang K, Sun L, Cheng B, Qiao S, Dai H, Shi W, Ma J, Liu H. 2020. Therapeutic manipulation of gut microbiota by polysaccharides of Wolfiporia cocos reveals the contribution of the gut fungi-induced PGE2 to alcoholic steatosis. Gut Microbes. 12(1):1830693. doi:10.1080/19490976.2020.1830693. PMID: 33106075
- 86. Moyes DL, Wilson D, Richardson JP, Mogavero S, Tang SX, Wernecke J, Höfs S, Gratacap RL, Robbins J, Runglall M, et al. 2016. Candidalysin is a fungal peptide toxin critical for mucosal infection. Nature. 532 (7597):64–68. PMID: 27027296. doi:10.1038/nature17625.
- 87. Chu H, Duan Y, Lang S, Jiang L, Wang Y, Llorente C, Liu J, Mogavero S, Bosques-Padilla F, Abraldes JG, et al. 2020. The Candida albicans exotoxin candidalysin promotes alcohol-associated liver disease. J Hepatol. 72(3):391-400. PMID: 31606552. doi:10.1016/j.jhep.2019.09.029.

- 88. Hill C, Guarner F, Reid G, Gibson GR, Merenstein DJ, Pot B, Morelli L, Canani RB, Flint HJ, Salminen S, et al. 2014. Expert consensus document. The international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. Nat Rev Gastroenterol Hepatol. 11 PMID: 24912386. doi:10.1038/ (8):506-514.nrgastro.2014.66.
- 89. Barssotti L, Abreu I, Brandao ABP, Albuquerque R, Ferreira FG, Salgado MAC, Dias DDS, De Angelis K, Yokota R, Casarini DE, et al. 2021. Saccharomyces boulardii modulates oxidative stress and renin angiotensin system attenuating diabetes-induced liver injury in Sci Rep. 11(1):9189. PMID: 33911129. doi:10.1038/s41598-021-88497-w.
- 90. Albuquerque R, Brandao ABP, De Abreu I, Ferreira FG, Santos LB, Moreira LN, Taddei CR, Aimbire F, Cunha TS. 2019. Saccharomyces boulardii Tht 500101 changes gut microbiota and ameliorates hyperglycaemia, dyslipidaemia, and liver inflammation in streptozotocin-diabetic mice. Benef Microbes. 10(8):901-912. doi:10.3920/ BM2019.0056. PMID: 31965836
- 91. Yu L, Zhao XK, Cheng ML, Yang GZ, Wang B, Liu HJ, Hu YX, Zhu LL, Zhang S, Xiao ZW, et al. 2017. Saccharomyces boulardii administration changes gut microbiota and attenuates D-Galactosamine-induced liver injury. Sci Rep. 7(1):1359. PMID: 28465509. doi:10.1038/s41598-017-01271-9.
- 92. Li M, Zhu L, Xie A, Yuan J. 2015. Oral administration of Saccharomyces boulardii ameliorates carbon tetrachloride-induced liver fibrosis in rats via reducing intestinal permeability and modulating gut microbial composition. Inflammation. 38(1):170-179. doi:10.1007/s10753-014-0019-7. PMID: 25227279
- 93. Fernandez-Pacheco P, Ramos Monge IM, Fernandez-Gonzalez M, Poveda Colado JM, Safety A-VM. 2021. Evaluation of yeasts with probiotic potential. Front PMID: 34095190. Nutr. 8:659328. doi:10.3389/ fnut.2021.659328.
- 94. Santino I, Alari A, Bono S, Teti E, Marangi M, Bernardini A, Magrini L, Di Somma S, Teggi A. 2014. Saccharomyces cerevisiae fungemia, a possible consequence of the treatment of clostridium difficile colitis with a probioticum. Int Immunopathol Pharmacol. 27(1):143-146. doi:10.1177/039463201402700120. PMID: 24674691
- 95. Cherifi S, Robberecht J, Miendje Y. 2004. Saccharomyces cerevisiae fungemia in an elderly patient with Clostridium difficile colitis. Acta Clin Belg. 59(4):223-224. doi:10.1179/ acb.2004.033. PMID: 15597730
- 96. Henry S, D'Hondt L, Andre M, Holemans X, Canon JL. 2004. Saccharomyces cerevisiae fungemia in a head and neck cancer patient: a case report and review of the literature. Acta Clin Belg. 59(4):220-222. doi:10.1179/ acb.2004.032. PMID: 15597729

- 97. Cesaro S, Chinello P, Rossi L, Zanesco L. 2000. Saccharomyces cerevisiae fungemia in a neutropenic patient treated with Saccharomyces boulardii. Support 8(6):504-505. Care Cancer. doi:10.1007/ s005200000123. PMID: 11094997
- 98. Lu H, Lou H, Hu J, Liu Z, Chen Q. 2020. Macrofungi: a review of cultivation strategies, bioactivity, and application of mushrooms. Compr Rev Food Sci Food Saf. 19 (5):2333-2356. doi:10.1111/1541-4337.12602. PMID: 33336985
- 99. Huang J, Ou Y, Yew TW, Liu J, Leng B, Lin Z,Su Y, Zhuang Y, Lin J, Li X, et al. 2016. Hepatoprotective effects of polysaccharide isolated from Agaricus bisporus industrial wastewater against CCl(4)-induced hepatic injury in mice. Int J Biol Macromol. 82:678-686. PMID: 26454111. doi:10.1016/j. ijbiomac.2015.10.014.
- 100. Li S, Li J, Zhang J, Wang W, Wang X, Jing H, Ren Z, Gao Z, Song X, Gong Z, et al. 2017. The antioxidative, antiaging, and hepatoprotective effects Alkali-Extractable polysaccharides by agaricus bisporus. Evid Based Complement Alternat Med. 2017:7298683. PMID: 29104605. doi:10.1155/2017/
- 101. Liu Y, Zheng D, Su L, Wang Q, Li Y. 2018. Protective effect of polysaccharide from Agaricus bisporus in Tibet area of China against tetrachloride-induced acute liver injury in mice. Int J Biol Macromol. 118:1488-1493. PMID: 29969638. doi:10.1016/j.ijbiomac.2018.06.179.
- 102. Duan Z, Zhang Y, Zhu C, Wu Y, Du B, Ji H. 2020. Structural characterization of phosphorylated Pleurotus ostreatus polysaccharide and its hepatoprotective effect on carbon tetrachloride-induced liver injury in mice. Int J Biol Macromol. 162:533-547. PMID: 32565302. doi:10.1016/j.ijbiomac.2020.06.107.
- 103. Bergh O, Borsheim KY, Bratbak G, Heldal M. High abundance of viruses found in aquatic environments. 1989;340(6233):467-468. doi:10.1038/ Nature. 340467a0. PMID: 2755508
- 104. Cobian Guemes AG, Youle M, Cantu VA, Felts B, Nulton J, Rohwer F. 2016. Viruses as winners in the game of life. Annu Rev Virol. 3(1):197-214. doi:10.1146/annurev-virology-100114-054952. PMID: 27741409
- 105. Shkoporov AN, Clooney AG, Sutton TDS, Ryan FJ, Daly KM, Nolan JA, McDonnell SA, Khokhlova EV, Draper LA, Forde A, et al. 2019. The human gut virome is highly diverse, stable, and individual specific. Cell Host Microbe. 26(4):527-41e5. PMID: 31600503. doi:10.1016/j.chom.2019.09.009.
- 106. Shkoporov AN, Hill C. 2019. Bacteriophages of the human gut: the "known unknown" of the microbiome. Cell Host Microbe. 25(2):195-209. doi:10.1016/j. chom.2019.01.017. PMID: 30763534

- 107. Roux S, Hallam SJ, Woyke T, Sullivan MB. 2015. Viral dark matter and virus-host interactions resolved from publicly available microbial genomes. Elife. 4 PMID: 26200428. doi:10.7554/eLife.08490.
- 108. Krishnamurthy SR, Wang D. 2017. Origins and challenges of viral dark matter. Virus Res. 239:136–142. PMID: 28192164. doi:10.1016/j.virusres.2017.02.002.
- 109. Clooney AG, Sutton TDS, Shkoporov AN, Holohan RK, Daly KM, O'Regan O, Ryan FJ, Draper LA, Plevy SE, Ross RP, et al. 2019. Whole-Virome analysis sheds light on viral dark matter in inflammatory bowel disease. Cell Host Microbe. 26(6):764–78e5. PMID: 31757768. doi:10.1016/j.chom.2019.10.009.
- 110. Gregory AC, Zablocki O, Zayed AA, Howell A, Bolduc B, Sullivan MB. 2020. The gut virome database reveals age-dependent patterns of virome diversity in the human gut. Cell Host Microbe. 28(5):724–40e8. doi:10.1016/j.chom.2020.08.003. PMID: 32841606
- 111. Camarillo-Guerrero LF, Almeida A, Rangel-Pineros G, Finn RD, Lawley TD. 2021. Massive expansion of human gut bacteriophage diversity. Cell. 184(4):1098– 109e9. doi:10.1016/j.cell.2021.01.029. PMID: 33606979
- Wommack KE, Colwell RR. 2000. Virioplankton: viruses in aquatic ecosystems. Microbiol Mol Biol Rev. 64:69–114.
 PMID: 10704475. doi:10.1128/mmbr.64.1.69-114.2000.
- 113. Dalmasso M, Hill C, Ross RP. 2014. Exploiting gut bacteriophages for human health. Trends Microbiol. 22(7):399–405. doi:10.1016/j.tim.2014.02.010. PMID: 24656964
- 114. Manrique P, Bolduc B, Walk ST, van der Oost J, de Vos WM, Young MJ. 2016. Healthy human gut phageome. Proc Natl Acad Sci U S A. 113 (37):10400–10405. doi:10.1073/pnas.1601060113. PMID: 27573828
- 115. Ross A, Ward S, Hyman P. 2016. More is better: selecting for broad host range bacteriophages. Front Microbiol. 7:1352. PMID: 27660623. doi:10.3389/fmicb.2016.01352.
- 116. Cahill J, Young R. 2019. Phage lysis: multiple genes for multiple barriers. Adv Virus Res. 103:33–70. PMID: 30635077. doi:10.1016/bs.aivir.2018.09.003.
- 117. Howard-Varona C, Hargreaves KR, Abedon ST, Sullivan MB. 2017. Lysogeny in nature: mechanisms, impact and ecology of temperate phages. ISME J. 11(7):1511–1520. doi:10.1038/ismej.2017.16. PMID: 28291233
- 118. Human Microbiome Jumpstart Reference Strains C. KE N, GM W, SK H, KC W, HH C, Wortman JR, Rusch DB, Mitreva M, Sodergren E, Chinwalla AT, et al. 2010. A catalog of reference genomes from the human microbiome. Science. 3285981: 994–999. 10.1126/science.1183605PMID: 20489017
- 119. Lengeling A, Mahajan A, Gally DL. 2013. Bacteriophages as pathogens and immune modulators?. mBio. 4(6):e00868–13. doi:10.1128/mBio.00868-13. PMID: 24222490

- 120. Scotland SM, Smith HR, Willshaw GA, Rowe B. 1983. Vero cytotoxin production in strain of Escherichia coli is determined by genes carried on bacteriophage. Lancet. 2(8343):216. doi:10.1016/s0140-6736(83) 90192-7. PMID: 6135046
- 121. Tetz GV, Ruggles KV, Zhou H, Heguy A, Tsirigos A, Tetz V. 2017. Bacteriophages as potential new mammalian pathogens. Sci Rep. 7(1):7043. doi:10.1038/s41598-017-07278-6. PMID: 28765534
- 122. Norman JM, Handley SA, Baldridge MT, Droit L, Liu CY, Keller BC, Kambal A, Monaco C, Zhao G, Fleshner P, et al. 2015. Disease-specific alterations in the enteric virome in inflammatory bowel disease. Cell. 160(3):447–460. PMID: 25619688. doi:10.1016/j.cell.2015.01.002.
- 123. Ma Y, You X, Mai G, Tokuyasu T, Liu C. 2018. A human gut phage catalog correlates the gut phageome with type 2 diabetes. Microbiome. 6(1):24. doi:10.1186/s40168-018-0410-y. PMID: 29391057
- 124. Monaco CL, Gootenberg DB, Zhao G, Handley SA, Ghebremichael MS, Lim ES, Lankowski A, Baldridge M, Wilen C, Flagg M, et al. 2016. Altered virome and bacterial microbiome in human immunodeficiency virus-associated acquired immunodeficiency syndrome. Cell Host Microbe. 19(3):311–322. PMID: 26962942. doi:10.1016/j.chom.2016.02.011.
- 125. Nakatsu G, Zhou H, Wu WKK, Wong SH, Coker OO, Dai Z, Li X, Szeto C-H, Sugimura N, Lam TYT, et al. 2018. Alterations in enteric virome are associated with colorectal cancer and survival outcomes. Gastroenterology. 155(2):529–41e5. PMID: 29689266. doi:10.1053/j.gastro.2018.04.018.
- 126. Koonin EV, Dolja VV, Krupovic M. 2021. The healthy human virome: from virus-host symbiosis to disease. Curr Opin Virol. 47:86–94. PMID: 33652230. doi:10.1016/j.coviro.2021.02.002.
- 127. Jiang L, Lang S, Duan Y, Zhang X, Gao B, Chopyk J, Schwanemann LK, Ventura-Cots M, Bataller R, Bosques-Padilla F, et al. 2020. Intestinal virome in patients with alcoholic hepatitis. Hepatology. 72(6):2182–2196. PMID: 32654263. doi:10.1002/hep.31459.
- 128. Rojo J, Simoes P, Krueger GR, Humberto CO, Ramon AM. 2003. Human herpesvirus-6 has no apparent influence on course of HCV hepatitis, but may complicate HBV hepatitis and alcoholic liver disease. A pilot study. Vivo. 17:29–33. PMID: 12655786.
- 129. Yurlov KI, Masalova OV, Kisteneva LB, Khlopova IN, Samokhvalov EI, Malinovskaya VV,Parfyonov VV, Shuvalov AN, Kushch AA. 2021. Human herpesviruses increase the severity of hepatitis. Biology (Basel). 10. PMID: 34072365. doi:10.3390/biology10060483.
- 130. Tsai JP, Tseng KC, Lin MN, Su CC. 2019. A high seroprevalence of human herpesvirus type 8 already present in patients with chronic hepatitis before the development of cirrhosis. Pathology. 51(1):86–90. doi:10.1016/j.pathol.2018.10.005. PMID: 30497802

- 131. Chou AL, Huang WW, Tsao SM, Li CT, Su CC. 2008. Human herpesvirus type 8 in patients with cirrhosis: correlation with sex, alcoholism, hepatitis B virus, disease severity, and thrombocytopenia. Am J Clin Pathol. 130(2):231-237. doi:10.1309/4FMBN316N4792UQ4. PMID: 18628092
- 132. Hu J, Zhang X, Yu G, Cai H, Gu J, Hu M, Xiang D, Lian J, Yu L, Jia H, et al. 2019. Epstein-Barr virus infection is associated with a higher Child-Pugh score and may predict poor prognoses for patients with liver cirrhosis. BMC Gastroenterol. 19(1):94. PMID: 31215410. doi:10.1186/s12876-019-1021-1.
- 133. Keen EC, Dantas G. 2018. Close encounters of three kinds: bacteriophages, commensal bacteria, and host immunity. Trends Microbiol. 26(11):943-954. doi:10.1016/j. tim.2018.05.009. PMID: 29909042
- 134. Bajaj JS, Sikaroodi M, Shamsaddini A, Henseler Z, Santiago-Rodriguez T, Acharya C, Fagan A, Hylemon PB, Fuchs M, Gavis E, et al. 2021. ssInteraction of bacterial metagenome and virome in patients with cirrhosis and hepatic encephalopathy. Gut. 70(6):1162-1173. PMID: 32998876. doi:10.1136/ gutjnl-2020-322470.
- 135. Pirnay JP, Blasdel BG, Bretaudeau L, Buckling A, Chanishvili N, Clark JR, Corte-Real S, Debarbieux L, Dublanchet A, De Vos D, et al. 2015. Quality and safety requirements for sustainable phage therapy products. Pharm Res. 32(7):2173-2179. PMID: 25585954. doi:10.1007/s11095-014-1617-7.
- 136. Cheroutre H, Madakamutil L. 2004. Acquired and natural memory T cells join forces at the mucosal front line. Nat Rev Immunol. 4(4):290-300. doi:10.1038/ nri1333. PMID: 15057787
- 137. Liu L, Gong T, Tao W, Lin B, Li C, Zheng X, Zhu S, Jiang W, Zhou R. 2019. Commensal viruses maintain intestinal intraepithelial lymphocytes via noncanonical RIG-I signaling. Nat Immunol. 20(12):1681-1691. doi:10.1038/s41590-019-0513-z. PMID: 31636462
- 138. Gogokhia L, Buhrke K, Bell R, Hoffman B, Brown DG, Hanke-Gogokhia C, Ajami NJ, Wong Ghazaryan A, Valentine JF, et al. 2019. Expansion of bacteriophages is linked to aggravated intestinal inflammation and colitis. Cell Host Microbe. 25(2):285-99e8. PMID: 30763538. doi:10.1016/j.chom.2019.01.008.
- 139. Metzger RN, Krug AB, Eisenacher K. 2018. Enteric virome sensing-its role in intestinal homeostasis and immunity. Viruses. 10(4):146. doi:10.3390/v10040146. PMID: 29570694
- 140. Broquet AH, Hirata Y, McAllister CS, MF K. 2011. RIG-I/MDA5/MAVS are required to signal a protective IFN response in rotavirus-infected intestinal epithelium. J Immunol. 186(3):1618-1626. doi:10.4049/jimmunol.1002862. PMID: 21187438
- 141. Norman JM, Handley SA, Virgin HW. 2014. Kingdomagnostic metagenomics and the importance of complete characterization of enteric microbial communities. Gastroenterology. 146(6):1459-1469. doi:10.1053/j.gastro.2014.02.001. PMID: 24508599

- 142. Duan Y, Llorente C, Lang S, Brandl K, Chu H, Jiang L, White RC, Clarke TH, Nguyen K, Torralba M, et al. 2019. Bacteriophage targeting of gut bacterium attenuates alcoholic liver disease. Nature. 575(7783):505-511. PMID: 31723265. doi:10.1038/s41586-019-1742-x.
- 143. Kortright KE, Chan BK, Koff JL, Turner PE. 2019. Phage therapy: a renewed approach to combat antibiotic-resistant bacteria. Cell Host Microbe. 25(2):219-232. doi:10.1016/j. chom.2019.01.014. PMID: 30763536
- 144. Loc-Carrillo C, Abedon ST. 2011. Pros and cons of therapy. Bacteriophage. 1(2):111-114. phage doi:10.4161/bact.1.2.14590. PMID: 22334867
- 145. DeLong EF. 1992. Archaea in coastal marine environments. Proc Natl Acad Sci U S A. 89 (12):5685-5689. doi:10.1073/pnas.89.12.5685. PMID: 1608980
- 146. Schleper C, Jurgens G, Jonuscheit M. 2005. Genomic studies of uncultivated archaea. Nat Rev Microbiol. 3 (6):479-488.doi:10.1038/nrmicro1159. PMID: 15931166
- 147. Auguet JC, Barberan A, Casamayor EO. 2010. Global ecological patterns in uncultured Archaea. ISME J. 4 (2):182-190.doi:10.1038/ismej.2009.109. PMID: 19847207
- 148. Pereira O, Hochart C, Auguet JC, Debroas D, Galand PE. 2019. Genomic ecology of Marine Group II, the most common marine planktonic Archaea across the surface ocean. Microbiologyopen. 8(9):e00852. doi:10.1002/mbo3.852. PMID: 31264806
- 149. Santoro AE, Richter RA, Dupont CL. 2019. Planktonic marine archaea. Ann Rev Mar Sci. 11(1):131-158. doi:10.1146/annurev-marine-121916-063141. PMID: 30212260
- 150. Albers SV, Meyer BH. 2011. The archaeal cell envelope. Nat Rev Microbiol. 9(6):414-426. doi:10.1038/nrmicro2576. PMID: 21572458
- 151. Koga Y, Morii H. 2007. Biosynthesis of ether-type polar lipids in archaea and evolutionary considerations. Microbiol Mol Biol Rev. 71(1):97-120. doi:10.1128/ MMBR.00033-06. PMID: 17347520
- 152. Valentine DL. 2007. Adaptations to energy stress dictate the ecology and evolution of the archaea. Nat Rev Microbiol. 5(4):316-323. doi:10.1038/nrmicro1619. PMID: 17334387
- 153. Borrel G, McCann A, Deane J, Neto MC, Lynch DB, Brugere JF,O'Toole PW. 2017. Genomics and metagenomics of trimethylamine-utilizing archaea in the human gut microbiome. ISME J. 11(9):2059-2074. PMID: 28585938. doi:10.1038/ismej.2017.72.
- 154. Dridi B, Henry M, El Khechine A, Raoult D, Drancourt M, Dobrindt U. 2009. High prevalence of methanobrevibacter smithii and methanosphaera stadtmanae detected in the human gut using an improved DNA detection protocol. PLoS One. 4(9): e7063. doi:10.1371/journal.pone.0007063. 19759898

967271.

- 155. Horz HP, Conrads G. 2010. The discussion goes on: what is the role of Euryarchaeota in humans? Archaea. 2010:967271. PMID: 21253553. doi:10.1155/2010/
- 156. Hajishengallis G, Darveau RP, Curtis MA. 2012. The keystone-pathogen hypothesis. Nat Rev Microbiol. 10 (10):717–725. doi:10.1038/nrmicro2873. PMID: 22941505
- 157. Gaci N, Borrel G, Tottey W, O'Toole PW, Brugere JF. 2014. Archaea and the human gut: new beginning of an old story. World J Gastroenterol. 20(43):16062–16078. doi:10.3748/wjg.v20.i43.16062. PMID: 25473158
- 158. Borrel G, Brugere JF, Gribaldo S, Schmitz RA, Moissl-Eichinger C. 2020. The host-associated archaeome. Nat Rev Microbiol. 18(11):622–636. doi:10.1038/s41579-020-0407-y. PMID: 32690877
- 159. Bang C, Schmitz RA, Robinson NP. 2018. Archaea: forgotten players in the microbiome. Emerg Top Life Sci. 2(4):459–468. doi:10.1042/ETLS20180035. PMID: 33525830
- 160. Pausan MR, Csorba C, Singer G, Till H, Schopf V, Santigli E, Klug B, Högenauer C, Blohs M, Moissl-Eichinger C, et al. 2019. Exploring the archaeome: detection of archaeal signatures in the human body. Front Microbiol. 10:2796. PMID: 31866971. doi:10.3389/fmicb.2019.02796.
- 161. Grine G, Lotte R, Chirio D, Chevalier A, Raoult D, Drancourt M, Ruimy R. 2019. Co-culture of Methanobrevibacter smithii with enterobacteria during urinary infection. EBioMedicine. 43:333–337. PMID: 31072770. doi:10.1016/j.ebiom.2019.04.037.
- 162. Lepp PW, Brinig MM, Ouverney CC, Palm K, Armitage GC, Relman DA. 2004. Methanogenic archaea and human periodontal disease. Proc Natl Acad Sci U S A. 101(16):6176–6181. doi:10.1073/pnas.0308766101. PMID: 15067114
- 163. Blais Lecours P, Marsolais D, Cormier Y, Berberi M, Hache C, Bourdages R, Duchaine C.
 2014. Increased prevalence of methanosphaera stadtmanae in inflammatory bowel diseases. PLoS One. 9(2):e87734. doi:10.1371/journal. pone.0087734. PMID: 24498365
- 164. Pimentel M, Lin HC, Enayati P, van den Burg B, Lee HR, Chen JH, Park S, Kong Y, Conklin J. 2006. Methane, a gas produced by enteric bacteria, slows intestinal transit and augments small intestinal contractile activity. Am J Physiol Gastrointest Liver Physiol. 290(6):G1089–95. doi:10.1152/ajpgi.00574.2004. PMID: 16293652

- 165. Triantafyllou K, Chang C, Pimentel M. 2014. Methanogens, methane and gastrointestinal motility. J Neurogastroenterol Motil. 20(1):31–40. doi:10.5056/jnm.2014.20.1.31. PMID: 24466443
- 166. Bang C, Weidenbach K, Gutsmann T, Heine H, Schmitz RA, Foligne B. 2014. The intestinal archaea methanosphaera stadtmanae and methanobrevibacter smithii activate human dendritic cells. PLoS One. 9(6):e99411. doi:10.1371/journal. pone.0099411. PMID: 24915454
- 167. Blais Lecours P, Duchaine C, Taillefer M, Tremblay C, Veillette M, Cormier Y, Marsolais D. 2011. Immunogenic properties of archaeal species found in bioaerosols. PLoS One. 6(8):e23326. doi:10.1371/journal.pone.0023326. PMID: 21858070
- 168. Vierbuchen T, Bang C, Rosigkeit H, Schmitz RA, Heine H. 2017. The human-associated archaeon methanosphaera stadtmanae is recognized through its RNA and induces TLR8-Dependent NLRP3 inflammasome activation. Front Immunol. 8:1535. PMID: 29181003. doi:10.3389/fimmu.2017.01535.
- 169. Bang C, Vierbuchen T, Gutsmann T, Heine H, Schmitz RA, Proost P. 2017. Immunogenic properties of the human gut-associated archaeon methanomassi-liicoccus luminyensis and its susceptibility to antimicrobial peptides. PLoS One. 12(10):e0185919. doi:10.1371/journal.pone.0185919. PMID: 28982164
- 170. Samuel BS, Hansen EE, Manchester JK, Coutinho PM, Henrissat B, Fulton R, Latreille P, Kim K, Wilson RK, Gordon JI, et al. 2007. Genomic and metabolic adaptations of Methanobrevibacter smithii to the human gut. Proc Natl Acad Sci U S A. 104(25):10643–10648. PMID: 17563350. doi:10.1073/pnas.0704189104.
- 171. Gao B, Zhang X, Schnabl B. 2021. Fungi-bacteria correlation in alcoholic hepatitis patients. Toxins (Basel). 13 (2):143. doi:10.3390/toxins13020143. PMID: 33672887
- 172. Gilbert JA, Quinn RA, Debelius J, Xu ZZ, Morton J, Garg N, Jansson JK, Dorrestein PC, Knight R. 2016. Microbiome-wide association studies link dynamic microbial consortia to disease. Nature. 535(7610):94–103. doi:10.1038/nature18850. PMID: 27383984
- 173. Wu X, Xia Y, He F, Zhu C, Ren W. 2021. Intestinal mycobiota in health and diseases: from a disrupted equilibrium to clinical opportunities. Microbiome. 9(1):60. doi:10.1186/s40168-021-01024-x. PMID: 33715629
- 174. Townsend EM, Kelly L, Muscatt G, Box JD, Hargraves N, Lilley D, Jameson E. 2021. The human gut phageome: origins and roles in the human gut microbiome. Front Cell Infect Microbiol. 11:643214. PMID: 34150671. doi:10.3389/fcimb.2021.643214.