

Status and trends of nutraceuticals from onion and onion by-products: A critical review

Muhammad Sajid Arshad, Muhammad Sohaib, Muhammad Nadeem, Farhan Saeed, Ali Imran, Ahsan Javed, Zaid Amjad & Syeda Mamoonah Batool |

To cite this article: Muhammad Sajid Arshad, Muhammad Sohaib, Muhammad Nadeem, Farhan Saeed, Ali Imran, Ahsan Javed, Zaid Amjad & Syeda Mamoonah Batool | (2017) Status and trends of nutraceuticals from onion and onion by-products: A critical review, Cogent Food & Agriculture, 3:1, 1280254, DOI: [10.1080/23311932.2017.1280254](https://doi.org/10.1080/23311932.2017.1280254)

To link to this article: <https://doi.org/10.1080/23311932.2017.1280254>



© 2017 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license



Published online: 09 Feb 2017.



Submit your article to this journal [↗](#)



Article views: 6274



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 18 View citing articles [↗](#)



Received: 24 October 2016
Accepted: 02 January 2017
First Published: 13 January 2017

*Corresponding author: Muhammad Sajid Arshad, Institute of Home and Food Sciences, Government College University, Faisalabad, Pakistan
E-mail: sajid_ft@yahoo.com

Reviewing editor:
Fatih Yildiz, Middle East Technical University, Turkey

Additional information is available at the end of the article

FOOD SCIENCE & TECHNOLOGY | REVIEW ARTICLE

Status and trends of nutraceuticals from onion and onion by-products: A critical review

Muhammad Sajid Arshad^{1*}, Muhammad Sohaib², Muhammad Nadeem³, Farhan Saeed¹, Ali Imran¹, Ahsan Javed¹, Zaid Amjad¹ and Syeda Mamoonah Batool¹

Abstract: The pivotal roles of vegetables are predominantly important for health due to availability of minerals, vitamins, phytochemicals and antioxidants. These foods and food products those are prepared from such kind of material having natural origin have great importance due to their versatile nature. Onion importance is greatly increasing and now it has become second most medicinal and horticultural crop after tomatoes. It has multiple functional compounds including organosulphur, anthocyanins, flavonoids, quercetin, kaempferol and polyphenols. Antibacterial, anti-inflammatory and antioxidant properties are the various diverse functions of onions, which have sound effects on human health. Fructo-oligosaccharides are present in abundant quantity in it due to which the growth of potentially harmful bacteria in the colon is retarded by the oligomers thus reducing the risk of emerging tumors in the colon and also initiates the growth of healthy bifidobacterium. In the nutshell, the onion and onion by-products have a lot of bioactive compounds against certain metabolic disorders.

Subjects: Environment & Agriculture; Food Science & Technology; Health and Social Care

Keywords: onion; anthocyanins; flavonoids; quercetin; kaempferol; polyphenols

1. Introduction

Onion (*Allium cepa* L.) is the main crop grown all over the world especially in the Asia. It belongs to the family, Amaryllidaceae, and the genus, Allium. Alliums are perennial herbs having bulbous and scented underground stems. This genus includes garlic, shallot, chives, leeks, and even a non-edible species grown only for its showy flower (Mollavali et al., 2016). The common garden onions are in the species, *A. cepa*. There are various varieties of onion each with their own unique flavor, ranging from mildly sweet to very strong i.e. red, yellow, white and green. It can be eaten fried, dried, raw, cooked or roasted. They are usually used to flavor salads, spreads, stir-fry, dips, soups and other dishes (van Wyk, 2014). Since ancient times in various cultures, onions on every continent have been growing in their natural habitat. Dates back to 3500 BC, onions were one of the only some foods that did not

ABOUT THE AUTHOR

Muhammad Sajid Arshad is Assistant Professor at Department of Food Science, Government College University, Faisalabad, Pakistan. He has published a number of research and review articles related to his research area in high repute Journals.

PUBLIC INTEREST STATEMENT

Onion and onion by-products have vital role against certain ailments due to the presence of bioactive compounds present in it. Before people did not use the by-products but now the concept of value addition is very popular. Pharmaceutically onion possesses many characteristics like anti-inflammatory, antioxidant and antibacterial properties. In decreasing pro-inflammatory diseases the effectiveness of onion is rooted in its nature as modulators of metabolism.

spoil during the winter seasons. Ancestors in Asia have become familiar with its stability and started to grow as a food. Swenson (2008) reported that the ancient Egyptians worshipped the onion, having faith in its spherical shape and concentric rings that represented eternity. Pakistan being an agricultural country, is producing onion as a major agricultural crop. The annual world production of onion is about 55 million tonnes and Pakistan ranked 8th in onion production with an area of 147.6 thousand hectares and production of 1,939.6 thousand tonnes (Mushtaq et al., 2013). Swenson (2008) proposed that in the mountainous areas of Pakistan, Tajikistan, Northern Iran and Afghanistan the bulb onion (*A. cepa* L.) has been domesticated and cultivated thousands of years ago. Allium species (onion bulbs and fresh shoots) are grown for the production of seeds, sets and as fresh shoots for green salad in open, sunny and dry mountain slopes of these countries. Onions vary in color, size, firmness, pungency shape, stiffness of outer neck and dry skins and may be consumed as uncooked or pickled (Akhtar, 2015).

Short-day and long-day onions are two major types being cultivated all over Pakistan. On the other hand, a third group named as an intermediary day length type is expressed between these two major groups. White and Zellner (2008) described that the onions which require 11–12 h to bulb are placed under short-day type category whereas those require more than 14 h to bulb are categorized as long day type of onions.

The chemical composition of onions for moisture, carbohydrate, total sugar, vitamin-C, Ca, P and K ranged from (82.99 and 82.77%), (14.146 and 14.772%), (4.74 and 2.32%), (6.5 and 5.7 mg), (46.9 and 25.7 mg), (50.6 and 30.3 mg) and (140 and 129 mg), respectively. Likewise, Bhattacharjee et al. (2013) revealed the level of protein and fat as (2.62 and 1.489%) and (0.4%) with other trace elements. Major components of commercially important volatile compounds of onion includes 3,5-Diethyl-1,2,4-trithionale and Propyl 1-propenyl disulphide (Farkaas, Hradsky, & Kovac, 1992). Whereas, amongst non-volatile components Sitosterol, Gitogenin, Oleanolic acid, Amyrin and Diosgenin are abundantly present (Ismailov & Aliev, 1974).

2. Onions and health interactions

Onion is the main component of condiments used in all households around the year. Small bulbs of onions are pickled in vinegar and are used for making various seasonings, sauces and soups. Onion is a spice with characteristic flavor and aroma as well as having pronounced medicinal importance such as rich in carbohydrates, phosphorus and calcium. Lee and Mitchell (2011) and Suleria, Butt, Anjum, Saeed, and Khalid (2013) have reported high levels of ACSOs (alk(en)yl cysteine sulfoxides) present in Allium plants. Onions are being used since primeval times as a traditional medicine. By having rich photochemistry, these reduce the peril of cardiovascular diseases, diabetes, cancer and atherosclerosis (Cazzola, Camerotto, & Cestaro, 2011; Suleria et al., 2013).

3. Role of onion as antioxidant

BHT (butylated hydroxytoluene) and BHA (butylated hydroxyanisole) are synthetic antioxidants which are needed to be exchanged with natural antioxidants, as several studies showed that a number of synthetic antioxidants were toxic and carcinogenic (Che Othman et al., 2011). Antioxidant compounds (flavonoids and phenolics), have protective effects against different degenerative pathologies are present in high amount in onions (Pérez-Gregorio, Regueiro, González-Barreiro, Rial-Otero, & Simal-Gándara, 2011; Siddiq, Roidoung, Sogi, & Dolan, 2013). Onion consider as a good antioxidant additive for food due to Cysteine derivatives (Ostrowska et al., 2004), enhancing its usability as in ethnomedicine and as functional food (Tram Ngoc et al., 2005). Chemical spoilage, resulting in deterioration or rancidity of the texture, flavor color, nutritional quality and safety of foods is caused by the oxidation (Antolovich, Prenzler, Patsalides, McDonald, & Robards, 2002). Flavonoid extractability is increased during onion processing. Nemeth and Piskula (2007) reported that enzymes released from disrupted plant tissue causes deglycosylation, and after ingestion, due to β -glucosidases of microbial origin and those of the consumer's body.

4. Disparity in antioxidant activity

Onion antioxidant activity is reduced after cooking, so onion is efficient in its raw form (Ali, Thomson, & Afzal, 2000). Yang, Meyers, van der Heide, and Liu (2004) reported that various kinds of onions were found which differ in their characteristics, with maximum *in vitro* tumor cell inhibition as well as highest total antioxidant activities seen in the onion variety Western Yellow and in shallots. The synthesis of amino acid and its transformation to leukotrienes and pro-inflammatory prostaglandins is inhibited by cepaene and thiosulphinate content present in onion (Ali et al., 2000). Cao, Sofic, and Prior (1996) suggest that many vegetables, fruits, and tea's antioxidant activity have been studied, which pointed out that the presence of vitamin C, β -carotene and flavonoids shows the antioxidant activity of these plant foods (Bors & Saran, 1987; Gey, 1990; Willett, 1994). Cao et al. (1996) studied two members of Allium family are also known to have antioxidant activity, which are Garlic bulb and onion (Yang, Yasaei, & Page, 1993).

There is little information available about the antioxidant activity of other members of the allium family; however, antioxidant capacities of Garlic and onion bulbs have been investigated briefly. For the antioxidant activity of garlic bulb, Allicin which is a scavenger of peroxy radicals was responsible (Zakarova et al., 2014). Thiosulfinates and allicin configuration in allium members and garlic have been studied shows allicin is absent in onion and rich in thiosulfinates compounds which play a key role in onion antioxidant activity (Block, Naganathan, Putman, & Zhao, 1992; Lawson, Wood, & Hughes, 1991).

5. Onion and onion by products as antimicrobial and antifungal

Significant amounts of phenolic-rich by-products are being produced in food processing industries, which could be important natural sources of antioxidants. Roldán, Sánchez-Moreno, de Ancos, and Cano (2008) reported that we should worried about the production of large quantities of industrial onion wastage and its disposal. Stabilizing and processing onion wastes (surpluses and residues of onion) could symbolize advantages (Roldán et al., 2008). Literature shows that onions contain the powerful antioxidants, sulfur and other numerous phenolic compounds which provoke great interests and holds anti-bacterial and anti-fungal activities (Onyeoziri, Romanus, & Onyekachukwu, 2016).

Onion and onion by-product has high amount of antioxidants which makes it a good free radical scavenging ability. So, it has good anti-inflammatory and anti-mutagenic properties. Lipid peroxidation was found to be inhibited in rat liver microcosms and also found to effectively scavenge superoxide anions (Krishnakantha & Lokesh, 1993; Reddy & Lokesh, 1992). The antioxidant power of onion by product has also been confirmed in applications where extracts of onion by product was added to meat products. Fresh, frozen and precooked pork patties were further tested to check the antioxidant effectiveness of onion by product extract. By the inclusion of onion by product extract, the shelf life of all products was enhanced which was determined by TBA value (Cao et al., 2013).

Onion by product has strong antibacterial. Active constituents of onion by product restrict the growth of *Streptococci*, *Staphylococci*, *Proteus* sp., *Escherichia coli* and *Salmonella* *in vitro* studies (Janes, Nannapaneni, & Johnson, 1999). Meena (1998) reported that at ambient temperatures fresh onion by product juice showed inhibitory action against *S. cerevisiae*, *A. niger*, *Mycoderma* spp. and *L. acidophilus* at 4, 10, 12 and 14% respectively.

In vitro and *in vivo* onion showed effective antioxidant activity (Jackson et al., 2002). Volatile oils (allicin, ajoene and alliin) consisting of sulphur, carbohydrates (sucrose, glucose), amino acids like methionine, glutamine, cysteine and isoleucine, minerals (selenium, zinc, germanium), bioflavonoids like cyanidin and quercetin and allistatin I and allistatin II, A, E and C vitamins, enzymes (miracynase, peroxidase and alliinase), and niacin, B₂ vitamins and beta carotene fall in >20 is a plant0 components present in onion (Gulsen & Ayaz, 2010). Onion lowers glycaemia and cholesterol and kills parasites, fungus, bacteria and has the ability to protect liver and contains antitumor agents. Onion has the ability of protecting human body against various illnesses with more than 200 chemical substances present in its body. For protection against infections and free radicals onion should

be consumed as fresh (Ayaz & Alpsoy, 2007). Harris (2001) reported that onion has a wide range of actions; not only as antiprotozoal, antifungal, antibacterial and antiviral, but also has positive effects on the immune systems and cardiovascular.

6. Anthocyanins as antioxidant in onion

There is little information regarding absolute amount of anthocyanins in red onions. Ferreres, Gil, and Tomás-Barberán (1999) and Clifford (2000) reported that up to 250 mg/kg anthocyanins present in red onion. Nevertheless, it is not sure that this amount relates only to the edible part or to the whole onion bulb. It is also reported that the analysis performed on dry skins shows that values extending from a minimum (109/100 g) to a maximum (219/100 g) (Donner, Gao, & Mazza, 1997; Fossen, Andersen, Øvstedal, Pedersen, & Raknes, 1996). Flavonoids occur in a variety of complex conjugates with sugars, so sugars are very substantial for anthocyanin biosynthesis, but data about this stuff is still controversial (Harborne & Mabry, 2013). In the context of a survey aimed on the typical Italian varieties characterization, produced in some areas of southern Italy and was investigated. Tropea red onion is known for its crispness and sweetness that is why it is consumed uncooked which preserve the flavonoid compounds (Gennaro et al., 2002; Ioku et al., 2001).

7. Onion as disease preventive

Antioxidants hunt free radicals and are related to minimize the risk of cancer and cardiovascular diseases. Stajner and Varga (2003) reported that *A. cepa* L. has antispasmodic, hypoglycaemic, antimicrobial, hypotensive, antiasthmatic, anticholesterolaemic, anticancer, and antioxidant properties. In onions polyphenols, anthocyanins, quercetin, kaempferol, flavonoids and their glycosides have been reported (Ashwini, Balaganesh, Balamurugan, Murugan, & Sathishkumar, 2013). The studies proposed that fruits and vegetables have a defensive action against coronary heart disease and cancer. Much attention has been given for lowering the oxidative stress prompted by chronic diseases (Crozier et al., 2000). Furthermore, *Allium* species are strongly resistant to diseases produced by nematodes. Onion is very helpful against free radicals and cancer due to antioxidant activity and antibacterial activity. It also consists of some terpenoids, carotenoids, flavonoids, vitamins, phytoestrogens and minerals etc. (Calucci, Pinzino, Zandomenighi, & Capocchi, 2003).

8. Phytochemical synthesis

Pasteurized Recas product contained 721 mg/100 g quercetin of dry weight whereas frozen Recas paste contain highest total quercetin content of 4,431 mg/100 g of dry weight i.e. 4%. Major products of lipid oxidation are free radicals which implicated over 100 important diseases including atherosclerosis, cancer, arthritis (Hajhashemi, Vaseghi, Pourfarzam, & Abdollahi, 2010). Taste and toxicity both are undesired which are formed by the oxidation of unsaturated fatty acids (Benkeblia, 2005). The total quercetin concentration in onion is from traces in white varieties to 2.5–3 mmol/kg of fresh weight in red varieties. There are four predominant forms of Q in onions exists in: Q3G (quercetin-3-O-glucoside), Q3,4'G (quercetin-3,4-di-O-glucoside), Q aglycone, Q4'G (quercetin-4-O-glucoside), and many forms of monoglucoside and diglucoside conjugates in lesser amounts have been reported (Wiczowski, Nemeth, Bucin'ski, & Piskula, 2003).

Among fruits and vegetables, onion rated maximum in quercetin (Q) content (Ko, Cheigh, Cho, & Chung, 2011). Lombard, Peffley, Geoffriau, Thompson, and Herring (2005) reported that bulb type and color vary the concentration of (Q) in onions and quercetin is spread typically in the outer rings and skins. Mainly aglycones are the end result of food processing but glycosylated, flavonols and other flavonoids are present in fresh vegetables (Rohn, Buchner, Driemel, Rauser, & Kroh, 2007). In onions secondary metabolites occur in two main classes; flavonoids and organosulfur compounds having beneficial health effects and methods of biosynthetic pathways and action are rather changed.

El-Hadidy, Mossa, and Habashy (2014) reported that by inhibiting lipoxygenase and cyclooxygenase enzymes the organosulfur compounds are thought to have anti-allergic, anti-thrombotic, antimicrobial and anti-inflammatory activity. Probably, the compounds work through sulfur-oxygen or

sulfur-sulfur bonds. When onion cell walls is disrupted and cut these compounds are produced. El-Hadidy et al. (2014) depicted that via S-alk (en)yl cysteine sulfoxides (ACSOs) Sulfenic acids are produced by Allinase enzymes which reposition to numerous compounds such as, cepaenes and thiosulfinates. Flavonoids comprise aromatic rings held together by a C₃ unit. The subclass of flavonoid is determined by the rate of C₃ unit oxidation such as anthocyanins, flavonols, and flavanones. Chalcone formation starts the Flavonoid biosynthesis by chalcone synthase from coumaroyl CoA and malonyl CoA, which is a derive form of amino acid phenylalanine. Flavanone is being isomerized by the chalcone. The formation of each flavonoid subclass is catalyzed by specific enzymes; Kaempferol and quercetin, in the flavonol subclass they are considered main flavonoids. The extent of hydroxylation differentiates them from one another. These compounds are associated with the beneficial health property which reduce various kinds of cancer and are believed to have anti-oxidative activity as well as metal ion chelation and risk of coronary heart disease and lipid peroxidation inhibition (El-Hadidy et al., 2014).

Singh et al. (2009) reported that many biologically active phyto-molecules are extracted from onions which include anthocyanins, thiosulfinates, flavonoids and phenolic acids. Prakash, Singh, and Upadhyay (2007) reported that dry peel of onion (consider as a waste product) contain major flavonoids which includes quercetin glycoside which is an effective antioxidants against oxidative stress. Patricia et al. (2005) reported that vegetables and fruits are excellent sources of antioxidant phytochemicals, such as terpenoids vitamins C and E, flavonoids, vegetable pigments carotenoids, glutathione, and polyphenolics, by defending against cellular damage due to their ability to reduce oxygen-derived free radicals by donating electron, inhibit lipooxygenases and chelate to redox-active metals (Dimitrios, 2006). Free hydroxyl and peroxy radicals can be scavenged by flavonoids, on the other hand, they may extract iron ions as metal-chelating agents and reduce the superoxide-driven Fenton reaction (Suh, Lee, Cho, Kim, & Chung, 1999).

Flavonoids and flavonols are present in high levels in Allium vegetables. Flavonoid ranges from <0.03 to >1 g/kg in the edible portions of Allium vegetables. Higher levels of flavonoids 2 ± 10 g/kg are present in onion skins rather than the edible portion. The most significant flavonoids were quercetin aglycone, quercetin diglucoside and quercetin 40-glucoside and kaempferol monoglycoside or isorhamnetin monoglycoside in some cases (Leighton et al., 1992). After onion processing, a huge amount of onion skins remains useless regardless onion skin contains higher levels of flavonoids. Nuutila, Puupponen-Pimiä, Aarni, and Oksman-Caldentey (2003) reported that onion skin extracts have an anti-oxidative activity.

9. Anticancer effect

Onion consumption has inhibitory action on human carcinoma which has been broadly researched. Excess uses of onions lower the risk for carcinoma (Corzomartinez, Corzo, & Villamiel, 2007). In those individuals which have the consumption of onion is more, they had a 50% reduced risk of cancers of the alimentary, stomach and respiratory tracts (Turati, Rossi, Pelucchi, Levi, & La Vecchia, 2015). Renal and colon carcinogenesis are inhibited by organosulfur compounds such as S-methylcysteine (SMC), diallyl disulfide (DDS), and S-allylcysteine (SAC) (Fukushima, Takada, Hori, & Wanibuchi, 1997; Hatono, Jimenez, & Wargovich, 1996). Allium vegetable consumption may have a strong effect for the prevention of stomach cancer. Stomach cancer may be reduce with high consumption of onions >0.5 onion/day. In China the risk of stomach cancer was decreased by consuming allium vegetables such as onion, Chinese chives and Welsh onion more than 1 time/week (Gao, Takezaki, Ding, Li, & Tajima, 1999). The risk of *Helicobacter pylori* infection, which causes stomach cancer by ulcer formation, may be decreased by higher intake of allium vegetables. Abourehab, Khaled, Sarhan, and Ahmed (2015) reported that quercetin not only increase the cure level of ulcers with the help of free radical scavenging, but also restrict the induction of gastric mucosal injury.

Dorant, van den Brandt, Goldbohm, and Sturmans (1996) reported that the risk of lung cancer was also be analyzed through stomach carcinoma risk. Dorant, Van Din Brandt, and Goldbohm (1994) reported that those who intake higher onion will have a mild risk of lung cancer. In the onions consumption it was

found that the quantity and/or dietary absorption of flavonoids compounds and organosulfur were insufficient to produce effective results. Hence, it can be concluded that onions intake have inhibitory effect on lung carcinoma risk, but the levels needed are feasible to consume is unknown.

It is reported that tobacco smoking is major cause of urinary bladder cancer in humans (Malaveille et al., 1996). Malaveille et al. (1996, 1998) reported that human urine have dietary phenols which have an ant mutagenic effect on a known tobacco-smoke correlated carcinogen. Extraction of phenols from urine was shown to have same effects as Extracted from wine and onions, resulting that after ingestion, flavonoids have the ability to protect against tobacco carcinogens which may cause through vegetable. By *in vivo* and *in vitro* analysis the impact of onion phytochemicals can be studied on colon cancer. Hatono et al. (1996) reported that from garlic, organosulfur compound such as S-allylcysteine (SAC) was shown to hinder the precursors of colon cancer when administered orally to rats. Cancer formation has been inversely related with antioxidant activity (Newmark, 1996).

Higher intake (>16 times/week) of onions showed a significant decrease in risk of breast cancer (Challier, Pernau, & Viel, 1998; Levi, La Vecchia, Gulie, & Negri, 1993). Rodgers and Grant (1998) reported that cytotoxic carcinogenic compounds can be deactivated by enhancing the role of reductase enzyme, on the cells of breast cancer quercetin may have a substantial anti-proliferative effect. Quercetin *in vitro* study showed to have anti-proliferative effect on ovarian carcinoma cells, major cause of death in women from cancer (Shen, Herenyiova, & Weber, 1999). Rzymowska, Gawron, Pawlikowska-Pawlega, Jakubowicz-Gil, and Wojcierowski (1999) suggested that in the signal transduction pathway dose-dependent inhibition of kinase activity was found as the mode of protective action. The risk of brain cancer was positively related to salt preserved foods and inversely associated with vegetable consumption (Guo, Linet, Chow, Li, & Blot, 1994). High consumption of onions has a strong inverse relationship to cancer risk. Both Hu et al. (1999) and Guo et al. (1994) postulated that the risk of brain cancer may be associated with N-nitroso compounds (NOCs) from salted foods (Shenoy & Choughuley, 1992).

Like ROS (reactive oxygen species), RNS (reactive nitrogen species) causes oxidative damage to tissues, cellular proteins and DNA (Juurlink & Paterson, 1998). Increased risk of brain cancer was associated with (NOCs) as described by Guo et al. (1994) and Hu et al. (1999). Strong oxidant in brain, peroxynitrate has been decreased due to quercetin by hunting superoxide anions (Shutenko et al., 1999). Shenoy and Choughuley (1992) reported that researchers on N-nitroso compound formation proposed that sulphur compounds present in onion juice were the causative agents in the observed inhibition of nitrosation reactions. Sulphydryl compounds show specific inhibitory effects. Hence, onions containing both sulphur and flavonoids compounds may decrease nitrosamine formation risk and in cancer development and also possible cellular damage.

10. Bioactivity of onion and onion by-products

In a number of agro-food residues, the regaining of bioactive polyphenolic phytochemicals is one of the higher value options. In Europe A. *cepa* is commonly cultivated horticultural crop and production on annual basis is more than 5 million tons. Schieber, Stintzing, and Carle (2001) reported that on an annual basis, 450,000 tons by-products of onion are produced during consumption and processing. Biodegradable substances are mainly present in these waste materials, excess in fibre and polyphenols. Outer, non-edible dry layer of onions bulbs contain a wide-range of polyphenolic components which are removed before processing (Ly et al., 2005; Ramos et al., 2006). As these substances are absent in bulb scales, but they characterize oxidation products of its quercetin and glucosides. Ly et al. (2005) and Ramos et al. (2006) reported that phenolic compounds present in onion and onion by-products possessed strong antioxidant activity *in vitro* and have anti-platelet properties.

The vast research related to exploitation of agro-food waste materials in which the major field of concern is the extraction of polyphenols with its functional properties is just because of the day by day increasing knowledge of health relevant potential of polyphenolic phytochemicals and its major role

as food antioxidant (Schieber et al., 2001). In this concern, usage of wide range of residual sources for the recovery of natural antioxidants has been under investigation (Balasundram, Sundram, & Samman, 2006; Moure et al., 2001). In this finding, water/ethanol mixtures were investigated by non-toxic means in extraction process. For maximum recovery of total flavonol, solvent composition was kept at the optimum level. In onions, polyphenols and fructo-oligosaccharides (FOS) are good sources of bioactive compounds. Onions by-products were characterized to develop as a possibly bioactive food ingredient. The main subject of our study was to find the safety and the effects on health of by-product of onion were shared by either of two derived fractions, a fraction containing mainly materials of cell wall and an extract containing the onion polyphenols and FOS. We studied here about feeding these products to rats and their signs on gut environment, protective enzymes potential toxicity. A diet containing control feed balanced in carbohydrate were fed to rats during 4 weeks. Anaemia is caused by the onion extracts and onion by-products in rodents as expected for *Allium* products. Including genotoxicity no other toxicity was detected. When extract of onion were fed to rats, the activities of glutathione peroxidase (GPx1) and Glutathione reductase (GR) in erythrocytes was increased. Hepatic gene expression of 5-aminolevulinate synthase, NADPH, Gpx1, Gr and catalase: quinone oxidoreductase does not altered when onion is fed in any groups. By contrast, when rats given the onion residue the catalytic subunit gene expression of gamma-glutamate cysteine ligase was up regulated. The insoluble and soluble fractions as well as by-products of onion had prebiotic effects which was verified by increased butyrate production, changed gut micro biota enzyme activities and decreased pH. Roldán-Marín et al. (2009) suggested that *in vivo*, the by-products of onion may support antioxidative defense with no genotoxicity and change the functionality of the rat gut microbiota.

11. Antitumor effect

Craig (2010) reported that variety of sulfides rich in onion extracts, provide protection against tumor growth. Natural constituents of onion sodium, NPTS (*n*-propyl thiosulfate) and Two alk(en)yl thiosulfates induced hemolytic anemia as causative agents of onion. Antitumor effects of these compounds were studied by studying the beneficial functions of *n*-propyl thiosulfate (NPTS). Study shows that *in vitro*, propagation of three human tumorigenic cell lines HL-60, 293 and WiDr was inhibited. General, 2PTS as compared to NPTS have strong activity for inhibiting cell growth, not occur in WiDr cells as they are sensitive for both compounds. Oxidative damage and induced apoptosis caused by 2PTS and NPTS to HL-60 cells. Oxidative damage was approximately proportional to the extent of apoptosis and also to that of the cytotoxicity produced by these compounds. Chang, Yamato, Yamasaki, Ko, and Maede (2004) demonstrated that apoptosis induction is initiated by oxidative stress as the alk(en)yl thiosulfates have an antitumor effect.

Cell cultures of an epidermoid carcinoma cell line derived from HCPC-1 (hamster buccal pouch carcinoma) were performed by adding various onion extract concentrations *in vitro* studies. Concentration of onion extracts (25% and above) in culture media cause inhibition of tumor growth after 24 h of incubation. After incubating 4–10 days tumor proliferation was decreased. Niukian, Schwartz, and Shkla (1987) suggested that there-suits shown here give *in vitro* sign of the cytotoxic and inhibitory activity on an oral carcinoma cell line.

A bioflavonoid, quercetin, usually distributed in vegetables (onion) shows to have a chemoprotective role in cancer, through compound effects on signal transduction involved in angiogenesis and cell proliferation. The effects of quercetin supplementation (30 mg per day) were observed in the study, incorporated into a blackcurrant drink. For 14 days healthy male subjects aged between 33 and 64 years received either placebo or quercetin. At baseline Blood samples were taken and upon completion of the study and examined for full blood count, MMP-2(matrix metalloproteinase-2), TIMP-1 and -2 (tissue inhibitor of matrix metalloproteinase-1 and -2) plasma levels using ELISA techniques. For carrying out matrix metalloproteinase-2 and tissue inhibitor of matrix metalloproteinase-1 and -2 gene expression determinations, RNA was taken out from the RT-PCR (reverse transcriptase-polymerase chain reaction) and peripheral blood lymphocytes. In this study plasma protein levels of the healthy subjects or matrix metalloproteinase-2 or tissue inhibitor of matrix metalloproteinase-2 gene transcription was not changed by supplementation of the diet with quercetin. In this study the

plasma protein levels (311 ± 70 ng/ml and tissue inhibitor of matrix metalloproteinase-1 gene transcription at baseline to 183 ± 35 ng/ml post-supplementation, $p < 0.05$) of the subjects were decreased following quercetin supplementation. This is an interesting result, the functions of tissue inhibitor of matrix metalloproteinase-1 in tumour progression has some disputes. Artificially increased tissue inhibitor of matrix metalloproteinase-1 levels decrease or prevent growth of tumour in certain model systems. Yet, high levels of tissue inhibitor of matrix metalloproteinase-1 have been associated with destructive diseases and poor diagnosis in patients with certain malignancies in other studies. Morrow, Fitzsimmons, Chopra, and McGlynn (2001) proposed that a potential role for the anti-tumour promoter quercetin as a dietary mediator of the carcinogenic cascade.

In a study using CH_2Cl_2 , crude thiosulfinates are isolated from *Allium tuberosum* L. and then with silica gel column chromatography purified S-methyl 2-propene-1-thiosulfinate and S-methyl methanthiosulfinate from the crude thiosulfinates. Consequently, *in vitro* as well as *in vivo* the thiosulfinates were examined against antitumor activities and cytotoxicities against human cancer cells. Cancer cells in human have strong cytotoxicities, in the order of S-methyl methanthiosulfinate, crude thiosulfinates, and S-methyl 2-propene-1-thiosulfinate. For 7 days, when thiosulfinates consecutively administered at 10, 30, and 50 mg/kg ip, in mice, an increase in the life spans of mice occur that had been immunized with Sacoma-180 tumor cells. In MCF-7 cancer cells crude thiosulfinates also induces apoptosis. These results suggest that via apoptosis thiosulfinates from *Allium tuberosum* L. prevent the propagation of cancer cells and have antitumor activities (Park et al., 2007).

12. Need for natural antioxidant

Synthetic antioxidants (BHA and BHT) are being restricted because of their carcinogenicity e.g. they cause liver swelling and influence liver enzyme activities. Thus there is a growing interest in search for natural antioxidants for the public perception. Spices and herbs contain high amounts of compounds capable of free radical scavenging activity (Dastmalchi et al., 2008; El-Ghorab et al., 2008). In this perspective, due to its natural high content of flavonoids and its widespread popularity all over the world, the onion is an interesting vegetable plant species (Griffiths, Trueman, Crowther, Thomas, & Smith, 2002).

Allium contains pharmaceutically interesting compounds that are mainly sulfur-based compounds. Tada, Hiroe, Kiyohara, and Suzuki (1988) reported that so far, antibacterial and nematicidal activities of these compounds against *E. coli* and *Bacillus* sp. have been already investigated. Growth of root of chrysanthemum cuttings was inhibited by Allelopathic substances present in Welsh onion roots (Choi, 1993). Havey (1995) suggested that *Alliums* have been historically maintained as open pollinated populations and the growth of edible *Alliums* are worldwide and are, traded, and consumed in many countries. Numerous scientific reports suggesting that onions and its relatives can be used to cure, to lessen, or avoid some of the health problems such as asthma, antibiosis, antidiabetic, cancer, cardiovascular diseases, and prebiotic effects. Onions and its relatives contains phenolic (Fossen, Pedersen, & Andersen, 1998), minerals with antioxidant capacity (Terry, Zayed, de Souza, & Tarun, 2000), sulfur compounds (Kubec, Svobodová, & Velišek, 2000), and polysaccharides with significant prebiotic properties (Biedrzycka & Bielecka, 2004).

Lanzotti (2006) proposed that researchers assessed the antioxidant and chemical properties of various types of onions and found high concentration of kaempferol, luteolin and quercetin. Tepe, Sokmen, Akpulat, and Sokmen (2005) studied methanol extracts of five different *Allium* species except *A. cepa* red green, violet, and white Four onion scale color, compared antioxidant capacity and phenolic content, with no specified variety (Prakash et al., 2007). They suggested that kaempferol and quercetin contents are present in highest amount in red onion than other types of onion. Santas, Carbo, Gordon, and Almajano (2008) compared the polyphenol content and antioxidant. More recently, attention are diverted towards antioxidant content of onions because regular consumption of onions reduces the risk of cataract formation, inhibition of heart and vascular diseases by stopping lipid peroxidation (LPO), reduction in symptoms associated with osteoporosis (NOA), lowering of low density lipoprotein (LDL) cholesterol levels, many forms of cancer, ulcer development, and

neurodegenerative disorders (Kaneko & Baba, 1999; Kawaii, Tomono, Katase, Ogawa, & Yano, 1999; Sanderson, McLauchlan, & Williamson, 1999; Shutenko et al., 1999). Generally flavonols occur in the edible vegetables, fruits and seeds (Herrmann, 1976, 1988), and their content may reach up to a few hundred mg/kg of fresh weight (Pierpoint, 1986). Häkkinen, Kärenlampi, Heinonen, Mykkänen, and Törrönen (1999) reported that Berries such as cranberry (*Vaccinium oxycoccos* and) bog whortleberry (*Vaccinium uliginosum*) contains 263 and 184 mg/kg fresh weight of total flavonols, respectively. Still, kale (*Brassica oleracea* L. cv. Alba DC.) and onions (*A. cepa* L.) considered as excellent dietary sources, as in total flavonols their content may reach 311 and 349 mg/kg fresh weight, respectively, determined as aglycons (Hertog, Hollman, & Katan, 1992).

An integral part of the human diet is flavonols which are consumed daily by humans and for the determination of flavonol levels few efforts have been made. Total flavonoids average intake on daily basis is about 1.0 g out of which 115 mg are, the share of flavonols and flavones (Kühnau, 1976). Intake of flavonoid vary from 2.6 to 68.2 mg daily and the quercetin percentage is 39 to 100% shown by “Seven Countries Study” (Hertog et al., 1995). From fourteen countries on seventeen volunteers showed that the main use of kaempferol and quercetin is 28 mg/day approximately (De Vries, Janssen, Hollman, van Staveren, & Katan, 1997). Quercetin concentration can be lowered in tomato and onion from 35 to 82% by microwave cooking, frying and boiling which are common domestic processes particularly in flavonols (Price, Bacon, & Rhodes, 1997). It is reported that boiling and frying process show an overall loss of 25% in quercetin glucosides in case of onions and 4 α -glucoside and quercetin 3,4 α -diglucoside also show the lower level of 50% which is done by the boiling of about 25 min (Hirota, Shimoda, & Takahama, 1998). Kaempferol and quercetin level can be lowered from 39 to 64% in onion by blanching, respectively (Ewald, Fjellkner-Modig, Johansson, Sjöholm, & Åkesson, 1999). It can also be lowered in sweet potato leaves from 19 to 50% in myricetin and quercetin respectively (Chu, Chang, & Hsu, 2000). Pharmacological properties like anti-stress, anti-infection, immune enhancement potential and many more was also observed (Valko et al., 2007). In outer layer rather than purple, white and green varieties of onion the highest antioxidant and antioxidants activities in red onion and free radical scavenging activities (Prakash et al., 2007). The significance of onion is also significant by the major sources of antioxidant intake, both for specific class of flavonoids and the specific sulfur-containing compounds (Hertog, Feskens, Hollman, Katan, & Kromhout, 1993). Rather than red grape and blackberry the anthocyanin content of red onion is quite low (Clifford, 2000), with respect to flavonols their level is ~10% (Fuleki, 1971). At the same time both anthocyanins and flavonols intake is regarded as particularly healthy.

13. Conclusion

Onion and onion by-products have vital role against certain ailments due to the presence of bioactive compounds present in it. Before people did not use the by-products but now the concept of value addition is very popular. Pharmaceutically onion possesses many characteristics like anti-inflammatory, antioxidant and antibacterial properties. In decreasing pro-inflammatory diseases the effectiveness of onion is rooted in its nature as modulators of metabolism. The onion also consist of anthocyanin, flavonoids and rich source of phenolic compounds which act as antioxidant compounds. The antioxidant compounds help in control of oxidation. It is also used as condiment in different dishes. Onion has effective role against fungi and bacteria and increase the shelf life of food. It also has medicinal value and effective against different diseases. In nutshell, use of nutraceutical are better than to consume drugs against certain diseases.

Funding

The authors received no direct funding for this research.

Competing Interest

The authors declare no competing interests.

Author details

Muhammad Sajid Arshad¹
E-mail: sajid_ft@yahoo.com

Muhammad Sohaib²

E-mail: ftsohaib@yahoo.com

Muhammad Nadeem³

E-mail: mnadeem11@gmail.com

Farhan Saeed¹

E-mail: f.saeed@gcuf.edu.pk

Ali Imran¹

E-mail: aliimran.ft@gmail.com

Ahsan Javed¹

E-mail: ahsanjvi08@gmail.com

Zaid Amjad¹

E-mail: zaidamjad2012@gmail.com

ORCID ID: <http://orcid.org/0000-0002-3971-2043>

Syeda Mamoonah Batool¹

E-mail: monasyed92@gmail.com

¹ Institute of Home and Food Sciences, Government College University, Faisalabad, Pakistan.

² Department of Food Science and Human Nutrition, University of Veterinary and Animal Sciences, Lahore, Pakistan.

³ Department of Environmental Sciences, COMSATS Institute of Information Technology, Vehari, Pakistan.

Citation information

Cite this article as: Status and trends of nutraceuticals from onion and onion by-products: A critical review, Muhammad Sajid Arshad, Muhammad Sohaib, Muhammad Nadeem, Farhan Saeed, Ali Imran, Ahsan Javed, Zaid Amjad & Syeda Mamoonah Batool, *Cogent Food & Agriculture* (2017), 3: 1280254.

References

- Abourehab, M. A., Khaled, K. A., Sarhan, H. A., & Ahmed, O. A. (2015). Evaluation of combined famotidine with quercetin for the treatment of peptic ulcer: *In vivo* animal study. *Drug Design, Development and Therapy*, 9, 2159.
- Ali, M., Thomson, M., & Afzal, M. (2000). Garlic and onions: Their effect on eicosanoid metabolism and its clinical relevance. *Prostaglandins, Leukotrienes and Essential Fatty Acids (PLEFA)*, 62, 55–73.
<http://dx.doi.org/10.1054/plef.1999.0124>
- Akhtar, S. H. I. R. I. N. (2015). Advances in conventional breeding approaches for postharvest quality improvement in vegetables. *Postharvest Biology and Technology of Horticultural Crops: Principles and Practices for Quality Maintenance*, 141–176.
- Antolovich, M., Prenzler, P. D., Patsalides, E., McDonald, S., & Robards, K. (2002). Methods for testing antioxidant activity. *The Analyst*, 127, 183–198.
<http://dx.doi.org/10.1039/b009171p>
- Ashwini, M., Balaganesh, J., Balamurugan, S., Murugan, S. B., & Sathishkumar, R. (2013). Antioxidant activity in *in vivo* and *in vitro* cultures of onion varieties (Bellary and CO 3). *Food and Nutrition Sciences*, 4, 6, Article ID: 36052.
- Ayaz, E., & Alpsoy, H. C. (2007). Onion (*Allium sativum*) and traditional medicine. *Acta Parasitologica Turcica*, 31, 145–149.
- Balasundram, N., Sundram, K., & Samman, S. (2006). Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. *Food Chemistry*, 99, 191–203.
<http://dx.doi.org/10.1016/j.foodchem.2005.07.042>
- Benkeblia, N. (2005). Free-radical scavenging capacity and antioxidant properties of some selected onions (*Allium cepa* L.) and garlic (*Allium sativum* L.) extracts. *Brazilian Archives of Biology and Technology*, 48, 753–759.
<http://dx.doi.org/10.1590/S1516-89132005000600011>
- Bhattacharjee, S., Sultana, A., Sazzad, M. H., Islam, M. A., Ahtashom, M., & Asaduzzaman, M. (2013). Analysis of the proximate composition and energy values of two varieties of onion (*Allium cepa* L.) bulbs of different origin: A comparative study. *International Journal of Nutrition and Food Sciences*, 2, 246–253.
<http://dx.doi.org/10.11648/j.ijnfs.20130205.16>
- Biedrzycka, E., & Bielecka, M. (2004). Prebiotic effectiveness of fructans of different degrees of polymerization. *Trends in Food Science & Technology*, 15, 70–175.
- Block, E., Naganathan, S., Putman, D., & Zhao, S. H. (1992). Allium chemistry: HPLC analysis of thiosulfates from onion, garlic, wild garlic (Ramsoms), Chinese leek, scallion, shallot, elephant (Great-Headed) garlic, chive, and Chinese chive. Uniquely high allyl-to-methyl ratios in some garlic samples. *Journal of Agricultural and Food Chemistry*, 40, 2418–2430.
<http://dx.doi.org/10.1021/jf00024a017>
- Bors, W., & Saran, M. (1987). Radical scavenging by flavonoid antioxidants. *Free Radical Research Communications*, 2, 289–294.
<http://dx.doi.org/10.3109/10715768709065294>
- Calucci, L., Pinzino, C., Zandomenighi, M., & Capocchi, A. (2003). Effects of gamma-irradiation on the free radical and antioxidant contents in nine aromatic herbs and spices. *Journal of Agricultural and Food Chemistry*, 51, 927–934.
<http://dx.doi.org/10.1021/jf020739n>
- Cao, G., Sofic, E., & Prior, R. L. (1996). Antioxidant capacity of tea and common vegetables. *Journal of Agricultural and Food Chemistry*, 44, 3426–3431.
<http://dx.doi.org/10.1021/jf9602535>
- Cao, Y., Gu, W., Zhang, J., Chu, Y., Ye, X., Hu, Y., & Chen, J. (2013). Effects of chitosan, aqueous extract of ginger, onion and garlic on quality and shelf life of stewed-pork during refrigerated storage. *Food Chemistry*, 141, 1655–1660.
<http://dx.doi.org/10.1016/j.foodchem.2013.04.084>
- Cazzola, R., Camerotto, C., & Cestaro, B. (2011). Anti-oxidant, anti-glycant, and inhibitory activity against α -amylase and α -glucosidase of selected spices and culinary herbs. *International Journal of Food Sciences and Nutrition*, 62, 175–184. <http://dx.doi.org/10.3109/09637486.2010.529068>
- Challier, B., Pernau, J., & Viel, J. (1998). Garlic, onion and cereal fibre as protective for breast cancer: A French case study. *European Journal of Epidemiology*, 14, 739–747.
- Chang, H. S., Yamato, O., Yamasaki, M., Ka, M., & Maede, Y. (2004). Growth inhibitory effect of alk(en)yl thiosulfates derived from onion and garlic in human immortalized and tumor cell lines. *Cancer Letters*, 223, 47–55.
- Che Othman, S. F., Idid, S. O., Idid, S. Z., Koya, M. S., Mohamed Rehan, A., & Kamarudin, K. R. (2011). Antioxidant study of garlic and red onion: A Comparative study. *Pertanika Journal of Tropical Agricultural Science*, 34, 253–261.
- Choi, S. T. (1993). Materials Research Society Symposium Proceedings. *Korean Journal of Horticultural Science*, 34, 355.
- Chu, Y. H., Chang, C. L., & Hsu, H. F. (2000). Flavonoid content of several vegetables and their antioxidant activity. *Journal of the Science of Food and Agriculture*, 80, 561–566.
[http://dx.doi.org/10.1002/\(SICI\)1097-0010\(200004\)80:5<1.0.CO;2-Q](http://dx.doi.org/10.1002/(SICI)1097-0010(200004)80:5<1.0.CO;2-Q)
- Clifford, M. (2000). Anthocyanins nature occurrence and dietary burden. *Journal of the Science of Food and Agriculture*, 80, 1063–1072.
[http://dx.doi.org/10.1002/\(ISSN\)1097-0010](http://dx.doi.org/10.1002/(ISSN)1097-0010)
- Corzomartinez, M., Corzo, N., & Villamiel, M. (2007). Biological properties of onions and garlic. *Trends in Food Science & Technology*, 18, 609–625.
<http://dx.doi.org/10.1016/j.tifs.2007.07.011>
- Craig, W. (2010). *Professor of nutrition*. Berrien Springs, MI: Andrews University.
- Crozier, A., Burns, J., Aziz, A. A., Stewart, A. J., Rabsiasz, H. S., Jenkins, G. I., ... Lean, M. E. (2000). Antioxidant flavonols from fruits, vegetables and beverages: Measurement and bioavailability. *Biological Research*, 33, 79–88.
- Dastmalchi, K., Dorman, D., Oinonen, P., Darwis, Y., Laakso, I., & Hiltunen, R. (2008). Chemical composition and *in vitro* antioxidative activity of a lemon balm (*Melissa officinalis* L.) extract. *LWT - Food Science and Technology*, 41, 391–400. <http://dx.doi.org/10.1016/j.lwt.2007.03.007>
- De Vries, J. H., Janssen, M. P. L. T. M., Hollman, P. C. H., van Staveren, W. A., & Katan, M. B. (1997). Consumption of quercetin and kaempferol in free-living subjects eating a

- p variety of diets.
- Cancer Letters*
- , 114, 141–144.
-
- [http://dx.doi.org/10.1016/S0304-3835\(97\)04645-4](http://dx.doi.org/10.1016/S0304-3835(97)04645-4)
- Dimitrios, B. (2006). Sources of natural phenolic antioxidants. *Trends in Food Science & Technology*, 17, 505–512.
<http://dx.doi.org/10.1016/j.tifs.2006.04.004>
- Donner, H., Gao, L., & Mazza, G. (1997). Separation and characterization of simple and malonylated anthocyanins in red onions, *Allium cepa* L. *Food Research International*, 30, 637–643.
[http://dx.doi.org/10.1016/S0963-9969\(98\)00011-8](http://dx.doi.org/10.1016/S0963-9969(98)00011-8)
- Dorant, E., van den Brandt, P., Goldbohm, R., & Sturmans, F. (1996). Consumption of onions and a reduced risk of stomach carcinoma. *Gastroenterology*, 110, 12–20.
<http://dx.doi.org/10.1053/gast.1996.v110.pm8536847>
- Dorant, E., Van Din Brandt, P., & Goldbohm, R. (1994). A prospective cohort study on Allium vegetable consumption, garlic supplement use, and the risk of lung carcinoma in the Netherlands. *Cancer Research*, 54, 6148–6153.
- El-Ghorab, A., Gaara, A., Nassar, M., Farrag, A., Shen, H., Huq, E., & Mabry, T. (2008). Chemical constituents of clove (*Syzygium aromaticum*, Fam. Myrtaceae) and their antioxidant activity. *Zeitschrift für Lebensmittel-Untersuchung und-Forschung*, 217, 215–220.
- El-Hadidy, E. M., Mossa, M. E., & Habashy, H. N. (2014). Effect of freezing on the pungency and antioxidants activity in leaves and bulbs of green onion in Giza 6 and Photon varieties. *Annals of Agricultural Sciences*, 59, 33–39.
<http://dx.doi.org/10.1016/j.aaoas.2014.06.005>
- Ewald, C., Fjellkner-Modig, S., Johansson, K., Sjöholm, I., & Åkesson, B. (1999). Effect of processing on major flavonoids in processed onions, green beans, and peas. *Food Chemistry*, 64, 231–235.
[http://dx.doi.org/10.1016/S0308-8146\(98\)00136-8](http://dx.doi.org/10.1016/S0308-8146(98)00136-8)
- Farkaas, P., Hradsky, P., & Kovac, M. (1992). Novel flavour components identified in the steam distillate of onion (*Allium cepa* L.). *Zeitschrift für Lebensmittel-Untersuchung und-Forschung*, 195, 459–462.
<http://dx.doi.org/10.1007/BF01191718>
- Ferreres, F., Gil, M. I., & Tomás-Barberán, F. A. (1999). Anthocyanins and flavonoids from shredded red onion and changes during storage in perforated films. *Food Research International*, 29, 389.
- Fossen, T., Andersen, Ø. M., Øvstedal, D. O., Pedersen, A. T., & Raknes, Å. (1996). Characteristic anthocyanin pattern from onions and other *Allium* spp. *Journal of Food Science*, 61, 703.
<http://dx.doi.org/10.1111/jfds.1996.61.issue-4>
- Fossen, T. A., Pedersen, A. T., & Andersen, Ø. M. (1998). Flavonoids from red onion (*Allium cepa*). *Phytochemistry*, 47, 281–285.
[http://dx.doi.org/10.1016/S0031-9422\(97\)00423-8](http://dx.doi.org/10.1016/S0031-9422(97)00423-8)
- Fukushima, S., Takada, N., Hori, T., & Wanibuchi, H. (1997). Cancer prevention by organosulfur compounds from garlic and onion. *Journal of Cellular Biochemistry*, 67, 100–105.
[http://dx.doi.org/10.1002/\(ISSN\)1097-4644](http://dx.doi.org/10.1002/(ISSN)1097-4644)
- Fuleki, T. (1971). Anthocyanins in red onion, *Allium cepa*. *Journal of Food Science*, 36, 101.
<http://dx.doi.org/10.1111/jfds.1971.36.issue-1>
- Gao, C., Takezaki, T., Ding, J., Li, M., & Tajima, K. (1999). Protective effect of *Allium* vegetables against both esophageal and stomach cancer: A simultaneous case-referent study of a high-epidemic area in Jiangsu province. *Japanese Journal of Cancer Research*, 90, 614–621.
<http://dx.doi.org/10.1111/cas.1999.90.issue-6>
- Gennaro, L., Leonardi, C., Esposito, F., Salucci, M., Maiani, G., Quaglia, G., & Fogliano, V. (2002). Flavonoid and carbohydrate contents in tropea red onions: Effects of homelike peeling and storage. *Journal of Agricultural and Food Chemistry*, 50, 1904–1910.
<http://dx.doi.org/10.1021/jf011102r>
- Gey, K. F. (1990). The antioxidant hypothesis of cardiovascular disease: Epidemiology and mechanisms. *Biochemical Society Transactions*, 18, 1041–1045.
<http://dx.doi.org/10.1042/bst0181041>
- Griffiths, G., Trueman, L., Crowther, T., Thomas, B., & Smith, B. (2002). Onions—A global benefit to health. *Phytotherapy Research*, 16, 603–615.
[http://dx.doi.org/10.1002/\(ISSN\)1099-1573](http://dx.doi.org/10.1002/(ISSN)1099-1573)
- Gulsen, G., & Ayaz, E. (2010). Antimicrobial effect of onion (*Allium sativum*) and traditional medicine. *Journal of Animal and Veterinary Advances*, 9, 1680–1593.
- Guo, W., Linet, M., Chow, W., Li, J., & Blot, W. (1994). Diet and serum markers in relation to primary brain tumor risk in China. *Nutrition and Cancer*, 22, 143–150.
<http://dx.doi.org/10.1080/01635589409514339>
- Hajhashemi, V., Vaseghi, G., Pourfarzam, M., & Abdollahi, A. (2010). Are antioxidants helpful for disease prevention? *Research in Pharmaceutical Sciences*, 5(1), 1–8.
- Häkkinen, S. H., Kärenlampi, S. O., Heinonen, I. M., Mykkänen, H. M., & Törrönen, A. R. (1999). Content of the flavonols quercetin, myricetin, and kaempferol in 25 edible berries. *Journal of Agricultural and Food Chemistry*, 47, 2274–2279.
<http://dx.doi.org/10.1021/jf9811065>
- Harborne, J. B., & Mabry, T. J. (2013). *The flavonoids: Advances in research*. London: Springer.
- Harris, P. (2001). *Food supplement*. Google Patents.
- Hatono, S., Jimenez, A., & Wargovich, M. (1996). Chemopreventive effect of S-allylcysteine and its relationship to the detoxification enzyme glutathione S-transferase. *Carcinogenesis*, 17, 1041–1044.
<http://dx.doi.org/10.1093/carcin/17.5.1041>
- Havey, M. J. (1995). Onion and other cultivated alliums *Allium* spp. (Liliaceae). In J. Smartt & N. W. Simmonds (Eds.), *Evolution of crop plants* (2nd ed., pp. 344–350). Madison, WI: Longman Scientific and Technical.
- Herrmann, K. (1976). Flavonols and flavones in food plants: A review. *International Journal of Food Science & Technology*, 11, 433–448.
<http://dx.doi.org/10.1111/ifs.1976.11.issue-5>
- Herrmann, K. (1988). On the occurrence of flavonol and flavone glycosides in vegetables. *Zeitschrift für Lebensmittel-Untersuchung und-Forschung*, 186, 1–5.
<http://dx.doi.org/10.1007/BF01027170>
- Hertog, M. G., Feskens, E. J., Hollman, P. C., Katan, M. B., & Kromhout, D. (1993). Dietary antioxidant flavonoids and risk of coronary heart disease: The Zutphen Elderly Study. *The Lancet*, 342, 1007–1011.
[http://dx.doi.org/10.1016/0140-6736\(93\)92876-U](http://dx.doi.org/10.1016/0140-6736(93)92876-U)
- Hertog, M. G. L., Hollman, P. C. H., & Katan, M. B. (1992). Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *Journal of Agricultural and Food Chemistry*, 40, 2379–2383.
<http://dx.doi.org/10.1021/jf00024a011>
- Hertog, M. G. L., Kromhout, D., Aravanis, C., Blackburn, H., Buzina, R., Fidanza, F., ... Katan, M. B. (1995). Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Archives of Internal Medicine*, 155, 381–386.
<http://dx.doi.org/10.1001/archinte.1995.00430040053006>
- Hirota, S., Shimoda, T., & Takahama, U. (1998). Tissue and spatial distribution of flavonol and peroxidase in onion bulbs and stability of flavonol glucosides during boiling of the scales. *Journal of Agricultural and Food Chemistry*, 46, 3497–3502.
<http://dx.doi.org/10.1021/jf980294w>

- Hu, J., La Vecchia, C., Negri, E., Chatenoud, L., Bosetti, C., Jia, X., ... Wang, C. (1999). Diet and brain cancer in adults: A case-control study in Northeast China. *International Journal of Cancer*, 81, 20–23.
[http://dx.doi.org/10.1002/\(ISSN\)1097-0215](http://dx.doi.org/10.1002/(ISSN)1097-0215)
- Ioku, K., Aoyama, Y., Tokuno, A., Terao, J., Nakatani, N., & Takei, Y. (2001). Various cooking methods and the flavonoid content in onion. *Journal of Nutritional Science and Vitaminology*, 47, 78–83. <http://dx.doi.org/10.3177/jnsv.47.78>
- Ismailov, A. I., & Aliev, A. M. (1974). *Uchen. Zap.*, 37, 60.
- Jackson, R., McNeil, B., Taylor, C., Holl, G., Ruff, D., & Gwebu, E. T. (2002). Effect of aged onion extract on casepase-3 activity in viro. *Nutritional Neuroscience*, 5, 287–290.
<http://dx.doi.org/10.1080/10284150290032012>
- Janes, M. E., Nannapaneni, R., & Johnson, M. G. (1999). Identification and characterization of two bacteriocin-producing bacteria isolated from onion and onion by products root. *Journal of Food Protection*, 62, 899–904. <http://dx.doi.org/10.4315/0362-028X-62.8.899>
- Juurlink, B., & Paterson, P. (1998). Review of oxidative stress in brain and spinal cord injury: Suggestions for pharmacological and nutritional management strategies. *The Journal of Spinal Cord Medicine*, 21, 309–334.
<http://dx.doi.org/10.1080/10790268.1998.11719540>
- Kaneko, T., & Baba, N. (1999). Protective effect of flavonoids on endothelial cells against linoleic acid hydroperoxide-induced toxicity. *Bioscience, Biotechnology, and Biochemistry*, 63, 323–328.
<http://dx.doi.org/10.1271/bbb.63.323>
- Kawai, S., Tomono, Y., Katase, E., Ogawa, K., & Yano, M. (1999). Antiproliferative activity of flavonoids on several cancer cell lines. *Bioscience, Biotechnology, and Biochemistry*, 63, 896–899. <http://dx.doi.org/10.1271/bbb.63.896>
- Ko, M. J., Cheigh, C. I., Cho, S. W., & Chung, M. S. (2011). Subcritical water extraction of flavonol quercetin from onion skin. *Journal of Food Engineering*, 102, 327–333.
<http://dx.doi.org/10.1016/j.jfoodeng.2010.09.008>
- Krishnakantha, T. P., & Lokesh, B. R. (1993). Scavenging of superoxide anions by spice principles. *Indian Journal of Biochemistry & Biophysics*, 30, 133–134.
- Kubec, R., Svobodová, M., & Velíšek, J. (2000). Distribution of S-alk(en)ylcysteine sulfoxides in some Allium species. Identification of new flavor precursor: Sethylcysteine sulfoxide (Ethiin). *Journal of Agricultural and Food Chemistry*, 48, 428–433. <http://dx.doi.org/10.1021/jf990938f>
- Kühnau, J. (1976). The flavonoids. A class of semi-essential food components: Their role in human nutrition. In G. H. Bourne (Ed.), *World review of nutrition and dietetics* (pp. 117–191). Atlanta, GA: Karger Publishers.
- Lanzotti, V. (2006). The analysis of onion and garlic. *Journal of Chromatography A*, 1112, 3–22.
<http://dx.doi.org/10.1016/j.chroma.2005.12.016>
- Lawson, L. D., Wood, S. G., & Hughes, B. G. (1991). HPLC analysis of Allicin and other Thiosulfonates in garlic clove homogenates. *Planta Medica*, 57, 263–270.
<http://dx.doi.org/10.1055/s-2006-960087>
- Lee, J., & Mitchell, A. E. (2011). Quercetin and isorhamnetin glycosides in onion (*Allium cepa* L.): Varietal comparison, physical distribution, coproduct evaluation, and long-term storage stability. *Journal of Agricultural and Food Chemistry*, 59, 857–863.
<http://dx.doi.org/10.1021/jf1033587>
- Leighton, T., Ginther, C., Fluss, L., Harter, W. K., Cansado, J., & Notario, V. (1992). Molecular characterization of quercetin and quercetin glycosides in Allium vegetables: Their effects on malignant cell transformation. In ACS symposium series (USA).
- Levi, F., La Vecchia, C., Gulie, C., & Negri, E. (1993). Dietary factors and breast cancer risk in Vaud, Switzerland. *Nutrition and Cancer*, 19, 327–335.
<http://dx.doi.org/10.1080/01635589309514263>
- Lombard, K., Peffley, E., Geoffria, E., Thompson, L., & Herring, A. (2005). Quercetin in onion (*Allium cepa* L.) after heat-treatment simulating home preparation. *Journal of Food Composition and Analysis*, 18, 571–581.
<http://dx.doi.org/10.1016/j.jfca.2004.03.027>
- Ly, T. N., Hazama, C., Shimoyamada, M., Ando, H., Kato, K., & Yamauchi, R. (2005). Antioxidative compounds from the outer scales of onion. *Journal of Agricultural and Food Chemistry*, 53, 8183–8189.
<http://dx.doi.org/10.1021/jf051264d>
- Malaveille, C., Hautefeuille, A., Pignatelli, B., Talaska, G., Vineis, P., & Bartsch, H. (1996). Dietary phenolics as anti-mutagens and inhibitors of tobacco-related DNA adduction in the urothelium of smokers. *Carcinogenesis*, 17, 2193–2200.
<http://dx.doi.org/10.1093/carcin/17.10.2193>
- Malaveille, C., Hautefeuille, A., Pignatelli, B., Talaska, G., Vineis, P., & Bartsch, H. (1998). Antimutagenic dietary phenolics as antigenotoxic substances in urothelium of smokers. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 402, 219–224.
[http://dx.doi.org/10.1016/S0027-5107\(97\)00300-X](http://dx.doi.org/10.1016/S0027-5107(97)00300-X)
- Meena, M. R. (1998). *Studies on antimicrobial activity of various spices and their oils* (MSc Thesis). Indian Agricultural Research Institute, New Delhi.
- Mollavali, M., Bolandnazar, S. A., Schwarz, D., Rohn, S., Riehle, P., & Zaare Nahandi, F. (2016). Flavonol glucoside and antioxidant enzyme biosynthesis affected by mycorrhizal fungi in various cultivars of onion (*Allium cepa* L.). *Journal of Agricultural and Food Chemistry*, 64, 71–77.
<http://dx.doi.org/10.1021/acs.jafc.5b04791>
- Morrow, D. M. P., Fitzsimmons, P. E. E., Chopra, M., & McGlynn, H. (2001). Dietary supplementation with the anti-tumour promoter quercetin: Its effects on matrix metalloproteinase gene regulation. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 480–481, 269–276.
[http://dx.doi.org/10.1016/S0027-5107\(01\)00184-1](http://dx.doi.org/10.1016/S0027-5107(01)00184-1)
- Moure, A., Cruz, J. M., Franco, D., Domínguez, J. M., Sineiro, J., Domínguez, H., ... Parajó, J. C. (2001). Natural antioxidants from residual sources. *Food Chemistry*, 72, 145–171.
[http://dx.doi.org/10.1016/S0308-8146\(00\)00223-5](http://dx.doi.org/10.1016/S0308-8146(00)00223-5)
- Mushtaq, S., Amjad, M., Ziaf, K., Cheema, K. L., Raza, M. A., & Hafeez, O. B. A. (2013). Productive and qualitative evaluation of onion cultivars under agro-climatic conditions of Faisalabad. *Pakistan Journal of Agricultural Sciences*, 50, 199–203.
- Nemeth, K., & Piskula, M. (2007). Food content, processing, absorption and metabolism of onion flavonoids. *Critical Reviews in Food Science and Nutrition*, 47, 397–409.
<http://dx.doi.org/10.1080/10408390600846291>
- Newmark, H. L. (1996). Plant phenolics as potential cancer prevention agents. In *Dietary phytochemicals in cancer prevention and treatment. Edited under the auspices of the American Institute for Cancer Research* (chap. 3, pp. 25–34). New York, NY: Plenum Press.
- Niukian, K., Schwartz, J., & Shkila, J. (1987). In vitro inhibitory effect of onion extract on hamster buccal pouch. *Carcinogenesis*, 10, 137–144.
- Nuutila, A. M., Puupponen-Pimiä, R., Aarni, M., & Oksman-Caldentey, K. M. (2003). Comparison of antioxidant activities of onion and garlic extracts by inhibition of lipid peroxidation and radical scavenging activity. *Food Chemistry*, 81, 485–493.
[http://dx.doi.org/10.1016/S0308-8146\(02\)00476-4](http://dx.doi.org/10.1016/S0308-8146(02)00476-4)
- Onyeoziri, U. P., Romanus, E. N., & Onyekachukwu, U. I. (2016). Assessment of antioxidant capacities and phenolic contents of Nigerian cultivars of onions (*Allium cepa* L.) and garlic (*Allium sativum* L.). *Pakistan Journal of Pharmaceutical Sciences*, 29, 1183–1188.

- Ostrowska, E., Gabler, N. K., Sterling, S. J., Tatham, B. G., Jones, R. B., & Eagling, D. R. (2004). Consumption of brown onions (*Allium cepa* var. cavalier and var. destiny) moderately modulates blood lipids, haematological and haemostatic variables in healthy pigs. *British Journal of Nutrition*, 91, 211–218.
<http://dx.doi.org/10.1079/BJN20031036>
- Park, K. W., Kim, S. Y., Jeong, Il-Yun, Byun, M. W., Park, K. H., Yamada, K., & Seo, K. (2007). Cytotoxic and antitumor activities of thiosulfates from *Allium tuberosum* L. *Journal of Agricultural and Food Chemistry*, 55, 7957–7961. <http://dx.doi.org/10.1021/jf0713051>
- Patricia, I., Oteiza, A. G., Erlejan, S., Verstraeten, V., Keen, C. L., & Fraga, C. S. (2005). Flavonoid-membrane interactions: A protective role of flavonoids at the membrane surface. *Clinical and Developmental Immunology*, 12, 23–25.
- Pérez-Gregorio, M., Regueiro, J., González-Barreiro, C., Rial-Otero, R., & Simal-Gándara, J. (2011). Changes in antioxidant flavonoids during freeze-drying of red onions and subsequent storage. *Food Control*, 22, 1108–1113. <http://dx.doi.org/10.1016/j.foodcont.2011.01.006>
- Pierpoint, W. S. (1986). Flavonoids in the human diet. In V. Cody, E. J. Middleton, & Jeffrey B. Harborne (Eds.), *Plant flavonoids in biology and medicine: Biochemical, pharmacological, and structure-activity relationships* (pp. 125–140). New York, NY: Allan R. Liss.
- Prakash, D., Singh, B. N., & Upadhyay, G. (2007). Antioxidant and free radical scavenging activities of phenols from onion (*Allium cepa*). *Food Chemistry*, 102, 1389–1393. <http://dx.doi.org/10.1016/j.foodchem.2006.06.063>
- Price, K. R., Bacon, J. R., & Rhodes, M. J. C. (1997). Effect of storage and domestic processing on the content and composition of flavonol glucosides in onion (*Allium cepa*). *Journal of Agricultural and Food Chemistry*, 45, 938–942. <http://dx.doi.org/10.1021/jf9605916>
- Ramos, F. A., Takaishi, Y., Shirotori, M., Kawaguchi, Y., Tsuchiya, K., Shibata, H., ... Takeuchi, M. (2006). Antibacterial and antioxidant activities of quercetin oxidation products from yellow onion (*Allium cepa*) skin. *Journal of Agricultural and Food Chemistry*, 54, 3551–3557. <http://dx.doi.org/10.1021/jf060251c>
- Reddy, A. C., & Lokesh, B. R. (1992). Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. *Molecular and Cellular Biochemistry*, 111, 117–124.
- Rodgers, E. H., & Grant, M. H. (1998). The effect of the flavonoids, quercetin, myricetin and epicatechin on the growth and enzyme activities of MCF7 human breast cancer cells. *Chemico-Biological Interactions*, 116, 213–228. [http://dx.doi.org/10.1016/S0009-2797\(98\)00092-1](http://dx.doi.org/10.1016/S0009-2797(98)00092-1)
- Rohn, S., Buchner, N., Driemel, G., Rauser, M., & Kroh, L. W. (2007). Thermal degradation of onion quercetin glucosides under roasting conditions. *Journal of Agricultural and Food Chemistry*, 55, 1568–1573. <http://dx.doi.org/10.1021/jf063221i>
- Roldán, E., Sánchez-Moreno, C., de Ancos, B., & Cano, M. (2008). Characterisation of onion (*Allium cepa* L.) by-products as food ingredients with antioxidant and antibrowning properties. *Food Chemistry*, 108, 907–916. <http://dx.doi.org/10.1016/j.foodchem.2007.11.058>
- Roldán-Marín, E., Krath, B. N., Poulsen, M., Binderup, M. L., Nielsen, T. H., Hansen, M., ... Dragsted, L. O. (2009). Effects of an onion by-product on bioactivity and safety markers in healthy rats. *British Journal of Nutrition*, 102, 1574–1582. <http://dx.doi.org/10.1017/S0007114509990870>
- Rzymowska, J., Gawron, A., Pawlikowska-Pawlega, B., Jakubowicz-Gil, J., & Wojcickowski, J. (1999). The effect of quercetin on induction of apoptosis. *Folia histochemica et cytobiologica/Polish Academy of Sciences, Polish Histochemical and Cytochemical Society*, 37, 125.
- Sanderson, J., McLauchlan, W., & Williamson, G. (1999). Quercetin inhibits hydrogen peroxide-induced oxidation of the rat lens. *Free Radical Biology and Medicine*, 26, 639–645. [http://dx.doi.org/10.1016/S0891-5849\(98\)00262-7](http://dx.doi.org/10.1016/S0891-5849(98)00262-7)
- Santas, J., Carbo, R., Gordon, M. H., & Almajano, M. P. (2008). Comparison of the antioxidant activity of two Spanish onion varieties. *Food Chemistry*, 107, 1210–1216. <http://dx.doi.org/10.1016/j.foodchem.2007.09.056>
- Schieber, A., Stintzing, F. C., & Carle, R. (2001). By-products of plant food processing as a source of functional compounds—Recent developments. *Trends in Food Science & Technology*, 12, 401–413. [http://dx.doi.org/10.1016/S0924-2244\(02\)00012-2](http://dx.doi.org/10.1016/S0924-2244(02)00012-2)
- Shen, F., Herenyiova, M., & Weber, G. (1999). Synergistic down-regulation of signal transduction and cytotoxicity by tiazofurin and quercetin in human ovarian carcinoma cells. *Life Sciences*, 64, 1869–1876. [http://dx.doi.org/10.1016/S0024-3205\(99\)00133-2](http://dx.doi.org/10.1016/S0024-3205(99)00133-2)
- Shenoy, N., & Choughuley, A. (1992). Inhibitory effect of diet related sulphhydryl compounds on the formation of carcinogenic nitrosamines. *Cancer Letters*, 65, 227–232. [http://dx.doi.org/10.1016/0304-3835\(92\)90236-0](http://dx.doi.org/10.1016/0304-3835(92)90236-0)
- Shutenko, Z., Henry, Y., Pinard, E., Seylaz, J., Potier, P., Berthet, F., Girard, P., & Sercombe, R. (1999). Influence of the antioxidant quercetin *in vivo* on the level of nitric oxide determined by electron paramagnetic resonance in rat brain during global ischemia and reperfusion. *Biochemical Pharmacology*, 57, 199–208. [http://dx.doi.org/10.1016/S0006-2952\(98\)00296-2](http://dx.doi.org/10.1016/S0006-2952(98)00296-2)
- Siddiq, M., Roidoung, S., Sogi, D. S., & Dolan, K. D. (2013). Total phenolics, antioxidant properties and quality of fresh-cut onions (*Allium cepa* L.) treated with mild-heat. *Food Chemistry*, 136, 803–806. <http://dx.doi.org/10.1016/j.foodchem.2012.09.023>
- Singh, B. N., Singh, B., Singh, R., Prakash, D., Singh, D., Sarma, B., ... Singh, H. (2009). Polyphenolics from various extracts/fractions of red onion (*Allium cepa*) peel with potent antioxidant and antimutagenic activities. *Food and Chemical Toxicology*, 47, 1161–1167. <http://dx.doi.org/10.1016/j.fct.2009.02.004>
- Stajner, D., & Varga, I. S. (2003). An evaluation of the antioxidant abilities of *Allium* species. *Acta Biologica Szegediensis*, 47, 103–106.
- Suh, H. J., Lee, J. M., Cho, J. S., Kim, Y. S., & Chung, S. H. (1999). Radical scavenging compounds in onion skin. *Food Research International*, 32, 659–664. [http://dx.doi.org/10.1016/S0963-9969\(99\)00141-6](http://dx.doi.org/10.1016/S0963-9969(99)00141-6)
- Suleria, H. A. R., Butt, M. S., Anjum, F. M., Saeed, F., & Khalid, N. (2013). Onion: Nature protection against physiological threats. *Critical Reviews in Food Science and Nutrition*, 1–17.
- Swenson, K. J. (2008). *U.S. Patent Application No. 12/053,975*.
- Tada, M., Hiroe, Y., Kiyohara, S., & Suzuki, S. (1988). Nematicidal and antimicrobial constituents from *Allium grayi* Regel. and *Allium fistulosum* L. var. caespitosum. *Agricultural and Biological Chemistry*, 52, 2381–2385.
- Tepe, B., Sokmen, M., Akpulat, H. A., & Sokmen, A. (2005). In vitro antioxidant activities of the methanol extracts of five species from Turkey. *Food Chemistry*, 92, 89–92. <http://dx.doi.org/10.1016/j.foodchem.2004.07.016>
- Terry, N., Zayed, A. M., de Souza, M. P., & Tarun, A. S. (2000). Selenium in higher plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 51, 401–432. <http://dx.doi.org/10.1146/annurev.arplant.51.1.401>
- Tram Ngoc, L., Chiharu, H., Makoto, M., Hiromune, A., Koji, K., & Ryo, Y. (2005). Antioxidative compounds from the outer scales of onion. *Journal of Agricultural and Food Chemistry*, 53, 8183–8189.
- Turati, F., Rossi, M., Pellicchi, C., Levi, F., & La Vecchia, C. (2015). Fruit and vegetables and cancer risk: A review of southern European studies. *British Journal of Nutrition*, 113(S2), S102–S110.

- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry & Cell Biology*, 39, 44–84. <http://dx.doi.org/10.1016/j.biocel.2006.07.001>
- van Wyk, B. E. (2014). *Culinary herbs and spices of the world*. Chicago, IL: University of Chicago Press. <http://dx.doi.org/10.7208/chicago/9780226091839.001.0001>
- White, K., & Zellner, J. (2008). *In college seminar 235 food for thought: The science, culture, & politics of food spring*.
- Wiczowski, W., Nemeth, K., Bucin'ski, A., & Piskula, M. K. (2003). Bioavailability of quercetin from flesh scales and dry skin of onion in rats. *Polish Journal of Food and Nutrition Sciences*, 12, 95–99.
- Willett, W. (1994). Diet and health: What should we eat? *Science*, 264, 532–537. <http://dx.doi.org/10.1126/science.8160011>
- Yang, G. C., Yasaei, P. M., & Page, S. W. (1993). Garlic as anti-oxidants and free radical scavengers. *Journal of Food and Drug Analysis*, 1, 357–364.
- Yang, J., Meyers, K. J., van der Heide, J., & Liu, R. H. (2004). Varietal differences in phenolic content and antioxidant and antiproliferative activities of onions. *Journal of Agricultural and Food Chemistry*, 52, 6787–6793. <http://dx.doi.org/10.1021/jf0307144>
- Zakarova, A., Seo, J. Y., Kim, H. Y., Kim, J. H., Shin, J. H., Cho, K. M., ... Kim, J. S. (2014). Garlic sprouting is associated with increased antioxidant activity and concomitant changes in the metabolite profile. *Journal of Agricultural and Food Chemistry*, 62, 1875–1880. <http://dx.doi.org/10.1021/jf500603v>



© 2017 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format
 Adapt — remix, transform, and build upon the material for any purpose, even commercially.
 The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.
 You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
 No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



***Cogent Food & Agriculture* (ISSN: 2331-1932) is published by Cogent OA, part of Taylor & Francis Group.**

Publishing with Cogent OA ensures:

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

Submit your manuscript to a Cogent OA journal at www.CogentOA.com

