



Comparison of the systemic bioavailability of mometasone furoate after oral inhalation from a mometasone furoate/formoterol fumarate metered-dose inhaler versus a mometasone furoate dry-powder inhaler in patients with chronic obstructive pulmonary disease

Teddy Kosoglou, James Hubbell, Fengjuan Xuan, David L Cutler, Alan G Meehan, Bhavna Kantesaria & Bret A Wittmer

To cite this article: Teddy Kosoglou, James Hubbell, Fengjuan Xuan, David L Cutler, Alan G Meehan, Bhavna Kantesaria & Bret A Wittmer (2013) Comparison of the systemic bioavailability of mometasone furoate after oral inhalation from a mometasone furoate/formoterol fumarate metered-dose inhaler versus a mometasone furoate dry-powder inhaler in patients with chronic obstructive pulmonary disease, International Journal of Chronic Obstructive Pulmonary Disease, , 107-116, DOI: [10.2147/COPD.S36592](https://doi.org/10.2147/COPD.S36592)

To link to this article: <https://doi.org/10.2147/COPD.S36592>



© 2013 Kosoglou et al, publisher and licensee Dove Medical Press Ltd.



Published online: 04 Mar 2013.



Submit your article to this journal [↗](#)



Article views: 112



View related articles [↗](#)

Comparison of the systemic bioavailability of mometasone furoate after oral inhalation from a mometasone furoate/formoterol fumarate metered-dose inhaler versus a mometasone furoate dry-powder inhaler in patients with chronic obstructive pulmonary disease

Teddy Kosoglou¹

James Hubbell²

Fengjuan Xuan³

David L Cutler¹

Alan G Meehan⁴

Bhavna Kantesaria⁵

Bret A Wittmer^{6,†}

¹Clinical Pharmacology, ²Exploratory Drug Metabolism, ³Early Development Statistics, ⁴Medical Communications, ⁵Drug Metabolism/Pharmacokinetics, Merck Sharp & Dohme Corp, Whitehouse Station, NJ, USA;

⁶Commonwealth Biomedical Research, LLC, Madisonville, KY, USA

[†]Dr Bret A Wittmer passed away on May 9, 2012.

Background: Coadministration of mometasone furoate (MF) and formoterol fumarate (F) produces additive effects for improving symptoms and lung function and reduces exacerbations in patients with asthma and chronic obstructive pulmonary disease (COPD). The present study assessed the relative systemic exposure to MF and characterized the pharmacokinetics of MF and formoterol in patients with COPD.

Methods: This was a single-center, randomized, open-label, multiple-dose, three-period, three-treatment crossover study. The following three treatments were self-administered by patients ($n = 14$) with moderate-to-severe COPD: MF 400 μ g/F 10 μ g via a metered-dose inhaler (MF/F MDI; DULERA®/ZENHALE®) without a spacer device, MF/F MDI with a spacer, or MF 400 μ g via a dry-powder inhaler (DPI; ASMANEX® TWISTHALER®) twice daily for 5 days. Plasma samples for MF and formoterol assay were obtained predose and at prespecified time points after the last (morning) dose on day 5 of each period of the crossover. The geometric mean ratio (GMR) as a percent and the corresponding 90% confidence intervals (CI) were calculated for treatment comparisons.

Results: Systemic MF exposure was lower (GMR 77%; 90% CI 58, 102) following administration by MF/F MDI compared to MF DPI. Additionally, least squares geometric mean systemic exposures of MF and formoterol were lower (GMR 72%; 90% CI 61, 84) and (GMR 62%; 90% CI 52, 74), respectively, following administration by MF/F MDI in conjunction with a spacer compared to MF/F MDI without a spacer. MF/F MDI had a similar adverse experience profile as that seen with MF DPI. All adverse experiences were either mild or moderate in severity; no serious adverse experience was reported.

Conclusion: Systemic MF exposures were lower following administration by MF/F MDI compared with MF DPI. Additionally, systemic MF and formoterol exposures were lower following administration by MF/F MDI with a spacer versus without a spacer. The magnitude of these differences with respect to systemic exposure was not clinically relevant.

Keywords: mometasone furoate, chronic obstructive pulmonary disease, pharmacokinetics, systemic exposure, metered-dose inhaler, dry-powder inhaler

Correspondence: Teddy Kosoglou
Clinical Pharmacology, Merck Sharp & Dohme Corp, 351 N Sumneytown Pike, UG4-D48, North Wales, PA 19454, USA
Tel +1 267 305 3187
Fax +1 908 259 2305
Email teddy.kosoglou2@merck.com

Introduction

Current treatment guidelines for the long-term management of chronic obstructive pulmonary disease (COPD) and asthma recommend, for certain degrees of severity,

combination therapy with an inhaled corticosteroid (ICS) and a long-acting β_2 -agonist.¹⁻⁵ Clinically, coadministration of an ICS and long-acting β_2 -agonist has been shown to have additive effects for improving symptoms and lung function and reducing the frequency of disease exacerbations.¹⁻⁵

Mometasone furoate (MF) is a potent ICS with relatively low potential to cause significant systemic side effects typically associated with oral corticosteroids, such as hypothalamic-pituitary-adrenal axis suppression.⁶⁻⁸ MF has been shown to produce clinical benefit for treating asthma and COPD by reducing symptoms and exacerbations, and improving lung function, with no significant safety risks.⁶⁻¹⁰ Formoterol fumarate (F) is a potent, selective, long-acting β_2 -agonist that exerts a preferential effect on β_2 -adrenergic receptors of bronchial smooth muscle.¹¹ Bronchodilator activity observed in patients with asthma after F inhalation is characterized by a rapid onset (within 3 minutes of inhalation) and long duration (at least 12 hours) of action.¹¹ F is approved for maintenance treatment in patients with asthma and COPD. Merck Sharp & Dohme Corp (Whitehouse Station, NJ, USA) and Novartis (East Hanover, NJ, USA) have jointly developed a fixed-dose combination product combining MF and F in a metered-dose inhaler device (MF/F MDI; marketed as DULERA® in the United States and ZENHALE® in Canada and elsewhere; Merck & Co. Inc., Whitehouse Station, NJ, USA) for the treatment of asthma. MF/F MDI is also in late-stage clinical development for the treatment of patients with COPD. In addition to producing additive beneficial effects on symptoms and lung function, an MF/F combination product is expected to be more convenient for patients with asthma or COPD.

Due to the effects of decreases in lung function on exposure to inhaled products in patients with COPD, the systemic exposure and pharmacokinetics of MF and formoterol in these patients were expected to differ from healthy volunteers, as has been reported for other ICSs.^{12,13} A previous pharmacokinetic study showed lower mean (area under the curve [AUC]; 25%; geometric mean ratio [GMR]: 75%; 90% confidence interval [CI]: 61%–91%; mean maximum concentration [C_{max}] 39% [GMR: 61%; 90% CI: 49%–75%]) systemic exposure of MF after steady-state dosing from the MDI compared to the dry-powder inhaler (DPI) ASMANEX® TWISTHALER® ([MF] Merck Sharp & Dohme Corp) in healthy patients (data on file, Merck Sharp & Dohme Corp, 2010). Similar differences in systemic drug exposure with MDIs and DPIs have been reported for other ICSs.^{14,15}

As part of the clinical development program for MF/F MDI for the treatment of patients with moderate to severe

COPD, the present study was conducted primarily to define the systemic exposure of MF when administered using a new combination product containing MF and F in an MDI device (MF/F MDI) versus the approved DPI monotherapy product (MF DPI) for which there is extensive clinical use and safety experience.¹⁶ As a secondary objective, this study examined the potential effect of using a spacer device in conjunction with the MF/F MDI on MF and formoterol exposure (versus use of MDI device without a spacer). In addition, this study provided descriptive multiple-dose pharmacokinetic data for MF and formoterol in patients with COPD.

Methods

Patient selection

This study enrolled men and women between the ages of 40 and 75 years with the following inclusion criteria: moderate to severe COPD (as defined by post-bronchodilator forced expiratory volume in 1 second [FEV_1] $\geq 30\%$ and $< 80\%$) within 1 week prior to the baseline visit (day 1); current smoker or ex-smoker with at least 10 pack-years of smoking history; and receiving only albuterol/salbutamol for relief of symptoms for at least 2 weeks prior to randomization. Subjects were excluded from participation in this study based on the following criteria: increase in absolute volume of FEV_1 of ≥ 400 mL within 30 minutes after administration of four inhalations of albuterol/salbutamol (total dose of 360 to 400 μ g), or nebulized 2.5 mg albuterol/salbutamol; inability to use the MF/F MDI device or the MF DPI device; female patients who were pregnant, intended to become pregnant (within 3 months of ending the study), or were breastfeeding; history of any infectious disease within 4 weeks prior to drug administration; or tested positive for hepatitis B surface antigen, hepatitis C antibodies, or human immunodeficiency virus.

Study design

This was a randomized, open-label, multiple-dose, three-period, three-treatment crossover study conducted at a single study center. Subjects were screened within 21 days prior to dosing.

All patients were trained in the use of the devices and proper inhalation techniques using placebo, MDI, and DPI. If necessary, patients could be retrained in the proper use of these inhaler devices prior to the start of each period. Subjects were instructed by the investigator regarding when and how to take the daily treatment.

Subjects were admitted to the study center on day 1 to confirm continued eligibility and for baseline assessments. The investigator or designee reviewed the inclusion/exclusion

criteria and recorded adverse events (AEs) and medications taken within the previous 14 days. A repeat drug and pregnancy screen, laboratory safety tests (hematology, blood chemistry, urinalysis, and electrocardiography), and vital signs also were performed on day 1. On day 1 of the first treatment period, after a 10-hour overnight fast, each patient was randomized to a crossover treatment sequence according to a computer-generated randomization schedule provided by the sponsor, and then received the first dose. The following three treatments were self-administered by patients: treatment A, MF 400 µg/F 10 µg twice a day (BID) via MDI oral inhalation (two puffs × 200 µg/5 µg MF/F per burst combination product); treatment B, MF 400 µg/F 10 µg BID via MDI oral inhalation and in conjunction with a spacer device (two puffs × 200 µg/5 µg MF/F per burst combination product); and treatment C, MF 400 µg BID via DPI oral inhalation (two puffs × 200 µg MF per oral inhalation from the MF DPI). Subjects self-administered the treatments under observation of the site staff for 4 days every 12 hours between approximately 8 am and 9 am and again between approximately 8 pm and 9 pm, and a single morning dose on day 5. After taking their treatment, subjects were instructed to rinse their mouth with water and then spit it out (not swallow it). Based on a previous pharmacokinetic study where the effective half-life (based on accumulation) of MF after administration from an MDI was approximately 25 hours, and since dosing for five half-lives is typically required to attain steady state conditions, a dosing period of 5 days was chosen for this study. Subjects were confined to the study center for the duration of treatment in each period of the crossover. After a 1-week washout between dosing, patients returned to start confinement for the next treatment period.

MF/F MDIs and placebo MDI devices (for the training of the inhalation technique) were manufactured by 3M Health Care Ltd (Loughborough, Leicestershire, UK, for Schering-Plough Corp [now Merck Sharp & Dohme Corp], Kenilworth, NJ, USA). The spacer device (AeroChamber Plus® Valved Holding Chamber; Monaghan Medical Corp, Plattsburgh, NY, USA) and MF DPI were obtained commercially by the site. Placebo DPI to match the MF DPI was manufactured and supplied by Schering-Plough Corp (now Merck Sharp & Dohme Corp).

The study population was identified from the surrounding urban and suburban communities of Madisonville, KY, USA. Subjects were recruited from a pool of volunteers obtained from the database of Commonwealth Biomedical Research, LLC and by word-of-mouth advertisement. The protocol and informed consent were approved by Independent

Investigational Review Board, Plantation, FL, USA, as required by the US Code of Federal Regulations and the internal Standard Operating Procedures of the sponsor. The study was conducted in accordance with good clinical practice and was approved by the appropriate institutional review boards and regulatory agencies. Written consent was obtained from all patients prior to the conduct of any study related procedures.

Pharmacokinetic assessments

Pharmacokinetic parameters were calculated via noncompartmental analysis. Day 5 plasma concentrations and actual sampling times were used to calculate the following pharmacokinetic parameters of MF and formoterol: area under the plasma concentration–time curve from 0 to 12 hours ($AUC_{0-12\text{ hr}}$), the maximum plasma concentration (C_{max}), trough plasma concentration (C_{trough}), and time to maximum plasma concentration (T_{max}). Pharmacokinetic parameters were calculated using WinNonlin® software (v 5.0.1; Pharsight Corporation, Mountain View, CA, USA). $AUC_{0-12\text{ hr}}$ was calculated using the linear trapezoidal method for ascending concentrations and log trapezoidal method for descending concentrations. Values for C_{max} , C_{trough} , and T_{max} were obtained by visual inspection of the blood concentration data.

For the determination of MF plasma concentration, whole blood was collected in K3-ethylenediaminetetraacetic acid-containing tubes at predose (0 hours) and at 0.5, 1, 2, 4, 8, and 12 hours after the morning dose on day 5 and centrifuged for 15 minutes at 1500 × g. Plasma was removed and stored in a freezer at −20°C. A 1 mL sample aliquot was fortified with 50 µL of internal standard (mometasone furoate-d3). Analytes were isolated through liquid–liquid extraction with 5.0 mL of 15:85 ethyl acetate/hexane, v/v. The extracts were further purified by solid phase extraction with Bond Elut® LRC NH2 cartridges (Agilent Technologies, Strathaven, Scotland). Analytes were eluted with 6.0 mL of 65:35 ethyl acetate/hexane, v/v. The solvent was evaporated under a nitrogen stream at approximately 50°C, and the remaining residue was reconstituted with 125 µL of methanol and 75 µL of 20 µM sodium acetate. The final extract was analyzed using a Sciex API 5000 triple quadrupole liquid chromatography with tandem mass spectrometry system equipped with an electrospray ionization source (AB SCIEX, Framingham, MA, USA) and having a lower limit of quantitation (LLOQ) of 0.250 pg/mL. The LC system employed a Luna (Phenomenex, Torrance, CA, USA) C18 3 × 150 mm (3 µm particle size) column and gradient elution. The retention time for both MF and the internal standard was approximately 14 minutes. The range of

the standard curve using a 1.00 mL sample of human plasma was 0.250 to 25.0 pg/mL. At the LLOQ (0.250 pg/mL) for the MF assay, the between-day mean (standard deviation) was 0.253 (0.023) pg/mL, mean % bias was 1.10%, and the mean coefficient of variation was 8.95%.

For the determination of the plasma concentration of formoterol, whole blood was collected in tubes containing lithium heparin with eserine (physostigmine) hemisulfate as a preservative at predose (0 hours) and at 0.167 (10 minutes), 0.25, 0.5, 1, 2, 4, 8, and 12 hours after the morning dose on day 5 and centrifuged for 15 minutes at $1500 \times g$. Plasma was removed and stored in a freezer at -20°C . A 500 μL plasma sample aliquot was fortified with 20 μL of internal standard (formoterol- d_6) and 200 μL of 2% ammonium hydroxide. Extraction solvent was added, and the tubes were vortexed and centrifuged. The aqueous layer was frozen and the organic layer was decanted to a clean tube containing keeper solution. The organic solution was evaporated and the remaining residue was reconstituted with 200 μL of reconstitution solution. A 40- μL volume of the final extract was injected and analyzed using a Sciex API 5000 triple quadrupole liquid chromatography with tandem mass spectrometry system equipped with a turbo ion spray source (AB SCIEX) and having an LLOQ of 1.45 pmol/L. The LC system employed a 10×3 mm (5 μm particle size) Thermo BETASIL Silica-100 loading column (Thermo Fisher Scientific, Waltham, MA, USA) and a 50×3 mm (5 μm particle size) Thermo BETASIL Silica-100 analytical column. Formoterol and the internal standard were separated from the other plasma components using a mobile phase A consisting of 0.1% formic acid in 10 mM ammonium formate and a mobile phase B consisting of a 95:5 acetonitrile:mobile Phase A. The retention time for both formoterol and the internal standard were approximately 2.5 minutes. The range of the standard curve using a 500 μL sample of human plasma was 1.45 pmol/L to 727 pmol/L. At the LLOQ (1.45 pmol/L) for the formoterol assay, the between-day mean (standard deviation) was 1.44 (0.135) pmol/L, mean % bias was -1.23% , and the mean % coefficient of variation was 9.37%.

Safety assessments

Clinical laboratory tests, vital signs, electrocardiography, and physical examinations were assessed at screening and clinical laboratory tests and vital signs at prespecified times during the study. Subjects were continually monitored for possible occurrence of AEs. At study conclusion (day 6 of period 3), vital signs, clinical laboratory tests, and physical examinations were repeated.

Statistical analysis

Summary statistics including means and coefficients of variation were provided for MF concentration data at each time point and the derived pharmacokinetic parameters. The primary objective was to compare the MF $\text{AUC}_{0-12 \text{ hr}}$ and C_{max} values for the MF/F MDI combination with those for MF DPI monotherapy (ie, treatment A versus treatment C, respectively). The $\text{AUC}_{0-12 \text{ hr}}$ and C_{max} values were log-transformed and analyzed using an appropriate analysis of variance (ANOVA) model for a three-period crossover design extracting sources of variation due to treatment, patient, sequence, and period. As a secondary objective, the MF $\text{AUC}_{0-12 \text{ hr}}$ and C_{max} values for the MF/F MDI combination administered without and with a spacer were compared (ie, treatment A versus treatment B). The geometric means of treatment A to treatment C or treatment A to treatment B were expressed as a ratio (GMR), and the corresponding 90% CIs were computed. In addition, the log-transformed $\text{AUC}_{0-12 \text{ hr}}$ and C_{max} values for formoterol were analyzed similarly for treatments A and B, and comparisons between treatment A and treatment B were performed using a ratio and the corresponding 90% CI.

Assuming an inpatient variability of 28% (based on the variability observed in a similarly designed crossover study in healthy adults), this study with a targeted sample size of 12 patients should have been able to detect a 30% difference in MF pharmacokinetics between treatments B or C versus A with 80% power and a 90% CI.

No inferential analysis of safety data was planned. The number of patients reporting any AEs, the occurrence of specific AEs, and discontinuation due to AEs were tabulated.

Results

Demographic and baseline characteristics

A total of 14 patients (five men and nine women) aged 45 to 72 years (mean, 62.7 years) were treated. Of these, 13 (93%) were white and one (7%) was black/African-American. Subjects had a mean (range) 50 (20–100) pack-year smoking history and a mean (range) predicted post-bronchodilator FEV_1 of 58% (42%–73%). All 14 patients completed the study.

Pharmacokinetic results

Mean plasma concentration-time profiles showed prolonged absorption of MF following administration of MF by MDI (Figure 1). Median MF T_{max} values were 3.00, 2.00, and 1.00 hours for treatments A, B, and C, respectively (Table 1). Median T_{max} values of formoterol were 1.02 and 0.52 hours for Treatments A and B, respectively (Table 1).




Figure 1 Mean (SD) MF plasma concentration-time profiles at day 5 in patients with chronic obstructive pulmonary disorder.

Notes: Subjects received the following treatments by oral inhalation twice-daily for 5 consecutive days: MF 400 µg/F 10 µg via MDI without a spacer device (MF/F MDI; treatment A); MF 400 µg/F 10 µg via MDI with a spacer device (MF/F MDI with spacer; treatment B); and MF 400 µg via DPI (MF DPI; treatment C). Inset shows graph with log scale for y-axis (MF plasma concentration).

Abbreviations: DPI, dry-powder inhaler; F, formoterol fumarate; MDI, metered-dose inhaler; MF, mometasone furoate; SD, standard deviation.

For the comparison of the MF exposure following inhalation of the DPI and MDI (primary objective), the ANOVA model included all treatments. Inpatient variabilities for MF AUC_{0–12 hr} and C_{max} of 44% and 47%, respectively, were obtained from the model (Table 2). Despite this large

observed variability, MF C_{max} values were significantly different between treatments, with the mean C_{max} for the MDI being 44% lower than that for DPI (Table 2). Mean MF AUC_{0–12 hr} following inhalation by MDI alone was 23% lower than the mean value following administration of MF by DPI. The

Table 1 MF and F pharmacokinetic parameters on day 5 following twice-daily administration of MF/F MDI with or without a spacer or MF DPI in patients with chronic obstructive pulmonary disease

Parameter	n	MF/F MDI Treatment A		MF/F MDI + spacer Treatment B		MF DPI Treatment C
		MF	Formoterol	MF	Formoterol	MF
T _{max} (median), h	14	3.00	1.02	2.00	0.52	1.00
Median (range)		(0.50–8.00)	(0.18–2.02)	(0.50–8.00)	(0.18–4.02)	(0.50–12.0)
C _{max} ^a mean	14	49.2 (55)	23.78 (24)	41.1 (45)	20.77 (47)	90.2 (48)
(CV [%])		(pg/mL)	(pmol/L)	(pg/mL)	(pmol/L)	(pg/mL)
AUC _{0–12 hr} ^a mean	14	484 (56)	152.9 (26)	353 (44)	101.1 (39)	646 (47)
(CV [%])		(pg · h/mL)	(pmol · h/L)	(pg · h/mL)	(pmol · h/L)	(pg · h/mL)

Notes: MF/F MDI (treatment A) = MF 400 µg/F 10 µg inhaled orally twice a day via MDI without a spacer device; MF/F MDI + spacer (treatment B) = MF 400 µg/F 10 µg inhaled orally twice a day via MDI with a spacer device; MF DPI (treatment C) = MF 400 µg inhaled orally twice a day via DPI.

Abbreviations: AUC_{0–12 hr}^a, area under the concentration–time curve from 0 hours to 12 hours on day 5; C_{max}^a, mean maximum concentration; CV, coefficient of variation; DPI, dry-powder inhaler; F, formoterol fumarate; MDI, metered-dose inhaler; MF, mometasone furoate; T_{max}^a, time to C_{max}.

Table 2 Day 5 exposures to MF following twice-daily administration of MF/F MDI with or without a spacer or MF DPI in patients with chronic obstructive pulmonary disorder

Parameter	n	Least squares geometric mean ^a			Inpatient CV	Treatment comparison	GMR (%)	90% CI
		MF/F MDI Treatment A	MF/F MDI + spacer Treatment B	MF DPI Treatment C				
AUC _{0–12 hr} (pg·hour/mL)	14	431	322	561	44%	A versus C B versus A	77 75	58–102 56–100
C _{max} (pg/mL)	14	44	37	77	47%	A versus C B versus A	56 86	41–77 63–117
C _{trough} (pg/mL)	14	28	17	26		A versus C B versus A	106 62	80–142 51–75

Notes: ^aModel-based (least squares) mean: ANOVA model extracting the effects due to treatment, sequence, period, and patient. MF/F MDI (treatment A) = MF 400 µg/F 10 µg inhaled orally twice a day via MDI without a spacer device; MF/F MDI + spacer (treatment B) = MF 400 µg/F 10 µg inhaled orally twice a day via MDI with a spacer device; MF DPI (treatment C) = MF 400 µg inhaled orally twice a day via DPI.

Abbreviations: ANOVA, analysis of variance; AUC_{0–12 hr}, area under the concentration–time curve from 0 hours to 12 hours on day 5; CI, confidence interval; C_{max}, mean maximum concentration; C_{trough}, trough plasma concentration; CV, coefficient of variation; DPI, dry-powder inhaler; F, formoterol fumarate; GMR, geometric mean ratio; MDI, metered-dose inhaler; MF, mometasone furoate; T_{max}, time to C_{max}.

large CI reflects the small sample size and the larger than expected inpatient variability (expected variability was approximately 28%; observed was 44%).

A secondary objective of the study was to compare MF and formoterol exposure following inhalation using the MDI with and without a spacer. Following inhalation by MDI with the spacer, MF exposures based on AUC_{0–12 hr} were lower than for MDI alone (Table 2). In the initial analysis with all treatments (Table 2), the ratio estimates for MF AUC_{0–12 hr} and C_{max} included 100%. However, because the larger inpatient variability was related to the DPI treatment group, a reanalysis without treatment C showed that the AUC_{0–12 hr} and C_{max} values were 28% and 18% lower, respectively, for MDI with a spacer compared with MDI alone (Table 3). Inpatient variability for MF AUC_{0–12 hr} and C_{max} values in the original three-treatment ANOVA ranged from 44% and 47%, respectively, to 23% and 24%, when comparing only the MDI data.

Mean plots of formoterol plasma concentrations showed rapid absorption of F (Figure 2). For formoterol, AUC_{0–12 hr} and C_{max} values were 38% and 20% lower, respectively, for MDI with the spacer compared to MDI alone (Table 3).

Safety

A total of ten (71.4%) patients reported at least one AE during the study: seven (50%) during treatment A (MF 400 µg + F 10 µg via MDI), three (21.4%) during treatment B (MF 400 µg + F 10 µg via MDI with spacer), and four (28.6%) during treatment C (MF 400 µg via DPI). The most common AEs were headache and dyspepsia, occurring in eight (57.1%) and two (14.3%) patients, respectively. All reported AEs were either mild or moderate in severity. No death or serious AEs occurred during the study.

There were no clinically significant changes in blood chemistry or hematologic parameters, vital signs, or electrocardiography in any of the treatment groups.

Table 3 Day 5 exposures to MF and formoterol following twice-daily administration of MF/F MDI with or without a spacer in patients with chronic obstructive pulmonary disorder

Parameter	n	Least squares geometric mean ^{a,b}		Inpatient CV	Treatment comparison	GMR (%)	90% CI (%)
		MF/F MDI Treatment A	MF/F MDI + spacer Treatment B				
Mometasone furoate							
AUC _{0–12 hr} (pg·h/mL)	14	429	309	23%	B versus A	72	61–84
C _{max} (pg/mL)	14	44	36	24%	B versus A	82	69–97
Formoterol							
AUC _{0–12 hr} (pmol·h/L)	14	150	93	–	B versus A	62	52–74
C _{max} (pmol/L)	14	23.3	18.5	–	B versus A	80	63–101

Notes: ^aModel-based (least squares) mean: ANOVA model extracting the effects due to treatment, sequence, period, and patient; ^bbased on analyses excluding treatment C (MF 400 µg inhaled orally twice a day via dry powder inhaler). MF/F MDI (treatment A) = MF 400 µg/F 10 µg inhaled orally twice a day via metered-dose inhaler without a spacer device; MF/F MDI + spacer (treatment B) = MF 400 µg/F 10 µg inhaled orally twice a day via metered-dose inhaler with a spacer device. Because data for treatment C had higher variability than data for treatments A and B, analyses excluding treatment C were performed as sensitivity analyses.

Abbreviations: ANOVA, analysis of variance; AUC_{0–12 hr}, area under the concentration–time curve from 0 hours to 12 hours postdose; CI, confidence interval; C_{max}, maximum plasma concentration; CV, coefficient of variation; F, formoterol fumarate; GMR, geometric mean ratio; MDI, metered-dose inhaler; MF, mometasone furoate.




Figure 2 Mean (SD) formoterol plasma concentration-time profiles at day 5 after twice-daily oral inhalation of MF/F MDI (MF 400 μ g/F 10 μ g) with (treatment B) and without (treatment A) a spacer device in patients with chronic obstructive pulmonary disorder.

Note: Inset shows graph with log scale for y-axis (formoterol plasma concentration).

Abbreviations: F, formoterol fumarate; MDI, metered-dose inhaler; MF, mometasone furoate; SD, standard deviation.

Discussion

Rapid and sustained relief from bronchoconstriction and improvement in lung function are critical for the long-term management of patients with persistent symptoms and exacerbation of COPD. Coadministration of MF and F has been shown to produce additive effects in rapidly improving symptoms and lung function and reducing the frequency of exacerbation of asthma and COPD.^{9,10,17–19} The decreases in lung function seen in patients with COPD may affect systemic exposure to drugs, such as MF/F MDI, which are administered via inhalation, and thereby may alter the pharmacokinetics of MF and formoterol. This has previously been described for ICS fluticasone propionate in patients with COPD compared to healthy controls.^{12,13} Therefore, the current study was conducted in patients with moderate-to-severe COPD to assess the pharmacokinetics of MF and formoterol in the intended target population. The rationale for including the MF DPI comparison arm in this

study was to assess the relative systemic exposure of MF as administered by the MF/F MDI to the MF DPI, for which there is extensive clinical use and safety experience.¹⁶ A previous pharmacokinetic study in healthy subjects showed lower systemic exposure to MF after steady state dosing from the MDI compared to the DPI device (data on file, Merck Sharp & Dohme Corp, 2006). Similar differences in the systemic exposure of ICSs between MDI and DPI devices have been reported for other ICSs. For example, for SYMBICORT® (AstraZeneca, London, UK), an approved fixed-dose combination product containing budesonide and formoterol, the systemic exposure of budesonide was approximately 30% lower in both pediatric and adult patients with asthma after administration from an MDI device compared to the same dose delivered from a DPI device.^{14,15}

The current study demonstrated that mean systemic exposures to MF were 23% lower following administration by MF/F MDI compared to MF DPI (primary objective).

Additionally, mean systemic exposures of MF and formoterol were 28% lower and 38% lower, respectively, following administration by MF/F MDI in conjunction with a spacer (AeroChamber Plus® Valved Holding Chamber) compared to MF/F MDI without a spacer (secondary objective). The high inpatient variability in MF exposures observed in this study may in part be attributed to differences in lung function over time (eg, reduced lung inflammation over time with observed dosing) and/or day-to-day variations in patient inhalation technique. In order to control for variations in inhalation technique, patients were extensively trained on the proper use of inhalation devices at baseline and, if necessary, prior to the start of each treatment period. However, even after allowing for differences in the inhalation techniques between the MDI and DPI for the observed inpatient variability, MF and formoterol exposure following MDI treatment were still lower than those following DPI administration. These observed differences were not due to differences between the two formulations/devices in the oral deposition and subsequent gastrointestinal absorption of MF, since patients were instructed to rinse their mouths with water and spit it out after treatment administration. Furthermore, after oral administration as a solution, MF has been shown to have very low systemic bioavailability due to extensive first pass metabolism (unpublished data). The magnitude of the observed differences in systemic exposure between the MDI and DPI are probably due to formulation differences, which may result in differences in regional lung deposition and clearance from the lungs. The high inpatient variability was related to inclusion of data from MF DPI treatment group (ie, treatment C) in the ANOVA model. A reanalysis of the results excluding treatment C showed lower overall mean exposures when a spacer was used with the MF/F MDI compared to when the MF MDI was used alone.

The present results are in agreement with previous studies showing that spacer devices reduce the systemic absorption of ICS in healthy volunteers.^{20,21} In those studies, a major factor that contributed to the lower dosage delivery using a spacer was the static charge of the spacer, which attracted medication particles. The authors also reported that multiple actuations and delayed inhalation of the drug after actuation may also cause reductions in dose delivery through a spacer. Application of antistatic material or washing the spacer was useful to reduce the static effect.²¹ It should be noted, however, that the demonstrated differences in systemic exposure seen in this study in the presence/absence of a spacer device were observed in COPD patients who were trained for good, reproducible inhalation techniques with an MDI device and

therefore would be unlikely to benefit from the use of a spacer in clinical practice. Spacer devices are indicated for patients with poor coordination and poor inhalation technique in order to improve drug delivery to the lungs. In clinical practice, pharmacotherapy with an inhalation product, such as MF/F MDI, is individualized, and each patient is titrated to a desired therapeutic response. Therefore, considering that patients who require a spacer device will be dosed with, and if necessary, titrated with a spacer, the use of a spacer device is not expected to have an efficacy implication in the target population.

In this study, MF/F MDI was shown to have a similar AE profile to MF DPI. All reported AEs were either mild or moderate in intensity and no serious AE was reported in this study. These safety findings are in agreement with those of previous studies conducted in asthma patients, which also reported that treatment with MF/F was generally well tolerated.^{22,23} Nevertheless, the current short term, multiple-dose pharmacokinetic study does not address the long-term safety and tolerability profile of chronic MF/F MDI therapy in patients with COPD. Two recently published articles demonstrated treatment with MF/F MDI was well tolerated in patients with moderate-to-severe COPD over 52 weeks.^{9,10}

Considering that this study demonstrated a lower systemic exposure to the MDI product compared with the DPI in patients with COPD, there may be some concern regarding the appropriate interchangeability of the DPI monotherapy for the MDI combination therapy with regard to the comparability of the delivery of MF dose in each device. While a comparison of the systemic exposure of inhaled drugs is an acceptable way to evaluate the relative risk of ICS with regard to their systemic safety, there is considerable debate whether similar systemic exposure reflects comparable localized drug concentrations in the lung. Therefore, the clinical development program for the MF/F fixed-dose combination product has focused on demonstrating the clinical efficacy and safety of the MF/F device compared with the MF MDI device and placebo rather than the DPI reference products.^{9,10}

In this study, the observed mean difference in systemic availability between MDI and DPI formulations was 23%. However, the difference between formulations in actual lung deposition may be smaller than those noted in systemic exposure and may be due to, at least in part, differences in lung retention. This hypothesis is supported by the apparently longer MF effective half-life (25 hours) after administration from the MDI versus the DPI (effective 13 hours; unpublished data). In addition, the majority of dose response

studies conducted with inhaled corticosteroids have failed to demonstrate clinically meaningful differences even between doubling of doses,^{24,25} let alone a 25% difference. This point is illustrated by formoterol/budesonide FDC (SYMBICORT®) where the same delivered doses are approved for both the MDI (aerosol inhaler) and DPI (Turbuhaler®; AstraZeneca) formulations despite the ~30% lower systemic budesonide exposure from the MDI.^{14,15} Consistent with the relatively flat dose-response relationship of ICS, the current European Medicines Agency therapeutic equivalence guideline for orally inhaled products (CPMP/EWP/4151/00) has expanded equivalence acceptance criteria margins of 0.67 and 1.5,²⁶ which assume a mean difference between treatments of up to 1.5-fold. In view of the aforementioned considerations, the 23% lower systemic exposure to MF from the MDI relative to the DPI formulation is not considered clinically important.

Conclusion

This multiple-dose pharmacokinetic study demonstrated that systemic exposure to MF was lower following administration by MF/F MDI compared to MF DPI in patients with mild-to-moderate COPD. Additionally, systemic exposures of MF and formoterol were lower following administration by MF/F MDI with an AeroChamber Plus® Valved Holding Chamber spacer compared to MF/F MDI without this spacer. The magnitude of the differences in systemic exposure to MF seen with MF/F MDI versus MF DPI as well as MF/F MDI administered with a spacer versus MF/F MDI without a spacer were not clinically relevant. There also was no clinically relevant difference in systemic exposure to formoterol seen with MF/F MDI administered in the presence and absence of a spacer. Finally, MF/F delivered twice daily by MDI was generally well tolerated among patients with moderate-to-severe COPD in this short-term study.

Acknowledgments

The authors wish to thank Amy O Johnson-Levonas (Merck Sharp & Dohme Corp.) for writing and editing sections of the manuscript as well as Kathleen Newcomb and Jennifer Pawlowski (Merck Sharp & Dohme Corp.) for assistance with the preparation of this manuscript for submission and publication.

Disclosures

This study was financially supported by Merck Sharp & Dohme Corp, a subsidiary of Merck and Co, Inc (Whitehouse Station, NJ, USA). T Kosoglou, J Hubbell, F Xuan, AG Meehan, BS Kantesaria, and DL Cutler all declare that

they are full-time employees of Merck Sharp & Dohme Corp. BA Wittmer received research grant support from Schering-Plough Corporation (now Merck Sharp & Dohme Corp). The authors report no other conflicts of interest in this work.

References

1. American Thoracic Society/European Respiratory Society Task Force. *Standards for the Diagnosis and Management of Patients with COPD 2004*. New York, NY: American Thoracic Society/European Respiratory Society Task Force; 2005.
2. National Collaborating Centre for Chronic Conditions. Chronic obstructive pulmonary disease: national clinical guideline on management of chronic obstructive pulmonary disease in adults in primary and secondary care. *Thorax*. 2004;59 Suppl 1:1–232.
3. *Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease*. Global Initiative for Chronic Obstructive Lung Disease, Inc; 2011. Available from: http://www.goldcopd.org/uploads/users/files/GOLD_Report_2011_Feb21.pdf. Accessed August 7, 2011.
4. GINA – the Global Initiative for Asthma [webpage on the Internet]. Global strategy for asthma management and prevention. National Heart, Lung, and Blood Institute. Global Initiative for Asthma (GINA); 2010. Available from: <http://www.ginasthma.org/guideline-report-2010.html>. Accessed August 7, 2011.
5. NHLBI, Diagnosis and Management of Asthma [webpage on the Internet]. Expert Panel Report 3 (EPR3): Guidelines for the diagnosis and management of asthma. National Heart Lung and Blood Institute; updated 2012. Available from: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.htm>. Accessed August 7, 2011.
6. Bernstein DI, Berkowitz RB, Chervinsky P, et al. Dose-ranging study of a new steroid for asthma: mometasone furoate dry powder inhaler. *Respir Med*. 1999;93(9):603–612.
7. Karpel JP, Nayak A, Lumry W, et al. Inhaled mometasone furoate reduces oral prednisone usage and improves lung function in severe persistent asthma. *Respir Med*. 2007;101(3):628–637.
8. Noonan M, Karpel JP, Bensch GW, et al. Comparison of once-daily to twice-daily treatment with mometasone furoate dry powder inhaler. *Ann Allergy Asthma Immunol*. 2001;86(1):36–43.
9. Doherty DE, Tashkin DP, Kerwin E, et al. Effects of mometasone furoate/formoterol fumarate fixed-dose combination formulation on chronic obstructive pulmonary disease (COPD): results from a 52-week Phase III trial in subjects with moderate-to-very severe COPD. *Int J Chron Obstruct Pulmon Dis*. 2012;7:57–71.
10. Tashkin DP, Doherty DE, Kerwin E, et al. Efficacy and safety characteristics of mometasone furoate/formoterol fumarate fixed-dose combination in subjects with moderate to very severe COPD: findings from pooled analysis of two randomized, 52-week placebo-controlled trials. *Int J Chron Obstruct Pulmon Dis*. 2012;7:73–86.
11. LaForce C, Prenner BM, Andriano K, Lavecchia C, Yegen U. Efficacy and safety of formoterol delivered via a new multidose dry powder inhaler (Certihaler) in adolescents and adults with persistent asthma. *J Asthma*. 2005;42(2):101–106.
12. Singh SD, Whale C, Houghton N, Daley-Yates P, Kirby SM, Woodcock AA. Pharmacokinetics and systemic effects of inhaled fluticasone propionate in chronic obstructive pulmonary disease. *Br J Clin Pharmacol*. 2003;55(4):375–381.
13. Edsbäcker S, Wollmer P, Selroos O, Borgström L, Olsson B, Ingelf J. Do airway clearance mechanisms influence the local and systemic effects of inhaled corticosteroids? *Pulm Pharmacol Ther*. 2008;21(2):247–258.
14. Tronde A, Gillen M, Borgström L, Lötvall J, Ankerst J. Pharmacokinetics of budesonide and formoterol administered via 1 pressurized metered-dose inhaler in patients with asthma and COPD. *J Clin Pharmacol*. 2008;48(11):1300–1308.

15. Symbicort® (budesonide/formoterol fumarate dehydrate) [package insert]. Wilmington, DE: AstraZeneca; 2010.
16. Zeidler M, Corren J, Tashkin DP. Use of mometasone furoate administered via a dry powder inhaler in the treatment of asthma. *Curr Med Res Opin.* 2010;26(6):1295–1305.
17. Nathan RA, Nolte H, Pearlman DS; for P04334 Study Investigators. Twenty-six-week efficacy and safety study of mometasone furoate/formoterol 200/10 microg combination treatment in patients with persistent asthma previously receiving medium-dose inhaled corticosteroids. *Allergy Asthma Proc.* 2010;31(4):269–279.
18. Weinstein SF, Murphy KR, Corren J, Nolte H, White MV. Efficacy and safety of medium and high doses of mometasone furoate/formoterol (MF/F) combination treatment in subjects with severe persistent asthma [abstract]. *J Allergy Clin Immunol.* 2010;125:AB196.
19. Weinstein SF, Corren J, Murphy K, Nolte H, White M; Study Investigators of P04431. Twelve-week efficacy and safety study of mometasone furoate/formoterol 200/10 microg and 400/10 microg combination treatments in patients with persistent asthma previously receiving high-dose inhaled corticosteroids. *Allergy Asthma Proc.* 2010; 31(4):280–289.
20. Wales D, Makker H, Kane J, McDowell P, O'Driscoll BR. Systemic bioavailability and potency of high-dose inhaled corticosteroids: a comparison of four inhaler devices and three drugs in healthy adult volunteers. *Chest.* 1999;115(5):1278–1284.
21. Clark DJ, Lipworth BJ. Effect of multiple actuations, delayed inhalation and antistatic treatment on the lung bioavailability of salbutamol via a spacer device. *Thorax.* 1996;51(10):981–984.
22. Maspero J, Cherrez I, Nolte H. Long-term safety and tolerability of two doses of mometasone furoate/formoterol (MF/F) combination, administered via a metered-dose inhaler, for the treatment of moderate-to-severe persistent asthma. *J Allergy Clin Immunol.* 2009;123(2):S159.
23. Maspero JF, Nolte H, Cherrez-Ojeda I; P04139 Study Group. Long-term safety of mometasone furoate/formoterol combination for treatment of patients with persistent asthma. *J Asthma.* 2010;47(10):1106–1115.
24. Adams N, Bestall J, Jones P. Budesonide at different doses for chronic asthma. *Cochrane Database Syst Rev.* 2001;4:CD003271.
25. Adams NP, Bestall JC, Jones P, Lasserson TJ, Griffiths B, Cates CJ. Fluticasone at different doses for chronic asthma in adults and children. *Cochrane Database Syst Rev.* 2008;4:CD003534.
26. Committee for Medicinal Products for Human Use (CHMP). *Guideline on the Requirements for Clinical Documentation for Orally Inhaled Products (OIP) Including the Requirements for Demonstration of Therapeutic Equivalence between Two Inhaled Products for Use in the Treatment of Asthma and Chronic Obstructive Pulmonary Disease (COPD) in Adults and for Use in the Treatment of Asthma in Children and Adolescents.* London: European Medicines Agency; 2009. Available from: http://www.ema.europa.eu/ema/pages/includes/document/open_document.jsp?webContentId=WC500003504. Accessed August 7, 2011.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-copd-journal>

Dovepress

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.