ERRATUM

Published online: 05 Apr 2013.

Submit your article to this journal ☑

Article views: 418

View related articles ☑

ERRATUM

Correction to: Prostate cancer treatment in Europe at the end of 1990s.

Gemma G et al. Prostate cancer treatment in Europe at the end of 1990s. Acta Oncol 2009;48(6):867–73.

Following publication of this article, an error has been identified Table III. Under the column 'Only hormone %'. The numbers 3411, 4820, 4722, 7112, 5413, 4922, and 4524 have to be substituted to the following percentages 34, 48, 47, 71, 54, 49, 45. Also, under the column 'None %' blank' have to be substitute with the percentages 11, 20, 22, 12, 13, 22 and 24.

The corrected Table III is shown below.

Table 3. Prostate cancer cases according to initial therapy, age, stage and tumor grade. Analysis performed on 3407 cases (without cases with unknown treatment).

		Treatment				
	N cases (%)	RP[1] %	RRT[2] %	TURP[3] %	Only hormone %	None %
Age (years)						
< 60	264 (8)	50	14	15	17	3
60–69	1102 (32)	38	18	18	21	5
70–79	1384 (41)	15	17	24	34	11
≥ 80	657 (19)	3	3	26	48	20
Stage						
T1	552 (16)	16	15	54	8	7
T2	1035 (30)	36	20	14	21	9
T1T2[†]	287 (8)	9	13	8	47	22
T3	416 (12)	34	19	17	26	4
T4	64 (2)	30	6	28	28	8
T3T4[¥]	66 (2)	9	3	5	71	12
N+	135 (4)	51	13	9	25	2
M+	673 (20)	4	8	21	54	13
Unknown	179 (5)	8	6	16	49	22
Grade						
Well diff.	847 (25)	20	15	32	24	9
Mod. diff.	1583 (46)	30	15	18	29	8
Poorly diff.	498 (15)	17	16	25	36	6
Unknown	479 (14)	8	11	13	45	24
Overall	**3407 (100)**	**23**	**14**	**22**	**31**	**10**

[1]RP: radical prostatectomy; [2]RRT: radical radiotherapy; [3]TURP: transurethral resection.
For cancers with exact T not available: [†]localized; [¥]advanced.