



Corrigendum

To cite this article: (2011) Corrigendum, Autoimmunity, 44:8, 664-664, DOI: [10.3109/08916934.2011.592354](https://doi.org/10.3109/08916934.2011.592354)

To link to this article: <https://doi.org/10.3109/08916934.2011.592354>



Published online: 02 Aug 2011.



Submit your article to this journal [↗](#)



Article views: 250



View related articles [↗](#)

Corrigendum

The authors would like to correct errors that occurred in the publication of *Autoimmunity*.

AID dysregulation in lupus-prone MRL/Fas^{lpr/lpr} mice increases class switch DNA recombination and promotes interchromosomal c-Myc/IgH loci translocations: Modulation by HoxC4

CLAYTON A. WHITE, J. SETH HAWKINS, EGEST J. PONE, ELLIOT S. YU, AHMED AL-QAHTANI, THACH MAI, HONG ZAN, & PAOLO CASALI

18 May 2011 [Epub ahead of print] (doi:10.3109/08916934.2011.577128)

Eleven of the 14 c-Myc/IgH junction sequences (TL01, TL02, TL6, TL10, TL13, *TL10*, TL30, TL38, TL42, TL158, and TL138—the second TL10, as italicized, was a mislabel of the sequence TL19) depicted in **Figure 7** are correct. These 11 sequences were obtained by nested PCR using degenerate IgH switch region primers 5'-AGCTCATTCCAGCTCAGCTCAGCCT-3' (first round PCR) and 5'-AGCTCAGCTCAGCCTARCCC-AGCTC-3' (second round PCR), rather than the 5'-TGAGGACCAGAGAGGGATAAAAGAGAA-3' (first round PCR) and 5'-CACCTGCTATTTCCTTGTTGCTAC-3' (second round PCR), as stated in *Materials and methods*. Only degenerate primers can anneal to DNA of different S regions, thereby allowing amplification of c-Myc/IgH junctions arising from IgH breakpoints at different S regions (Potter et al., *Blood* 97: 260–269, 1997; Kovalchuk et al., *J. Exp. Med.* 204: 2989–3001, 2007). The three remaining sequences, TL-125, TL-187, and TL-162, possibly resulted from a PCR artifact involving primers 5'-TGAGGACCAGAGAGGGATAAAAGAGAA-3' (first round PCR) and 5'-CACCTGCTATTTCCTTGTTGCTAC-3' (second round PCR), and, therefore, should be excluded from data analysis.