



Antioxidant potential of lichen species and their secondary metabolites. A systematic review

Carlos Fernández-Moriano, María Pilar Gómez-Serranillos & Ana Crespo

To cite this article: Carlos Fernández-Moriano, María Pilar Gómez-Serranillos & Ana Crespo (2016) Antioxidant potential of lichen species and their secondary metabolites. A systematic review, *Pharmaceutical Biology*, 54:1, 1-17, DOI: [10.3109/13880209.2014.1003354](https://doi.org/10.3109/13880209.2014.1003354)

To link to this article: <https://doi.org/10.3109/13880209.2014.1003354>



Published online: 17 Apr 2015.



Submit your article to this journal [↗](#)



Article views: 6348



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 35 View citing articles [↗](#)

REVIEW ARTICLE

Antioxidant potential of lichen species and their secondary metabolites. A systematic review

Carlos Fernández-Moriano¹, María Pilar Gómez-Serranillos¹, and Ana Crespo²

¹Department of Pharmacology and ²Department of Plant Biology II, Faculty of Pharmacy, Universidad Complutense Madrid, Madrid, Spain

Abstract

Context: Pharmacological interest of lichens lies in their capacity to produce bioactive secondary metabolites, being most of them phenolic compounds with reactive hydroxyl groups that confer antioxidant potential through various mechanisms. Increasing incidence and impact of oxidative stress-related diseases (i.e., neurodegenerative disorders) has encouraged the search of new pharmacological strategies to face them. Lichens appear to be a promising source of phenolic compounds in the discovery of natural products exerting antioxidant activity.

Objective: The present review thoroughly discusses the available knowledge on antioxidant properties of lichens, including both *in vitro* and *in vivo* studies and the parameters assessed so far on lichen constituents.

Methods: Literature survey was performed by using as main databases PubMed, Google Scholar, Scopus, Science Direct, and Recent Literature on Lichens. We reviewed 98 highlighted research articles without date restriction.

Results: Current report collects data related to antioxidant activities of more than 75 lichen species (from 18 botanical families) and 65 isolated metabolites. Much information comes from *in vitro* investigations, such as chemical assays evaluating radical scavenging properties, lipid peroxidation inhibition, and reducing power of lichen species and compounds; similarly, research on cellular substrates and animal models generally measures antioxidant enzymes levels and other antioxidant markers, such as glutathione levels or tissue peroxidation.

Conclusion: Since consistent evidence demonstrated the contribution of oxidative stress on the development and progression of several human diseases, reviewed data suggest that some lichen compounds are worthy of further investigation and better understanding of their antioxidant and neuroprotective potentials.

Keywords

Antioxidants, neurodegenerative diseases, lichens, oxidative stress, scavenging properties

History

Received 9 July 2014

Revised 1 October 2014

Accepted 23 December 2014

Published online 17 April 2015

Introduction

A lichen is a stable, ecologically obligate, self-supporting mutualism between an exhabitant fungus (the mycobiont) and one or more extracellularly located inhabitants, which can be either unicellular or filamentous photoautotrophic partners (the photobiont: alga or cyanobacterium) (Hawksworth & Honegger, 1994). Lichens have been used with medicinal purposes since the ancient times. For instance, *Usnea barbata* (L.) Weber ex F.H. Wigg (Parmeliaceae) and other *Usnea* species were used to treat hair-related diseases, *Lobaria pulmonaria* (L.) Hoffm. (Lobariaceae) and *Parmelia sulcata* Taylor (Parmeliaceae) for pulmonary and cranial diseases, respectively, yellow-orange colored *Xanthoria parietina* (L.) Th. Fr. (Teloschistaceae) for jaundice, *Peltigera aphthosa* (L.) Willd. (Peltigeraceae) for aphta, *Parmelia saxatilis* (L.) Ach.

(Parmeliaceae) for epilepsy, etc. (Brodo et al., 2001; Malhotra et al., 2008).

Pharmaceutical importance of lichens lies in their capacity to produce a great variety of secondary metabolites, many of which only appear in these lichenised fungi. Phenolic compounds are the most relevant secondary metabolites of lichen samples, and the best studied metabolites can be principally classified as depsides, depsidones, dibenzofurans, and pulvinic acid derivatives (Huneck, 1999). Chemical structures of some representative compounds of each group are shown in Figure 1. Systematic study of pharmacological properties of lichen compounds has recently started and they have attracted much attention in recent investigations because of their antiviral, antibiotic, antitumor, allergenic, and plant growth inhibitory activities (Dias & Urban, 2009; Einarsdóttir et al., 2010; Esimone et al., 2007; Honda & Vilegas, 1999; Nishitoba et al., 1987).

In the last years, there is an increasing interest in new bioactive natural products for the prevention and treatment of various human diseases, with remarkable attention to neurodegenerative diseases and compounds exerting

Correspondence: M. Pilar Gómez-Serranillos, Department of Pharmacology, Faculty of Pharmacy, University Complutense Madrid, Plaza de Ramón y Cajal s/n 28040, Madrid, Spain. Tel: +34 913941767. Fax: +34 913941624. E-mail: pserra@ucm.es




Figure 1. Chemical structures of depside atranorin (a), dibenzofuran usnic acid (b), depsidone protocetraric acid (c), and pulvinic acid (d).

neuroprotective potential and oxidative stress reversion (Gonzalez-Burgos et al., 2013; Wang et al., 2013). This is due to the fact that compounds of natural origin normally have beneficial effects on the human organism with lower incidence of unwanted effects. Regarding this point, lichens are the subject of many research teams (Karakus et al., 2009; Kosanić et al., 2012b).

The present review aims to summarize and discuss the available information about the antioxidant properties of lichens and their isolated secondary metabolites in order to facilitate and guide future research on these natural products.

Oxidative stress

Free radicals (including reactive oxygen species, such as the hydroxyl radical, superoxide anion, hydrogen peroxide, and reactive nitrogen species, such as nitric oxide) play important roles in many chemical processes of the cells under physiological conditions, but they are also associated with pathology and cell damage. Oxidative stress is defined as an imbalance between biochemical processes leading to the production of reactive oxygen species (ROS) and those responsible for the removal of ROS, the antioxidant cascade. In that situation, free radicals will be able to attack nucleic acids and proteins, as well as unsaturated fatty acids in the cell membrane; several human chronic diseases (such as neurodegenerative diseases) are related to this problem (Molnár & Farkas, 2010; Sayre et al., 2008).

On the contrary, antioxidant agents inhibit and prevent those ROS that can cause degenerative diseases. Since many synthetic antioxidants have shown toxic and/or mutagenic effects (Grice, 1986; Wichi, 1988), the scientific attention shifted towards the discovery of naturally occurring antioxidants. With this regard, numerous plant constituents have been shown to exert antioxidant activity, being flavonoids and other phenolic compounds such as hydroxycinnamic derivatives, catechins, theaflavins, curcumins and terpenoids remarkable among them (Gonzalez-Burgos et al., 2012; Sundararajan et al., 2006). They are mostly phenolic compounds containing reactive hydroxyl groups and their antioxidant properties might be based on their ability to scavenge ROS, chelate metal ions (i.e., iron and copper),

stabilize unpaired electrons, and modulate the endogenous enzymatic and non-enzymatic antioxidant defense systems.

There is unquestionable evidence for some participation of oxidative stress in all neurodegenerative diseases (such as Alzheimer's, Parkinson's and Huntington's diseases or amyotrophic lateral sclerosis among them). It therefore suggests possibilities of intervention into etiology or disease progression through individual or combined use of antioxidants capable to enhance endogenous enzymatic or non-enzymatic defense processes (Sayre et al., 2008). Thus, since natural antioxidants are preferred over synthetic ones and there is solid basis in thinking of a potential neuroprotective activity for phenolic compounds, investigation of the antioxidant potential of lichen metabolites becomes an interesting strategy for the prevention or treatment of various oxidative stress-related diseases.

Detailed mechanisms and pathways involved in the antioxidant activity of lichens still need further investigation. But classification of these issues known so far could help to understand the pharmacology of lichens secondary metabolites, and their possibilities in the treatment of neurodegenerative disorders, thus promoting the development of neuroprotective natural products.

Literature search

Current report is intended to discuss past and current research on antioxidant properties of lichens and their secondary metabolites. With this aim, an extensive review of scientific literature was carried out by considering all highlighted research articles and other reviews on the issue, without date or language restriction. Five main databases (PubMed, Google Scholar, Scopus, Science Direct, and Recent Literature on Lichens) were used as information sources through the inclusion of the search terms "lichens", "lichen metabolites", "antioxidant activities", "oxidative stress", and their combinations. As a result, a total of 98 bibliographic references are included in the present work.

Antioxidant properties of lichens

Some lichen extracts and metabolites have already been reported for antioxidant properties due to their phenolic content; for instance, the antioxidant activities of some depsides and depsidones isolated from several lichen species have been demonstrated (de Paz et al., 2010; Hidalgo et al., 1994; Jayaprakasha & Rao, 2000), as well as the *in vitro* properties of some lichen extracts (Gülçin et al., 2002; Stojanović et al., 2010). Even so, both studies on intracellular ROS modulation by lichen metabolites/extracts and their *in vivo* effects have been recently started.

Table 1 actually includes the antioxidant activity as revealed by *in vitro* assays of more than 75 species divided into 18 botanical families, among which Parmeliaceae family is the best studied, as it is one of the most rife and widespread. Some of the reflected studies are macrostudies in which authors evaluated the same antioxidant parameter in numerous species. Similarly, in Table 2, we gather all available data related to *in vitro* antioxidant activity of isolated lichen metabolites, referring to more than 65 compounds. In general, antioxidant activity has been mainly evaluated based on some

Table 1. Antioxidant activities of lichen extracts.

Lichen species	Solvent	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO [•] -RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Phenolic/flavonoid content	Isolated compounds/ composition	Reference
<i>Arthoniaceae</i> Reichenb. ex Reichenb.								
<i>Arthothelium avastii</i> Patw. & Makhlaja	ME	2.1–68.4% 15.7 µg/ml	2.6–68.4% 13.2 µg/ml	Not studied (NS)	NS	Different content depending on lichen culture conditions	NS	Verma et al. (2008b)
<i>Catillariaceae</i> Hafellner:								
<i>Toniina candida</i> (Weber) Th.Fr.	ChE, ME, PE	ChE: 41.7 µg/ml ME: 46.5 µg/ml PE: 21.5 µg/ml	ChE: 48.9 µg/ml ME: 51.5 µg/ml PE: 50.1 µg/ml 115.8 µg/ml	NS	ChE: 56.7 µg/ml ME: 78.5 µg/ml PE: 51.5 µg/ml 0.055 (500 µg/ml)	ChE: 45.2 mg GA/g ME: 76.3 mg GA/g PE: 42.9 mg GA/g 49.8 µg PE/mg	Norstictic acid, stictic acid, usnic acid, protocetraric acid, atranorin	Manojlovic et al. (2012b)
	AcE	NS		221.5 µg/ml			Norstictic acid, stictic acid, protocetraric acid, usnic acid, atranorin	Ranković et al. (2012)
<i>Cladoniaceae</i> Zenker:								
<i>Cladonia uniaurocranea</i> (Flörke) Schaerer	AcE, DmE, EE, ME	AcE: 80.3% DmE, EE, ME: numeric value not reported (NVNR)	NVNR	NS	NS	AcE: 0.018 mg/g DmE: 0.016 mg/g EE: 0.005 mg/g ME: 0.174 mg/g 371.8 mg GA/g	Atanorin, usnic acid	Singh et al. (2011)
<i>Cladonia clathrata</i> Ahl & L. Xavier	EE 70%	NS	50.2% 69.3 µg/ml	NS	NS		NS	Silva et al. (2010)
<i>Cladonia digitata</i> (L.) Hoffm.	ME	9.8 %	NS	NS	0.085	13.7 mg GA/g	NS	Ranković et al. (2010b)
<i>Cladonia fimbriata</i> (L.) Fr.	ME	20.4%	NS	NS	0.114	26.4 mg GA/g	NS	Ranković et al. (2010b)
<i>Cladonia foliacea</i> (Huds.) Willd.	ME	NS	>1000 µg/ml	NS	NS	78.1 mg GA/g (28.2 mg Rutin/g)	NS	Mitrovic et al. (2011)
<i>Cladonia furcata</i> (Hudson) Schrad.	AcE, ME	NVNR	AcE: 471.3 µg/ml ME: >1000 µg/ml	NVNR ME > AcE	NVNR AcE > ME		NS	Luo et al. (2009)
	AcE, ME, WE	NS	NVNR ME > AcE > WE	NVNR ME > AcE > WE	NVNR ME > AcE > WE	AcE: 12.1 µg PE ME: 52.7 µg PE WE: 5.8 µg PE	NS	Kosanic et al. (2011)
	AcE	NS	44.8%	23.9%	0.051	12.9 µg PE/mg (10.6 µg RE/mg)	NS	Ranković et al. (2011)
<i>Cladonia mediterranea</i> P.A.Duvign. & Abbeyes	AcE, DmE, EE, ME	AcE: 73.7% DmE, EE, ME: NVNR	NVNR	NS	NS	AcE: 0.006 mg/g DmE: 0.001 mg/g EE: 0.008 mg/g ME: 0.009 mg/g	Usnic acid	Singh et al. (2011)
<i>Cladonia rangiformis</i> Hoffm	ChE, ME, WE	ChE: 48.3% ME: 1.6% WE: – 11.5%	NS	NS	ChE: 0.083 ME: 0.115 WE: 0.034	ChE: 0.048 µg GA/ml ME: 0.010 µg GA/ml WE: 0.004 µg GA/ml	NS	Yücel et al. (2007)
<i>Graphidaceae</i> Dumort.								
<i>Graphis gumatrana</i> Vain.	ME 10%	NS	NS	19.7–23.2 µg/ml	NS	NS	NS	Behara et al. (2006a)
<i>Graphis nakanishiana</i> Patw. & C.R. Kulk.	ME 10%	NS	NS	11.2–14.2 µg/ml	NS	NS	NS	Behara et al. (2006a)
<i>Graphis schizograpia</i> Müll. Arg.	ME 10%	NS	NS	12.3–25.6 µg/ml	NS	NS	NS	Behara et al. (2006a)
<i>Imadophilaceae</i> Triebel								
<i>Thamnolia vermicularis</i> (Sw.) Schaer.	ME	2 mg/ml: 67.0% 0.2 mg/ml: 36.0%	2 mg/ml: 72.0% 0.2 mg/ml: 36.0%	NVNR	NVNR	NS	NS	Luo et al. (2006)
<i>Lecanoraceae</i> Körb.								
<i>Lecanora atra</i> (Hudson) Ach.	AcE, ME, WE	NS	AcE: 94.7% ME: 93.3% WE: 93.2%	AcE: 84.5% ME, WE: NVNR	NVNR	AcE: 73.0 µg PE ME: 71.0 µg PE WE: 69.8 µg PE (AcE: 54.8 µg RE	NS	Kosanić and Ranković (2011)

(continued)

Table 1. Continued

Lichen species	Solvent	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO ² -RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Phenolic/flavonoid content	Isolated compounds/ composition	Reference
<i>Lecanora muralis</i> (Schreber) Rabenh.	AcE	NS	94.7%	84.5%	0.109	ME: 53.7 µg RE WE: 52.6 µg RE 73.0 µg PE/mg	NS	Ranković et al. (2011)
<i>Lobariaceae</i> Chevall. <i>Lobaria pulmonaria</i> (L.) Hoffm.	AcE	NS	52.3%	33.6%	0.061	(54.8 µg RE/mg) 43.2 µg PE/mg (34.6 µg RE/mg)	NS	Ranković et al. (2011)
<i>Nephromataceae</i> Wetm. ex J. C. David & D. Hawksw. <i>Nephroma parile</i> (Ach.) Ach.	ME, WE	ME: 87.5% WE: – 17.1%	NS	NS	ME: 0.417 WE: 0.233	ME: 87.9 mg GA/g WE: 39.2 mg GA/g	NS	Odabasoglu et al. (2004)
<i>Ochrolechiaceae</i> R. C. Harris ex Lumbsch & I. Schmitt <i>Ochrolechia parella</i> (L.) A. Massal.	ME	3.6%	NS	NS	0.098	6.2 mg GA/g	NS	Ranković et al. (2010a)
<i>Ochrolechia tartarea</i> (L.) A. Massal.	ME	26.6%	NS	NS	0.077	5.6 mg GA/g	NS	Ranković et al. (2010b)
<i>Parmeliaceae</i> Zenker <i>Bryoria fuscescens</i> (Gyeln.) Brodo & D. Hawksw	ME, WE	ME: 12.6% WE: 18.4%	NS	NS	0.202	29.4 mg GA/g	NS	Ranković et al. (2010a)
<i>Cetraria aculeata</i> (Schreber) Fr.	AcE, ME	NVNR	AcE: > 1000 µg/ml ME: > 1000 µg/ml NVNR	NVNR AcE > ME NS	ME: 0.215 WE: 0.201	ME: 48.6 mg GA/g WE: 30.3 mg GA/g	NS	Odabasoglu et al. (2005)
<i>Cetraria fastigiata</i> (Nyl.) Kärnefelt	AcE, DmE, EE, ME	AcE: 80.3% DmE, EE, ME: NVNR	NVNR	NVNR	NVNR AcE > ME NS	NS	NS	Luo et al. (2009)
<i>Cetraria islandica</i> (L.) Ach.	WE	50 µg: 96.0% 100 µg: 99.0% 200 µg: 100.0% 500 µg: 100.0%	NVNR	NVNR	NVNR	AcE: 0.019 mg/g DmE: 0.048 mg/g EE: 0.009 mg/g ME: 0.050 mg/g 0.039 µg PE/mg	Lichensterinic acid	Singh et al. (2011)
<i>Cetraria pinastri</i> (Scop.) Gray	ME	48.8%	NS	NS	0.188	AcE: 25.0 µg PE ME: 38.0 µg PE WE: 18.2 µg PE (AcE: 7.7 µg RE, ME: 25.8 µg RE, WE: 1.4 µg RE) 32.9 mg GA/g	NS	Gülçin et al. (2002)
<i>Evernia prunastri</i> (L.) Ach.	BE, DE, EE, ME	Crude wax: weak activity DE: activity EE: activity ME: activity	NS	NS	NVNR	NS	NS	Kosanić and Ranković (2011)
<i>Flavocetraria nivalis</i> (L.) Kärnefelt & A. Thell	ME	NS	727.7 µg/ml	NS	NS	AcE: 25.0 µg PE ME: 38.0 µg PE WE: 18.2 µg PE (AcE: 7.7 µg RE, ME: 25.8 µg RE, WE: 1.4 µg RE) 32.9 mg GA/g	NS	Ranković et al. (2010b)
	ME	NS	> 1000 µg/ml	NS	NS	NS	NS	Racine et al. (1980)
	AcE	NS	663.1 µg/ml	1033.6 µg/ml	500 µg/ml: 0.029	18.2 µg GA/mg 80.7 mg GA/g (27.46 mg rutin/g) 34.1 µg PE/mg	NS	Stojanovic et al. (2010)
	AcE, DmE, EE, ME	NVNR	NVNR	NS	NS	AcE: 0.013 mg/g DmE: 0.012 mg/g	NS	Mitrović et al. (2011)
						Evermic acid, physodic acid, usnic acid, atranorin, chloroatranorin	NS	Kosanić et al. (2013)
						Usnic acid	NS	Singh et al. (2011)

<i>Flavoparmelia caperata</i> (L.) Hale	ME	NS	347.2 µg/ml	NS	21.6 µg AA/g 19.3 µg T/g	EE: 0.007 mg/g ME: 0.162 mg/g 11.9 µg GA/mg	NS	Sojanović et al. (2010)
	ME	NS	549.0 µg/ml	NS	NS	90.8 mg GA/g (33.6 mg rutin/g)	NS	Mitrović et al. (2011)
<i>Hypogymnia physodes</i> (L.) Nyl.	ME	NS	79.7 µg/ml	NS	96.9 µg AA/g 25.3 µg T/g	17.5 µg GA/mg	NS	Sojanović et al. (2010)
	ME	NS	45.6 µg/ml	NS	NS	141.6 mg GA/g (20.1 mg rutin/g)	NS	Mitrović et al. (2011)
	AcE, ME, WE	NS	AcE: 60.2% ME: 73.2% WE: 30.9%	NVNR ME > AcE > WE	NVNR ME > AcE > WE	AcE: 30.1 µg PE ME: 86.8 µg PE WE: 6.3 µg PE	NS	Kosanić et al. (2011)
<i>Lethariella cashmeriana</i> Krog	Hot WE	NS	NS	NS	Activity: 35.4	NS	Chlororubrocashmerquinone	Kinoshita et al. (2010)
<i>Lethariella sernanderi</i> (Most.) Obermayer	Hot WE	NS	NS	NS	Activity: 163.9	NS	Rubrocashmerquinone	Kinoshita et al. (2010)
<i>Lethariella sinensis</i> J.C.Wei & Y.M.Jiang	Hot WE	NS	NS	NS	Activity: 23.8	NS	Rubrocashmerquinone	Kinoshita et al. (2010)
	AcE, ME, WE	NS	NVNR AcE > ME > WE	NVNR AcE > ME > WE	NVNR AcE > ME > WE	AcE: 40.4 µg PE ME: 15.3 µg PE WE: 14.9 µg PE	NS	Kosanić et al. (2011)
<i>Parmelia caperata</i> (L.) Ach.	AcE	NS	46.4%	61.5%	0.057	40.4 µg PE/mg (15.3 µg RE/mg)	NS	Kosanić et al. (2012a)
<i>Parmelia centrifuga</i> (L.) Ach.	ME	54.2%	NS	NS	0.375	49.8 mg GA/g	NS	Ranković et al. (2010a)
<i>Parmelia cinitia</i> Ach.	ME	16.0%	NS	NS	0.172	12.8 mg GA/g	NS	Ranković et al. (2010b)
<i>Parmelia pertusa</i> Schaerer	AcE, ME, WE	NS	NVNR	NVNR Weak activity	NVNR Weak activity	AcE: 18.2 µg PE ME: 34.6 µg PE WE: 12.0 µg PE (AcE: 6.6 µg RE, WE: 18.2 µg RE, WE: 5.3 µg RE)	NS	Kosanić and Ranković (2011)
<i>Parmelia saxatilis</i> (L.) Ach.	ME	25.0%	No activity	NS	NS	1.0% (w/w)	NS	Güllüce et al. (2006)
	ME, WE	Ferric thiocyanate method WE: 86.1% ME: 94.8% TBA test WE > ME	WE: 19.8% ME: 63.6%	WE: 83.0% ME: 80.0%	ME > WE	WE: 24.3 mg PE/mg ME: 55.1 mg PE/mg	Atanorin, chloratanorin, salazinic acid	Özen and Kinaloglu (2008)
<i>Parmelia sulcata</i> Taylor	AcE	NS	55.3%	35.3%	0.066	40.6 µg PE/mg (20.6 µg RE/mg)	NS	Kosanić et al. (2012a)
	ME	NS	493.6 µg/ml	NS	25.3 µg AA/g 41.9 µg T/g	7.9 µg GA/g	NS	Sojanović et al. (2010)
	ME	NS	584.2 µg/ml	NS	NS	88.3 mg GA/g (44.9 mg rutin/g)	NS	Mitrović et al. (2011)
	AcE, ME, WE	NS	NVNR AcE > ME > WE	NVNR AcE > ME > WE	NVNR AcE > ME > WE	AcE: 38.2 µg PE ME: 25.1 µg PE WE: 9.6 µg PE 18.2 µg PE/mg (6.6 µg RE/mg)	NS	Kosanić et al. (2011)
	AcE	NS	38.8%	33.8%	0.034	NS	NS	Kosanić et al. (2012a)
<i>Parmotrema pseudotinctorum</i> (Abbayes) Hale	ME	NS	500 µg/ml: 90.7% + honey: 81.5%	NS	NS	NS	Atanorin, lecanoric acid	Prashith Kekuda et al. (2009)
	ME	NS	62.1–89.4%	NS	0.265–0.776	NS	Atanorin, lecanoric acid	Praveen Kumar et al. (2010)
<i>Parmotrema stuppeum</i> (Taylor) Hale	BE, AcE	BE: 30.0–65.0% AcE: 35.0–68.0%	NS	NS	NS	NS	Methyl orsellinate, orsellinic acid, atanorin, lecanoric acid	Jayaprakasha and Rao (2000)
	ME	NS	NS	NS	NS	NS	NDR	Verma et al. (2008b)

(continued)

Table 1. Continued

Lichen species	Solvent	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO ²⁻ RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Phenolic/flavonoid content	Isolated compounds/ composition	Reference
<i>Parmotrema tinctorum</i> (Delise ex Nyl.) Hale	ME	3.6–71.3% 11.5 µg/ml 27.0%	1.1–57.2% 13.6 µg/ml No activity	NS	NS	Depends on lichen culture conditions 1.1 % (w/w)	NS	Güllüce et al. (2006)
<i>Platismatia glauca</i> (L.) Culb. & C. Culb.	ME, WE	ME: 26.4% WE: 48.2%	NS	NS	ME: 0.229 WE: 0.229 NVNR	ME: 75.1 mg GA/g WE: 61.7 mg GA/g AcE: 76.4 µg PE ME: 37.0 µg PE WE: 18.7 µg PE (AcE: 37.6 µg RE, ME: 21.0 µg RE, WE: 1.8 µg RE)	NS	Odabasoglu et al. (2005)
<i>Pseudoevernia</i> <i>furfuracea</i> (L.) Zopf	AcE, ME, WE	NS	AcE: 87.3% ME: 57.9% WE: 33.9%	NVNR	NS	NS	NS	Kosanić et al. (2011)
<i>Pseudoevernia furfuracea</i> (L.) Zopf	BuE, DcmE, EaE, ME	BuE: No activity DcmE: 17.8% EaE: 23.8% ME: 17.2%	TLC spray	NS	NS	NS	Ataric acid, metil hematommate, methyl chlorohematommate	Glivenç et al. (2012)
<i>Pseudophebe pubescens</i> (L.) M. Choisy	AcE	NS	401.7 µg/ml	632.0 µg/ml	500 µg/ml: 0.058	76.4 µg PE/mg	Physodic acid, physodalic acid, atranorin, chloratranorin, 3- hydroxyphysodic Atranorin	Kosanić et al. (2013)
<i>Usnea antarctica</i> Du Rietz	AcE, ME	AcE: 82.4% DmE, EE, ME: NVNR	AcE: 51.8% DmE, EE, ME: NVNR	NS	NS	AcE: 0.017 mg/g DmE: 0.012 mg/g EE: 0.001 mg/g ME: 0.027 mg/g	NS	Singh et al. (2011)
<i>Usnea aurantiacoatra</i> (Jacq.) Bory	AcE, ME	NVNR	AcE: >1000 µg/ml ME: >1000 µg/ml	NVNR	NVNR	NS	NS	Luo et al. (2009)
<i>Usnea barbata</i> Motyka	AcE	NVNR	AcE: 445.7 µg/ml ME: >1000 µg/ml	NVNR	NVNR	NS	NS	Luo et al. (2009)
<i>Usnea complanata</i> (Müller Arg.) Motyka	WE, EE, EaE, ME	WE: 132.4 µg/ml EE: 125.0 µg/ml EaE: 157.9 µg/ml ME: 74.6 µg/ml	WE: 25.0 µg/ml EaE: 25.0 µg/ml EE: 22.9 µg/ml ME: 80.0 µg/ml	NS	500 µg/ml: 0.019	31.3 µg PE/mg	Norstictic acid, usnic acid, atranorin, chloratranorin	Ranković et al. (2012)
<i>Usnea florida</i> (L.) Weber ex F.H.Wigg.	AcE, ME	WE: 15.0% ME: 13.4% AcE: 55.0% DmE: 17.0% ME: 87.0% PE: 38.0%	AcE: 31.0% DmE: 11.0% ME: 73.0% PE: 43.0%	NS	NS	TLC technique	Usnic acid, psoromic acid	Behera et al. (2012)
<i>Usnea ghattensis</i> G. Awasthi	AcE, DmE, ME, PE	AcE: 55.0% DmE: 17.0% ME: 87.0% PE: 38.0%	AcE: 31.0% DmE: 11.0% ME: 73.0% PE: 43.0%	NS	ME: 0.069 WE: 0.083	ME: 10.5 mg GA/g WE: 10.4 mg GA/g AcE: 14.0 mg/g DmE: 12.0 mg/g ME: 35.0 mg/g PE: 9.0 mg/g 13.0 µg PE/mg	NS	Odabasoglu et al. (2004)
<i>Usnea longissima</i> Ach.	ME	2–20 mg/ml: 3.8–73.3% 67.0% ME: 82.4% WE: 25.6% 6 mg/ml: 97.3%	NS	NS	NS	NS	NS	Behera et al. (2005a)
<i>Usnea longissima</i> Ach.	ME, WE	ME: 82.4% WE: 25.6% 6 mg/ml: 97.3%	NS	NS	NS	NS	NS	Behera et al. (2006c)
<i>Usnea longissima</i> Ach.	ME	NS	NS	NS	NS	NS	NS	Verma et al. (2008a)
<i>Usnea longissima</i> Ach.	ME, WE	ME: 74.0% WE: 26.4%	NS	NS	ME: 0.178 WE: 0.100	ME: 38.6 mg GA/g WE: 18.3 mg GA/g 2.62%	NS	Odabasoglu et al. (2004)
<i>Usnea longissima</i> Ach.	ME	NS	NS	NS	NS	NS	NS	Kim and Cho (2007)
<i>Usnea longissima</i> Ach.	ME, WE	ME: 74.0% WE: 26.4%	NS	NS	ME: 0.542 WE: 0.218	ME: 66.4 mg GA/g WE: 38.1 mg GA/g	NS	Odabasoglu et al. (2005)
<i>Usnea longissima</i> Ach.	ME	22.6%	NS	NS	0.195	27.6 mg GA/g	NS	Ranković et al. (2010a)
<i>Usnea longissima</i> Ach.	ME	3.4–52.7% 12.7 µg/ml	1.6–53.7% 18.4 µg/ml	NS	NS	Different contents depending on lichen culture conditions	NDR	Verma et al. (2008b)

<i>Physcia caesia</i> (Hoffm.) Fürr.	AcE, DmE, EE, ME	NVNR	NS	NS	AcE: 0.065 mg/g EE: 0.015 mg/g ME: 0.128 mg/g DmE: 0.016 mg/g	Atanorin, zeorin	Singh et al. (2011)
<i>Ramalinaceae</i> C. Agardh							
<i>Ramalina conduplicans</i> Vain.	ME	NS	NS	NS	NS	Usnic acid, salazinic acid, sekkikaic acid	Vinayaka et al. (2009)
<i>Ramalina hossei</i> Vain.	ME	NS	NS	NS	NS	Usnic acid, sekikaic acid	Prashith Kekuda et al. (2009)
<i>Ramalina pollinaria</i> (Westr.) Ach.	ME	26.0%	NS	NS	1.0 % (w/w)	NS	Güllüce et al. (2006)
<i>Ramalina polymorpha</i> (Lilj.) Ach.	ME	19.0%	NS	NS	0.8 % (w/w)	NS	Güllüce et al. (2006)
<i>Ramalina terebrata</i> Hook. F. & Taylor	ME-WE (90/10)	NS	NS	NS	NVNR	NS	Bhattarai et al. (2008)
<i>Sphaerophoraceae</i> Fr.							
<i>Sphaerophorus globosus</i> (Huds.) Vain.	AcE, ME	NVNR	NVNR AcE > ME	NS	NS	NS	Paudel et al. (2008)
<i>Stereocaulaceae</i> Chevall.							
<i>Stereocaulon alpinum</i> Laurer ex Funck	ME-WE (90/10)	NS	NS	NS	NVNR	NS	Bhattarai et al. (2008)
<i>Teloschistaceae</i> Zahlbr.							
<i>Calopaca regalis</i> (Vain.) Zahlbr.	ME-WE (90/10)	NS	NS	NS	NVNR	NS	Bhattarai et al. (2008)
<i>Fulgensia fulgens</i> (Sw.) Elenkin	ME-WE (90/10) ME	NS 21.6%	NS NS	NS 0.136	NS 12.4 mg G/A/g	NS NS	Paudel et al. (2008) Ranković et al. (2010b)
<i>Xanthoria elegans</i> (Link) Th. Fr.	AcE, DmE, EE, ME	NVNR	NS	NS	AcE: 0.008 mg/g EE: 0.003 mg/g ME: 0.151 mg/g DmE: 0.036 mg/g	Xanthorina	Singh et al. (2011)
<i>Trypetheliaceae</i> Zenker							
<i>Laurera benguelensis</i> Zahlbr.	ChE and benzene (BF) and methanol fractions (MF)	NS	NS	NS	NS	Lichexanthone, secalonic acid D, norlichexanthone, parietin, enodin, teloschistin and citreorosein	Manojlovic et al. (2010)
<i>Umbilicariaceae</i> Chevall.							
<i>Lasallia pustulata</i> (L.) Méret	AcE, ME, WE	NS	AcE: 67.4% ME, WE: NVNR	NVNR AcE > ME > WE	AcE: 84.3 µg PE ME: 49.6 µg PE WE: 23.9 µg PE	NS	Kosanić et al. (2011)
<i>Umbilicaria Antarctica</i> Frey & Lamb	AcE, ME	NVNR	AcE: 91.1% ME: NVNR	NVNR AcE > ME	NS	Lecanoric acid	Luo et al. (2009)
<i>Umbilicaria aprina</i> var. <i>hutei</i> Nyl.	ME	NS	NS	NS	40.6 mg G/A/g	Lecanoric acid	Bucukoglu et al. (2012)
<i>Umbilicaria crustulosa</i> (Ach.) Lamy	AcE	NS	Moderate 42.0%	Low 0.050	39.6 µg PE/mg (28.1 µg RE/mg)	NS	Kosanić et al. (2012b)
<i>Umbilicaria cylindrica</i> (L.) Delise ex Duby	AcE, ME, WE	NS	NVNR	NVNR	AcE: 19.1 µg PE ME: 42.5 µg PE WE: 19.1 µg PE	NS	Kosanić and Rankovic (2011)

(continued)

Table 1. Continued

Lichen species	Solvent	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO ⁻ RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Phenolic/flavonoid content	Isolated compounds/ composition	Reference
	ME	NS	21.1% (5 mg/ml)	NS	NS	(AcE: 12.9 µg RE, ME: 19.1 µg RE, WE: 11.1 µg RE) 0.9 mg GA/g	NS	Buçukoglu et al. (2012)
	AcE	NS	39.4%	Moderate 33.9%	Low 0.035	19.1 µg PE/mg (12.9 µg RE/mg)	NS	Kosami et al. (2012b)
	ChE, ME	ChE: 29.3 µg/ml ME: 35.4 µg/ml	ChE: 31.3 µg/ml ME: 34.4 µg/ml	NS	NS	ME: 79.2 mg GA/g ChE: 71.3 mg GA/g	Salazinic acid, norstictic acid, methyl-β-orcinol carboxylate, ethyl haematommate, usnic acid, atranorin	Manojlović et al. (2012b)
<i>Umbilicaria esculenta</i> (Miyoshi) Minks	ME	6 mg/ml: 92.1%	6 mg/ml: 1.3 mg/ml	NS	NS	1.5%	NS	Kim and Cho (2007)
<i>Umbilicaria decussata</i> (Vill.) Zahlbr.	ME	NS	5 mg/ml: 23.4%	NS	NS	3.3 mg GA/g	NS	Buçukoglu et al. (2012)
<i>Umbilicaria hyperborea</i> (Ach.) Hoffm.	AcE, DE, EE, ME	NVNR	NVNR	NS	NS	AcE: 0.001 mg/g DmE: 0.029 mg/g EE: 0.001 mg/g ME: 0.046 mg/g 35.9 mg GA/g	Gyrophoric acid	Singh et al. (2011)
<i>Umbilicaria leiocarpa</i> DC.	ME	NS	5 mg/ml: 32.8%	NS	NS	3.0 % (w/w)	NS	Buçukoglu et al. (2012)
<i>Umbilicaria nylanderiana</i> (Zahlbr.) H. Magn.	ME	53.0%	400.2 µg/ml	NS	NS		NS	Güllüce et al. (2006)
<i>Umbilicaria polyphylla</i> (L.) Baumg.	AcE	NS	5 mg/ml: 32.9%	NS	NS	48.3 mg GA/g	Umbilicic acid	Buçukoglu et al. (2012)
<i>Umbilicaria virginis</i> Schaerer	ME	NS	72.8%	Moderate 55.4%	Low 0.065	47.0 µg PE/mg (30.3 µg RE/mg)	NS	Kosami et al. (2012b)
			5 mg/ml: 20.3%	NS	NS	15.3 mg GA/g	Gyrophoric acid	Buçukoglu et al. (2012)
<i>Verrucariaceae</i> Zenker	ME, WE	ME: 5.3% WE: 9.5%	NS	NS	ME: 0.119 WE: 0.131	ME: 18.6 mg GA/g WE: 9.2 mg GA/g	NS	Odabasoglu et al. (2005)
<i>Dermacarpion inextinctum</i> (Körber) Hasse								
Some macrostudies:								
Macrostudy of 46 lichen species	ME	S.n.: 85.5% P.p.: 85.4%	S.n.: 90.4% P.p.: 87.8%	NS	S.n.: 1.5 P.p.: 1.0	S.n.: 156.1 µg CE/mg P.p.: 109.3 µg CE/mg	<i>Sticta nylanderiana</i> : lecanoric acid	Luo et al. (2010)
Highest activities:								
- <i>Sticta nylanderiana</i> Zahlbr. (S.n.)								
- <i>Peltigera praetextata</i> (Flörke ex Sommerf.) Zopf (P.p.)								
Macrostudy of 77 lichen species (<i>Graphidaceae</i> sp.)	DmE, ME	NS	NS	Screening of 77 extracts by two methods. Highest SO ⁻ RSA (>50%): <i>Graphina multistriata</i> <i>Graphina salaciniabata</i> <i>Graphis assamensis</i> <i>Graphis guianana</i> <i>Graphis sikkimensis</i>	NS	NS	NS	Behera et al. (2003)

Macrostudy of 85 lichen species	EE	NS	Screening of 99 extracts. Highest DPPH scavenging activities: <i>Hypogymnia vittata</i> <i>Peltigera aphthosa</i> <i>Nephromopsis ornata</i> <i>Pseudoevernina furfuracea</i> <i>Cladonia vulcani</i> <i>Peltigera elizabethae</i>	NS	NS	NS	<i>Peltigera aphthosa</i> : solorinine	Hara et al. (2011)
---------------------------------	----	----	--	----	----	----	--	--------------------

NS, not studied; NVNR, numeric value not reported; LP, lipoperoxidation; %INH, percentage of inhibition; SO⁻ RSA, superoxide radical scavenging activity; GA, gallic acid; PE, pyrocatechol equivalent; RE, rutin equivalent; AA, ascorbic acid; T, tocopherol; TBA, thiobarbituric acid; CE, catechol equivalent; MDA, malondialdehyde; ChE, chloroform extract; ME, methanol extract; PE, petrol ether extract; AcE, acetone extract; DmE, dimethyl sulfoxide extract; EE, ethanol extract; WE, water extract; DE, diethyl-ether extract; BE, benzene extract; BuE, butanol extract; DemE, dichlorometane extract; EaE, ethyl acetate extract.

chemical assays, such as DPPH free radical scavenging activity, superoxide anion radical scavenging activity, reducing power, and lipid peroxidation inhibition. Methanol arises as one of the most used solvents for an efficient extraction of lichens bioactive compounds with antioxidant activities and, therefore, many antioxidant activity assays have been performed on methanol extracts (Stojanović et al., 2010; Kosanić & Ranković, 2011; Zambare & Christopher, 2012).

In the last part of our review, we focus on antioxidant responses of these natural products to oxidative stress occurring at intracellular level and *in vivo* trials. Therefore, we collect the more recent investigations on antioxidant activity of lichens species (both purified metabolites and extracts) on *in vitro* cellular substrates (Table 3) and *in vivo* animal models (Table 4).

Apart from all species and assays shown in the tables, some authors have measured similar parameters by other methods (e.g., other radical scavenging properties) and also different parameters related to the antioxidant potential of the aforementioned lichens and compounds. Regarding other radical scavenging properties, Papadopoulou et al. (2007) evaluated the hydroxyl radical scavenging activity of β-orcinol metabolites of the lichen *Hypotrachyna revoluta* (Flörke) Hale (Parmeliaceae); this OH⁻radical scavenging activity has also been measured in *Toninia candida* (Weber) Th.Fr. (Ramalinaceae) (Manojlovic et al., 2012b), *Usnea ghattensis* G. Awasthi (Parmeliaceae) (Verma et al., 2008a), and *Umbilicaria cylindrica* (L.) Delise ex Duby (Umbilicariaceae) (Manojlovic et al., 2012c). Nitric oxide radical scavenging activity was assayed on *Usnea complanata* (Müll. Arg.), Motyka (Parmeliaceae) (Behera et al., 2012), *Usnea ghattensis* (Verma et al., 2008b), psoromic and usnic acids (Behera et al., 2012), and on other 14 lichen-purified metabolites (Thadhani et al., 2011). Moreover, hydrogen peroxide scavenging activity was investigated on *Parmelia saxatilis* (Özen & Kinalioglu, 2008) and the TEAC value (Trolox equivalent antioxidant capacity) has been determined using the ABTS radical assay in several polar lichen species (Paudel et al., 2008; Singh et al., 2011), as well as in *Usnea ghattensis* (Behera et al., 2005a; Verma et al., 2008a), *Laurera benguelensis* Zahlbr. (Zahlbr.) (Manojlovic et al., 2010), and ramalin compound (Paudel et al., 2011).

Behera et al. (2003, 2006a) assessed the xanthine oxidase inhibitory capacity for many species of the family Graphidaceae; the ferrous ion-chelating activity was investigated in *Umbilicaria cylindrica* (Manojlovic et al., 2012c) and *Toninia candida* (Manojlovic et al., 2012b); and the tyrosinase inhibitory activity has been studied in some lichens and isolated compounds (Behera et al., 2006b; Kim & Cho, 2007; Paudel et al., 2011).

Finally, it is remarkable that the work conducted by Lopes et al. (2008) assayed the radical scavenging properties for many semisynthetic derivatives from the lecanoric acid obtained from a *Parmotrema tinctorum* (Delise ex Nyl.) Hale (Parmeliaceae) specimen and treated with alcohols. They obtained the three most active compounds at scavenging DPPH radical that were orsellinic acid, orcinol and resorcinol, and the lowest activity was displayed by methyl orsellinate.

Table 2. Antioxidant activities of isolated compounds from lichens.

Compound	Origin	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Reference
1,8-Dihydroxy-5-methoxy-3-methylxanthone	<i>Pyrenula japonica</i> Kurok [DE]	Not studied (NS)	Numeric Value Not Reported (NVNR)	NS	NS	Takenaka et al. (2000)
1,5,8-Trihydroxy-3-methylxanthone	<i>Pyrenula japonica</i> Kurok [DE]	NS	NVNR	NS	NS	Takenaka et al. (2000)
1,7-Dihydroxy-3-methylxanthone	<i>Pyrenula japonica</i> Kurok [DE]	NS	NVNR	NS	NS	Takenaka et al. (2000)
1,2,8-Trihydroxy-5-methoxy-3-methylxanthone	<i>Pyrenula japonica</i> Kurok [DE]	NS	NVNR	NS	NS	Takenaka et al. (2000)
1-Chloropannarin	<i>Erioderma chilense</i> Mont.	30% In rat brain homogenate: 27–66%	NS	NS	NS	Hidalgo et al. (1994)
7-Chlorocanarione	<i>Lehariella cashmeriana</i> Krog [ME]	NS	NS	Activity: 142.7	NS	Kinoshita et al. (2010)
2-O-Methylseckiaic acid	<i>Ramalina aschinae</i> W.L. Culb & C.F. Culb. [ME, AcE]	NS	1/IC ₅₀ = 0.052	NS	NS	Valencia-Islas et al. (2007)
Atranol	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	NVNR	NS	NS	NS	Toledo-Marante et al. (2003)
Atranorin	<i>Placopsis</i> sp.	6.5% In rat brain homogenate: 7.3–9.5%	NS	NS	NS	Hidalgo et al. (1994)
	<i>Parmotrema stippeum</i> (Taylor) Hale [BE, AcE]	14.0%	NS	NS	NS	Jayaprakash et al. (2000)
	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	Unable to prevent lipid peroxidation	NS	NS	NS	Toledo-Marante et al. (2003)
	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
Barbatic acid	<i>Parmotrema stippeum</i> (Taylor) Hale [ME, AcE]	NS	1/IC ₅₀ = 0.576	NS	NS	Valencia-Islas et al. (2007)
	<i>Parmotrema grayana</i> Hue. [DcmE], <i>Heterodermia obscurata</i> (Nyl.) Trevisan [DcmE], and <i>Cladonia</i> sp. [DcmE]	NS	No activity	1.6%	NS	Thadhani et al. (2011)
	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
Boninic acid	<i>Ramalina aschinae</i> W.L. Culb & C.F. Culb. [ME, AcE]	NS	1/IC ₅₀ = 0.014	NS	NS	Valencia-Islas et al. (2007)
Canarione	<i>Lehariella seranderi</i> (Motyka) Obermayer [AcE]	NS	NS	Activity: 137.2	NS	Kinoshita et al. (2010)
Chloratranol	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	50.0–60.0%	NS	NS	NS	Toledo-Marante et al. (2003)
Chloratranorin	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	50.0–57.0%	NS	NS	NS	Toledo-Marante et al. (2003)
	<i>Parmotrema stippeum</i> (Taylor) Hale [ME, AcE]	NS	1/IC ₅₀ = 0.589	NS	NS	Valencia-Islas et al. (2007)
Chlorobenatommic acid	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	NVNR	NS	NS	NS	Toledo-Marante et al. (2003)
Chlororubrocashmerquinone	<i>Lehariella seranderi</i> (Motyka) Obermayer [AcE], <i>L. sinensis</i> J.C. Wei & Y.M. Jiang [ME], and <i>L. cashmeriana</i> Krog [ME]	NS	NS	Activity: 277.4	NS	Kinoshita et al. (2010)
Cryptostictinolide	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	18.0%	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
Diffractaic acid	<i>Usnea longissima</i> Ach. [EE]	NVNR – no antioxidant activity	No scavenging activity	NS	NS	Atalay et al. (2011)
Divaricatic acid	<i>Protousnea magellanica</i> (Mont.) Krog <i>Protousnea malacea</i> (Sirt.) Krog	NS 8.6% In rat brain homogenate: 1.7–8.4%	No scavenging activity NS	NS NS	NS NS	Brisdelli et al. (2013) Hidalgo et al. (1994)
Ergosterol peroxide	<i>Parmotrema grayana</i> Hue [DcmE] <i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS NS	27.0% NVNR	92.5% NDR	NS NS	Thadhani et al. (2011) Lohézic-Le Dévéhat et al. (2007)
Erythrin	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	LNVNR	NVNR	NS	NS	Atalay et al. (2011)
Ethyl chlorohematommate	<i>Rocella montagnei</i> Bel. [AcE]	NS	No scavenging activity	84.1%	NS	Thadhani et al. (2011)
Ethyl hematommate	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	50.0–60.0%	NS	NS	NS	Toledo-Marante et al. (2003)
Evemic acid	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	NVNR	NS	NS	NS	Toledo-Marante et al. (2003)
Fumarprotocetraric acid	<i>Evernia prunastri</i> Ach. [AcE] <i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS NS	322.4 µg/ml 20.0%	561.8 µg/ml 566.0 µM	500 µg/ml NS	Kosani et al. (2013) Lohézic-Le Dévéhat et al. (2007)

Gyrophoric acid	<i>Umbilicaria virginis</i> Schaerer	NS	50.9%	NS	NS	Bucukoglu et al. (2012)
Hematommic acid	<i>Lehmanniella canariensis</i> (Ach.) Krog [ME]	50.0–60.0%	NS	NS	NS	Toledo-Maranete et al. (2003)
Isidiphoric acid	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NVNR – high scavenging activity	NS	NS	Atalay et al. (2011)
Lecanoric acid	<i>Parmotrema stipuleum</i> (Taylor) Hale [BE, AcE]	12.0–36.0%	NS	NS	NS	Jayaprakash et al. (2000)
Lecanoric acid	<i>Parmotrema tinctorum</i> (Nyl.) Hale	NS	42.87 mM	NS	NS	Lopes et al. (2008)
Lecanoric acid	<i>Sticta nyländeriana</i> Zahlbr. [ME]	51.5%	35.4%	NS	NS	Luo et al. (2010)
Lecanoric acid	<i>Parmotrema grayana</i> Hue. [ME]	NS	34.0%	98.4%	NS	Thadhani et al. (2011)
Loboric acid	<i>Umbilicaria aprina</i> var. <i>halei</i> Nyl. [ME]	NS	32.5%	NS	NS	Bucukoglu et al. (2012)
Loboric acid	<i>Cladonia</i> sp. [ME]	NS	No scavenging activity	96.8%	NS	Thadhani et al. (2011)
Loboric acid	<i>Stereocaulon alpinum</i> Laur.	NS	No scavenging activity	NS	NS	Brisdelli et al. (2013)
Methyl chlorohematommate	<i>Lehmanniella canariensis</i> (Ach.) Krog [ME]	66.0–70.0%	NS	NS	NS	Toledo-Maranete et al. (2003)
Methyl hematommate	<i>Lehmanniella canariensis</i> (Ach.) Krog [ME]	NVNR	NS	NS	NS	Toledo-Maranete et al. (2003)
Methyl hematommate	<i>Parmotrema grayana</i> Hue. [DcmE], and <i>Heterodermia obscurata</i> (Nyl.) Trevisan [DcmE]	NS	No scavenging activity	20.4%	NS	Thadhani et al. (2011)
Methyl orcinol carboxylate	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
Methyl orsellinate	<i>Heterodermia obscurata</i> (Nyl.) Trevisan [DcmE], and <i>Cladonia</i> sp. [DcmE]	NS	No scavenging activity	1.5%	NS	Thadhani et al. (2011)
Methyl orsellinate	<i>Parmotrema stipuleum</i> (Taylor) Hale [BE, AcE]	18.0–40.0%	NS	NS	NS	Jayaprakash et al. (2000)
Methyl orsellinate	<i>Lehmanniella canariensis</i> (Ach.) Krog [ME]	NVNR	NS	NS	NS	Toledo-Maranete et al. (2003)
Methyl orsellinate	<i>Parmotrema grayana</i> Hue. [ME], <i>Heterodermia obscurata</i> (Nyl.) Trevisan [DcmE], and <i>Cladonia</i> sp. [ME]	NS	2.4%	21.2%	NS	Thadhani et al. (2011)
Montagnitol	<i>Rocella montagnei</i> Bél. [AcE]	NS	No scavenging activity	9.9%	NS	Thadhani et al. (2011)
Norstictic acid	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	580.0 µM	NS	Lohézic-Le Dévéhat et al. (2007)
Orcinol	<i>Toninia candida</i> (Weber) Th. Fr. [AcE]	NS	102.7 µg/ml	133.5 µg/ml	500 µg/ml: 0.547	Rankovic et al. (2012)
Orcellinic acid	<i>Parmotrema grayana</i> Hue. [ME]	NS	49.7%	38.2%	NS	Thadhani et al. (2011)
Orcellinic acid	<i>Parmotrema stipuleum</i> (Taylor) Hale. [BE, AcE]	26–50%	NS	NS	NS	Jayaprakash et al. (2000)
Pannarin	<i>Parmotrema grayana</i> Hue. [ME]	NS	39.7%	7.9%	NS	Thadhani et al. (2011)
Pannarin	<i>Erioderma chilense</i> Mont.	23.0%	NS	NS	NS	Hidalgo et al. (1994)
Physodic acid	<i>Pseudovernia fufriraceae</i> (L.) Zopf. [AcE]	NS	69.1 µg/ml	118.2 µg/ml	500 µg/ml: 0.664	Kosanić et al. (2013)
Protolichsterinic acid	<i>Comicularia aculeata</i> (Schreb.) Ach.	NS	No scavenging activity	NS	NS	Brisdelli et al. (2013)
Protocetraric acid	<i>Parmelia caperata</i> (L.) Ach. [AcE]	NS	138.2 µg/ml	177.6 µg/ml	500 µg/ml: 0.090	Manojlovic et al. (2012a)
Psoromic acid	<i>Usnea complanata</i> (Müll. Arg.) Motyka [total extract]	0.2 mg/ml	0.271 mg/ml	NS	NS	Behra et al. (2012)
Pulmonarianin	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NVNR – low scavenging activity	NS	NS	Atalay et al. (2011)
Ramalin	<i>Ramalina terebrata</i> Hook. f. & Taylor. [ME]	NS	0.990 µg/ml	10.2 µg/ml	0.4 µg of BHT equivalent	Paudel et al. (2011)
Rhizaldehyde	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NVNR	NS	NS	Atalay et al. (2011)
Rhizonil alcohol	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NVNR	NS	NS	Atalay et al. (2011)
Rubrocassimerquinone	<i>L. seranderi</i> (Motyka) Obermayer [AcE], and <i>L. sinensis</i> Wei & Jiang. [ME]	NS	NS	Activity: 87.2	NS	Kinoshita et al. (2010)
Salazinic acid	<i>Parmelia</i> sp. (<i>P. saxatilis</i> (L.) Ach. and <i>P. sulcata</i> Taylor.) [AcE]	NS	91.6 µg/ml	119.1 µg/ml	500 µg/ml: 0.184	Manojlovic et al. (2012a)
Salazinic acid	<i>Parmotrema stipuleum</i> (Taylor) Hale. [ME, AcE]	NS	1/IC ₅₀ = 0.014	NS	NS	Valencia-Islas et al. (2007)
Sekikic acid	<i>Heterodermia obscurata</i> (Nyl.) Trevis. [ME]	NS	32.6%	98.2%	NS	Thadhani et al. (2011)
Solorinic acid	<i>Peltigera aphthosa</i> (L.) Willd. [EE]	NS	120.0 µmol/ml	NS	NS	Hara et al. (2011)
Stictic acid	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
Umbilicic acid	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NVNR	NS	NS	Atalay et al. (2011)
Umbilicic acid	<i>Umbilicaria nyländeriana</i> (Zahlbr.) H. Magn [ME]	NS	68.1%	NS	NS	Bucukoglu et al. (2012)

(continued)

Table 2. Continued

Compound	Origin	LP inhibition (IC ₅₀ or %INH)	DPPH (IC ₅₀ or %INH)	SO RSA (IC ₅₀ or %INH)	Reducing power (A ₇₀₀ or IC ₅₀)	Reference
Usnic acid	<i>Lehariella canariensis</i> (Ach.) Krog [ME]	Unable to prevent lipid peroxidation	NS	NS	NS	Toledo-Marante et al. (2003)
	<i>Usnea articulata</i> (L.) Hoffm. [various extracts]	NS	NVNR	NVNR	NS	Lohézic-Le Dévéhat et al. (2007)
	<i>Rumalina aschmiae</i> W. L. Culb. & C. F. Culb. [ME, AcE]	NS	1/IC ₅₀ = 0.112	NS	NS	Valencia-Islas et al. (2007)
	<i>Usnea longissima</i> Ach. [EE]	NVNR – no antioxidant activity	No scavenging activity	NS	NS	Atalay et al. (2011)
	<i>Parmotrema grayana</i> Hue. [DcmE]	NS	No scavenging activity	59.9%	NS	Thadhani et al. (2011)
	<i>Usnea complanata</i> (Müll. Arg.) Motyka [Total extract]	0.214 mg/ml	0.195 mg/ml	NS	NS	Behera et al. (2012)
	<i>Parmelia caperata</i> (L.) Ach. [AcE]	NS	60.7 µg/ml	97.3 µg/ml	500 µg/ml: 0.547	Manojlovic et al. (2012a)
	<i>Usnea barbata</i> Motyka [AcE]	NS	130.7 µg/ml	197.3 µg/ml	500 µg/ml: 0.664	Rankovic et al. (2012)
	<i>Cladonia lepidophora</i> Ahti & Kashiw	NS	No scavenging activity	NS	NS	Brisdelli et al. (2013)
	<i>Ochrolechia deceptivensis</i> (Hue) Darb.	NS	No scavenging activity	NS	NS	Brisdelli et al. (2013)
Variolaric acid	<i>Lobaria pulmonaria</i> L. (Hoffm.) [AcE]	NVNR	NS	NS	NS	Atalay et al. (2011)
Vesuvianic acid	<i>Psoroma pallidum</i> Nyl	NS	No scavenging activity	NS	NS	Brisdelli et al. (2013)
Vicianicin	<i>Cladonia</i> sp. [DcmE]	NS	No scavenging activity	1.6%	NS	Thadhani et al. (2011)
Zeorin						

NS, not studied; NVNR, numeric value not reported; LP, lipoperoxidation; %INH, percentage of inhibition; SO RSA, superoxide radical scavenging activity; GA, gallic acid; PE, pyrocatechol equivalent; RE, rutin equivalent; AA, ascorbic acid; T, tocopherol; TBA, thiobarbituric acid; CE, catechol equivalent; MDA, malondialdehyde; ChE, chloroform extract; ME, metanol extract; PE, petrol ether extract; AcE, acetone extract; DmE, dimethyl sulfoxide extract; EE, ethanol extract; WE, water extract; DE, diethyl-ether extract; BE, benzene extract; BuE, butanol extract; DcmE, dichloromethane extract; Eae, ethyl acetate extract.

Table 3. Evaluations of antioxidant parameters on cellular substrates.

Lichen specie/isolated compound	Solvent/origin	Cell line	LP inhibition	Antioxidant activity markers	Reference
Atranorin	Comercial	Human neuron-like cells (SH-SY5Y)	↑ Peroxyl radical-induced lipoperoxidation <i>in vitro</i>	Good antioxidant capacity in TRAP/TAR assays ↑ H ₂ O ₂ and NO production and SO ⁻ scavenging activity Protection of SH-SY5Y cells against H ₂ O ₂ -induced cell viability impairment	Melo et al. (2011)
<i>Cetraria islandica</i> (L.) Ach. Diffraitaic acid	Methanol <i>Protosnea magellanica</i> (Mont.) Krog	Human lymphocytes HeLa cell line	↓ MDA levels Not studied (NS)	↑ SOD and GPx activities No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Kotan et al. (2011) Brisdelli et al. (2013)
<i>Evernia prunastri</i> (L.) Ach. Lobaric acid	Methanol <i>Stereocaulon alpinum</i> Laurer ex Funck	Human lymphocytes HeLa cell line	↓ MDA levels NS	↑ GSH levels, SOD and GPx activities No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Alpsoy et al. (2015) Brisdelli et al. (2013)
<i>Peltigera horizontalis</i> (Hudson) Baumg.	Methanol	Human lymphocytes	↓ MDA levels	↑ GSH levels, SOD and GPx activities	Nardemir et al. (2013)
<i>Peltigera praetextata</i> (Flörke ex Sommerf.) Zopf	Methanol	Human lymphocytes	↓ MDA levels	↑ GSH levels, SOD and GPx activities	Nardemir et al. (2013)

Protolichesterric acid	<i>Cornicularia aculeata</i> (Schreb.) Ach.	HeLa cell line	NS	No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Brisdelli et al. (2013)
Ramalin	<i>Ramalina terebrata</i> Hook. f. & Taylor. [ME]	Murine macrophage Raw 264.7 cells	NS	↓ NO release and H ₂ O ₂ production after LPS stimulation	Paudel et al. (2011)
Salazinic acid	<i>Xanthoparmelia cantschadalis</i> (Ach.) Hale [ME]	Astrocyte cell line U373-MG	NS	↑ Cell viability in H ₂ O ₂ -treated cells ↓ ROS production ORAC value: 2.7 µmol TE/mg	de Paz et al. (2010)
Stictic acid	<i>Xanthoparmelia conspersa</i> (Ach.) Hale [ME]	Astrocyte cell line U373-MG	NS	↑ Cell viability in H ₂ O ₂ -treated cells ↓ ROS production ORAC value: 2.3 µmol TE/mg	de Paz et al. (2010)
<i>Umbilicaria vellea</i> (L.) Ach.	Methanol	Human lymphocytes	↓ MDA levels	↑ SOD and GPx activities (doses: 5 and 10 mg/ml)	Aslan et al. (2011)
Usnic acid (UA)	<i>Xanthoparmelia conspersa</i> (Ach.) Hale and <i>X. cantschadalis</i> (Ach.) Hale [ME] Comercial	Astrocyte cell line U373 MG	NS	↑ Cell viability in H ₂ O ₂ -treated cells ↓ ROS production ORAC value: 2.9 µmol TE/mg	de Paz et al. (2010)
		Human neuron-like cells (SH-SY5Y)	↑ Lipoperoxidation	Good antioxidant capacity in TRAP/TAR assays ↓ OH [•] and NO formation No effect on CAT and SOD-like activities No protection against H ₂ O ₂ -induced cell death	Rabelo et al. (2012)
Usnic acid (UA)	<i>Cladonia lepidophora</i> Ahti & Kashiw.	HeLa cell line	NS	↑ ROS production No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Brisdelli et al. (2013)
Variolaric acid	<i>Ochrolechia deceptionis</i> (Hue) Darb.	HeLa cell line	NS	No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Brisdelli et al. (2013)
Vicanicin	<i>Psoroma pallidum</i> Nyl.	HeLa cell line	NS	No effect on intracellular ROS level No protection against t-BHP-induced increase in intracellular ROS level	Brisdelli et al. (2013)
<i>Xantho somloensis</i> (Gyelnik) Hale.	Methanol	Human lymphocytes	↓ MDA levels	↑ SOD and GPx activities (doses: 5 and 10 mg/ml)	Aslan et al. (2011)
<i>Xanthoparmelia cantschadalis</i> (Ach.) Hale	Methanol	Astrocyte cell line U373 MG	NS	↑ Cell viability in H ₂ O ₂ -treated cells ↓ ROS production ORAC value: 4.9 µmol TE/mg	de Paz et al. (2010)
<i>Xanthoparmelia conspersa</i> (Ach.) Hale	Methanol	Astrocyte cell line U373 MG	NS	↑ Cell viability in H ₂ O ₂ -treated cells ↓ ROS production ORAC value: 8.8 µmol TE/mg	de Paz et al. (2010)
<i>Xanthoria elegans</i> (Link) Th. Fr.	Water	Human lymphocytes	NS	↑ TAC level without changing TOS level	Turkez et al. (2011)

TBARS, thiobarbituric acid reactive substances; SOD, superoxide dismutase; GPx, glutathione peroxidase; GSH, reduced glutathione; AFB₁, aflatoxin B₁; AE, water extract; ME, methanol extract; ROS, reactive oxygen species; OH[•], hydroxyl radical; CAT, catalase; TAC, total antioxidant capacity; TOS, total oxidative stress; MPx, myeloperoxidase; cNOS, constitutive nitric oxide synthase; iNOS, inducible nitric oxide synthase; GR, glutathione reductase; GST, glutathione transferase; DE, diethyl ether extract; AcE, acetone extract.

Table 4. *In vivo* evaluations of antioxidant activities.

Lichen specie/isolated compound	Solvent/origin	LP inhibition	Antioxidant enzymes and other antioxidant markers	Reference
<i>Cetraria islandica</i> (L.) Ach.	–	↓ MDA levels	↑ GSH levels ↓ CAT and GPx activities Association of magnesium augmented the antioxidant effect	Cernescu et al. (2011)
Diffraicta acid	<i>Usnea longissima</i> Ach. [DE]	↓ lipoperoxidation level in tissues	↑ SOD and GPx activities and GSH levels ↓ CAT and MPx activities	Bayir et al. (2006)
Fumarprotocetraric acid	<i>Cladonia verticillaris</i> (Raddi) Fr. [AcE]	↓ endotoxin-induced lipid peroxidation	↑ cNOS activity and ↓ iNOS activity Not studied (NS)	de Barros Alves et al. (2014)
<i>Lobaria pulmonaria</i> (L.) Hoffm.	Methanol	↓ lipoperoxidation level in tissues	↑ SOD, GPx and GSH levels in tissues	Karakus et al. (2009)
<i>Peltigera rufescens</i> (Weiss) Humb.	Methanol	↓ lipoperoxidation	No effect on CAT and MPx levels	Tanas et al. (2010)
<i>Usnea ghattensis</i> G. Awasthi	Methanol	↓ MDA formation in liver tissue	↑ SOD, CAT, GSH, GR and GPx levels ↓ MPx and iNOS activities	Verma et al. (2008a)
<i>Usnea longissima</i> Ach.	Water	47.1% inhibition	Depletion in antioxidant enzymes (SOD, CAT, GPx) and GSH	Halici et al. (2005)
Usmic acid	<i>Usnea longissima</i> Ach. [DE]	↓ lipoperoxidation level in tissues	↑ SOD and GST levels ↓ CAT activity. Reducing power (A_{700}): 0.1; phenolic content: 18.3 mg GA/g	Odabasoglu et al. (2006)

TBARS, thiobarbituric acid reactive substances; SOD, superoxide dismutase; GPx, glutathione peroxidase; GSH, reduced glutathione; AFB₁, aflatoxin B₁; AE, water extract; ME, methanol extract; ROS, reactive oxygen species; OH[•], hydroxyl radical; CAT, catalase; TAC, total antioxidant capacity; TOS, total oxidative stress; MPx, myeloperoxidase; cNOS, constitutive nitric oxide synthase; iNOS, inducible nitric oxide synthase; GR, glutathione reductase; GST, glutathione transferase; DE, diethyl ether extract; AcE, acetone extract.

Conclusions

The present review reports the biological activity of more than 75 different lichen species, as well as more than 65 purified metabolites, isolated from these or other species. The study of their antioxidant activities has recently been started and they have been determined by various chemical *in vitro* assays as first approach, with some of them showing interesting results. Further knowledge of this potential implies deeper research on their activities in order to understand the implied mechanisms. Thus, in this report, we also reflect the few available data about *in vitro* antioxidant activities of some lichen species and purified metabolites on cellular substrates and *in vivo* on animal models.

Concerning antioxidation, the most interesting compounds are polyphenols. The antioxidant properties of polyphenols are due to the presence of their many phenolic hydroxyl groups, which confer high potential for scavenging free radicals (Dai & Mumper, 2010; Sawa et al., 1999). For instance, phenolic compounds are able to donate hydrogen to reactive radicals and break the chain reaction of lipid oxidation at the initiation step (Gülçin et al., 2004).

Then, the strong antioxidant activity shown by some lichen extracts or metabolites, and assessed by different systems, can be attributed to their high total polyphenolic contents (specially depsides, depsidones, dibenzofurans, etc.), since a positive correlation between phenolic composition and antioxidant activity has been proved for most of them (Kosanić et al., 2011; Manojlovic et al. 2012c); at least, it suggests that polyphenols might be the major antioxidant compounds in studied lichens. Nevertheless, there have been other studies in which results did not show any positive correlation between antioxidant activity of certain lichens and total phenolic contents (Odabasoglu et al., 2004; Stojanović et al., 2010). This fact implies that other minor compounds should not be ignored but antioxidant activity might be as well attributed to the presence of non-phenolic compounds, antagonistic or synergistic interactions between constituents, and even distinct antioxidant activities of individual phenolics.

In the previous tables, the great diversity of lichens and their substances is shown, and one might deduce that the increasing interest in the study of its pharmacological properties is promoting further phylogenetic studies in an evolutionary context. Based on molecular data mainly, they are leading to a more complex classification of lichen families and species (Crespo et al., 2010). Moreover, phylogenetic analysis of biosynthetic genes can facilitate the discovery of novel compounds, novel genes, and, therefore, unknown producers of pharmaceutical relevant compounds, including antioxidants: the greatest challenges would be to find the biosynthetic gene of interest or assign function to each of the biosynthetic genes found in a lichen genome (Schmitt & Barker, 2009). Considering the difficulties still found for the *in vitro* culture of lichens and different culture conditions result in different antioxidant activities of lichen extracts (due to the production of different amount and type of secondary metabolites depending on culture characteristics) (Behera et al., 2005b), a global approach to the lichen metabolomic features seems to be crucial for the development of new and viable biotechnological processes. These will allow

production of suitable amounts of unique isolated antioxidant compounds from lichens (Boustie & Grube, 2005).

Through this review of literature, we can conclude that lichens are a potential source of natural antioxidants but, at the same time, there is still a need for a deeper research in order to establish their possibilities and a better understanding of their mechanisms of action. This goal can be achieved by the better isolation of purified metabolites and more studies on appropriate cell lines and *in vivo*, in order to identify molecular targets, active compounds, and structure–activity correlations.

Declaration of interest

The authors report that they have no conflicts of interest. The authors alone are responsible for the content and writing of this article. This work was supported by a Doctoral Grant from the Spanish Ministry of Education, Culture and Sports (FPU programme), awarded to Carlos Fernández Moriano (No. FPU12/03824).

References

- Alpsoy L, Orhan F, Nardemir G, et al. (2015). Antigenotoxic potencies of a lichen species, *Evernia prunastri*. *Toxicol Ind Health* 31:153–61.
- Aslan A, Agar G, Alpsoy L, et al. (2011). Protective role of methanol extracts of two lichens on oxidative and genotoxic damage caused by AFB1 in human lymphocytes *in vitro*. *Toxicol Ind Health* 28:505–12.
- Atalay F, Halici MB, Mavi A, et al. (2011). Antioxidant phenolics from *Lobaria pulmonaria* (L.) Hoffm. and *Usnea longissima* Ach. lichen species. *Turk J Chem* 35:647–66.
- Bayir Y, Odabasoglu F, Cakir A, et al. (2006). The inhibition of gastric mucosal lesion, oxidative stress and neutrophil-infiltration in rats by the lichen constituent diffractaic acid. *Phytomedicine* 13:584–90.
- Behera BC, Adawadkar B, Makhija U. (2003). Inhibitory activity of xanthine oxidase and superoxide-scavenging activity in some taxa of the lichen family Graphidaceae. *Phytomedicine* 10:536–43.
- Behera BC, Adawadkar B, Makhija U. (2006a). Tissue-culture of selected species of the *Graphis* lichen and their biological activities. *Fitoterapia* 77:208–15.
- Behera BC, Adawadkar B, Makhija U. (2006b). Tyrosinase-inhibitory activity in some species of the lichen family Graphidaceae. *J Herb Pharmacother* 6:55–69.
- Behera BC, Mahadik N, Morey M. (2012). Antioxidative and cardiovascular-protective activities of metabolite usnic acid and psoromic acid produced by lichen species *Usnea complanata* under submerged fermentation. *Pharm Biol* 50:968–79.
- Behera BC, Verma N, Sonone A, Makhija U. (2005a). Antioxidant and antibacterial activities of lichen *Usnea ghattensis* *in vitro*. *Biotechnol Lett* 27:991–5.
- Behera BC, Verma N, Sonone A, Makhija U. (2005b). Evaluation of antioxidant potential of the cultured mycobiont of a lichen *Usnea ghattensis*. *Phytother Res* 19:58–64.
- Behera BC, Verma N, Sonone A, Makhija U. (2006c). Determination of antioxidative potential of lichen *Usnea ghattensis* *in vitro*. *LWT* 39: 80–5.
- Bhattarai HD, Paudel B, Hong SG, et al. (2008). Thin layer chromatography analysis of antioxidant constituents of lichens from Antarctica. *J Nat Med* 62:481–4.
- Boustie J, Grube M. (2005). Lichens: A promising source of bioactive secondary metabolites. *Plant Genet Resour* 3:273–87.
- Brisdelli F, Perilli M, Sellitri D, et al. (2013). Cytotoxic activity and antioxidant capacity of purified lichen metabolites: An *in vitro* study. *Phytother Res* 27:431–7.
- Brodo MI, Sharnon SD, Sharnon S. (2001). *Lichens of North America*. New Haven, London: Yale University Press.
- Buçukoglu TZ, Albayrak S, Halici MG, Tay T. (2012). Antimicrobial and antioxidant activities of extracts and lichen acids obtained from some Umbilicaria species from Central Anatolia, Turkey. *J Food Process Pres* 37:1103–10.

- Cernescu I, Tarțău L, Macavei A, Lupușoru CE. (2011). Experimental research on the effects of a *Cetraria islandica* extract on oxidative stress in laboratory animals. *Rev Med Chir Soc Med Nat Iasi* 115: 899–904.
- Crespo A, Kauff F, Divakar PK, et al. (2010). Phylogenetic generic classification of parmelioid lichens (Parmeliaceae, Ascomycota) based on molecular, morphological and chemical evidence. *Taxon* 59:1735–53.
- Dai J, Mumper RJ. (2010). Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules* 15:7313–52.
- de Barros Alves GM, de Sousa Maia MB, de Souza Franco E, et al. (2014). Expectorant and antioxidant activities of purified fumarprotocetraric acid from *Cladonia verticillaris* lichen in mice. *Pulm Pharmacol Ther* 27:139–43.
- de Paz G, Raggio J, Gómez-Serranillos MP, Palomino OM, et al. (2010). HPLC isolation of antioxidant constituents from *Xanthoparmelia* spp. *J Pharm Biomed Anal* 53:165–71.
- Dias DA, Urban S. (2009). Phytochemical investigation of the Australian lichens *Ramalina glaucescens* and *Xanthoria parietina*. *Nat Prod Commun* 4:959–64.
- Einarsdóttir E, Groeneweg J, Björnsdóttir GG, et al. (2010). Cellular mechanisms of the anticancer effects of the lichen compound usnic acid. *Planta Med* 76:969–74.
- Esimone CO, Ofokansi KC, Adikwu MU, et al. (2007). *In vitro* evaluation of the antiviral activity of extracts from the lichen *Parmelia perlata* (L.) Ach. against three RNA viruses. *J Infect Dev Ctries* 1: 315–20.
- Gonzalez-Burgos E, Carretero ME, Gomez-Serranillos MP. (2012). Diterpenoids isolated from *Sideritis* species protect astrocytes against oxidative stress via Nrf2. *J Nat Prod* 75:1750–8.
- Gonzalez-Burgos E, Carretero ME, Gómez-Serranillos MP. (2013). Kaurane diterpenes from *Sideritis* spp. exert a cytoprotective effect against oxidative injury that is associated with modulation of the Nrf2 system. *Phytochemistry* 93:116–23.
- Grice HC. (1986). Enhanced tumour development by butylated hydroxytoluene (BHT) in the liver, lung and gastro-intestinal tract. *Food Chem Toxicol* 24:1127–30.
- Gülçin I, Beydemir S, Alici HA, et al. (2004). *In vitro* antioxidant properties of morphine. *Pharmacol Res* 49:59–66.
- Gülçin I, Oktay M, Küfrevioğlu I, Aslan A. (2002). Determination of antioxidant activity of lichen *Cetraria islandica* (L.) Ach. *J Ethnopharmacol* 79:325–9.
- Güllüce M, Aslanc A, Sokmend M, et al. (2006). Screening the antioxidant and antimicrobial properties of the lichens *Parmelia saxatilis*, *Platismatia glauca*, *Ramalina pollinaria*, *Ramalina polymorpha* and *Umbilicaria nylanderiana*. *Phytomedicine* 13:515–21.
- Güvenç A, Akkol EK, Süntar I, et al. (2012). Biological activities of *Pseudevernia furfuracea* (L.) Zopf extracts and isolation of the active compounds. *J Ethnopharmacol* 144:726–34.
- Halici M, Odabasoglu F, Suleyman H, et al. (2005). Effects of water extract of *Usnea longissima* on antioxidant enzyme activity and mucosal damage caused by indomethacin in rats. *Phytomedicine* 12: 656–62.
- Hara K, Endo M, Kawakami H, et al. (2011). Anti-oxidation activity of ethanol extracts from natural thalli of lichens. *Mycosystema* 30: 950–54.
- Hawksworth DL, Honegger R. (1994). The lichen thallus: A symbiotic phenotype of nutritionally specialized fungi and its response to gall producers. In: Williams MAJ, ed. *Plant Galls*. Special vol. 49. Oxford: The Systematics Association, 77–98.
- Hidalgo ME, Fernández E, Quilhot W, Lissi E. (1994). Antioxidant activity of depsides and depsidones. *Phytochemistry* 37: 1585–7.
- Honda NK, Vilegas W. (1999). The chemistry of lichens. *Quimica Nova* 22:110–25.
- Huneck S. (1999). The significance of lichens and their metabolites. *Naturwissenschaften* 86:559–70.
- Jayaprakasha GK, Rao LJ. (2000). Phenolic constituents from the lichen *Parmotrema stippeum* (Nyl.) Hale and their antioxidant activity. *Z Naturforsch* 55:1018–22.
- Karakus B, Odabasoglu F, Cakir A, et al. (2009). The effects of methanol extract of *Lobaria pulmonaria*, a lichen species, on indomethacin-induced gastric mucosal damage, oxidative stress and neutrophil infiltration. *Phytother Res* 23:635–9.
- Kim MS, Cho HB. (2007). Melanogenesis inhibitory effects of methanolic extracts of *Umbilicaria esculenta* and *Usnea longissima*. *J Microbiol* 45:578–82.
- Kinoshita K, Togawa T, Hiraishi A, et al. (2010). Antioxidant activity of red pigments from the lichens *Lethariella sernanderi*, *L. cashmeriana*, and *L. sinensis*. *J Nat Med* 64:85–8.
- Kosanic M, Manojlovic N, Jankovic S, et al. (2013). *Evernia prunastri* and *Pseudevernia furfuracea* lichens and their major metabolites as antioxidant, antimicrobial and anticancer agents. *Food Chem Toxicol* 53:112–18.
- Kosanić M, Ranković B, Stanojković T. (2012a). Antioxidant, antimicrobial and anticancer activities of three *Parmelia* species. *J Sci Food Agric* 92:1909–16.
- Kosanić M, Ranković B, Stanojković TJ. (2012b). Antioxidant, antimicrobial, and anticancer activity of 3 *Umbilicaria* species. *Food Sci* 71:20–5.
- Kosanić M, Ranković B, Vukojević J. (2011). Antioxidant properties of some lichen species. *J Food Sci Technol* 48:584–90.
- Kosanić M, Ranković B. (2011). Lichens as possible sources of antioxidants. *Pak. J Pharm Sci* 24:165–70.
- Kotan E, Alpsoy L, Anar M, et al. (2011). Protective role of methanol extract of *Cetraria islandica* (L.) against oxidative stress and genotoxic effects of AFB1 in human lymphocytes *in vitro*. *Toxicol Ind Health* 27:599–605.
- Lohézic-Le Dévéhat F, Tomasi S, Elix JA, et al. (2007). Stictic acid derivatives from the lichen *Usnea articulata* and their antioxidant activities. *J Nat Prod* 70:1218–20.
- Lopes TI, Coelho R, Yoshida N, Honda NK. (2008). Radical-scavenging activity of *Orsellinates*. *Chem Pharm Bull* 56:1551–4.
- Luo H, Ren M, Lim K, et al. (2006). Antioxidative activity of lichen *Thamnia vermicularis* *in vitro*. *Mycobiology* 34:124–7.
- Luo H, Yamamoto Y, Kim JA, et al. (2009). Lecanoric acid, a secondary lichen substance with antioxidant properties from *Umbilicaria antarctica* in maritime Antarctica (King George Island). *Polar Biol* 32:1033–40.
- Luo H, Yamamoto Y, Liu Y, et al. (2010). The *in vitro* antioxidant properties of Chinese highland lichens. *J Microbiol Biotechnol* 20: 1524–8.
- Malhotra S, Subban R, Singh A. (2008). Lichens – Role in traditional medicine and drug discovery. *Internet J Altern Med* 5:1–5.
- Manojlovic NT, Ranković B, Kosanić M, et al. (2012a). Chemical composition of three *Parmelia* lichens and antioxidant, antimicrobial and cytotoxic activities of some their major metabolites. *Phytomedicine* 19:1166–72.
- Manojlovic NT, Vasiljevic PJ, Gritsanapan W, et al. (2010). Phytochemical and antioxidant studies of *Laurera benguelensis* growing in Thailand. *Biol Res* 43:169–76.
- Manojlovic NT, Vasiljevic PJ, Maskovic PZ. (2012b). Chemical composition and antioxidant activity of lichen *Toninia candida*. *Rev Bras Farmacogn* 22:291–8.
- Manojlovic NT, Vasiljevic PJ, Maskovic PZ, et al. (2012c). Chemical composition, antioxidant, and antimicrobial activities of lichen *Umbilicaria cylindrica* (L.) Delise (Umbilicariaceae). *Evid Based Complement Alternat Med* 2012:452431.
- Melo MG, dos Santos JP, Serafini MR, et al. (2011). Redox properties and cytoprotective actions of atranorin, a lichen secondary metabolite. *Toxicol In Vitro* 25:462–8.
- Mitrović T, Stamenković S, Cvetković V, et al. (2011). Antioxidant, antimicrobial and antiproliferative activities of five lichen species. *Int J Mol Sci* 12:5428–48.
- Molnár K, Farkas E. (2010). Current results on biological activities of lichen secondary metabolites: A review. *Z Naturforsch* 65c:157–73.
- Nardemir G, Yanmis D, Alpsoy L, et al. (2013). Genotoxic, antigenotoxic and antioxidant properties of methanol extracts obtained from *Peltigera horizontalis* and *Peltigera praetextata*. *Toxicol Ind Health*. [Epub ahead of print]. doi: 10.1177/0748233713480207.
- Nishitoba Y, Nishimura I, Nishiyama T, Mizutani J. (1987). Lichen acids, plant growth inhibitors from *Usnea longissima*. *Phytochemistry* 26:3181–5.
- Odabasoglu F, Aslan A, Cakir A, et al. (2004). Comparison of antioxidant activity and phenolic content of three lichen species. *Phytother Res* 18:938–41.
- Odabasoglu F, Aslan A, Cakir A, et al. (2005). Antioxidant activity, reducing power and total phenolic content of some lichen species. *Fitoterapia* 76:216–19.

- Odabasoglu F, Cakir A, Suleyman H, et al. (2006). Gastroprotective and antioxidant effects of usnic acid on indomethacin-induced gastric ulcer in rats. *J Ethnopharmacol* 103:59–65.
- Özen T, Kinalioglu K. (2008). Determination of antioxidant activity of various extracts of *Parmelia saxatilis*. *Biologia* 63:211–16.
- Papadopolou P, Tzakou O, Vagias C, et al. (2007). β -Orcinol metabolites from the lichen *Hypotrachyna revolute*. *Molecules* 12:997–1005.
- Paudel B, Bhattarai HD, Koh HY, et al. (2011). Ramalin, a novel nontoxic antioxidant compound from the Antarctic lichen *Ramalina terebrata*. *Phytomedicine* 18:1285–90.
- Paudel B, Bhattarai HD, Lee JS, et al. (2008). Antioxidant activity of polar lichens from King George Island (Antarctica). *Polar Biol* 31: 605–8.
- Prashith Kekuda TR, Vinayaka KS, Praveen Kumar SV, Sudharshan SJ. (2009). Antioxidant and antibacterial activity of lichen extracts, honey and their combination. *J Pharm Res* 2:1875–78.
- Praveen Kumar SV, Prashith Kekuda TR, Vinayaka KS, et al. (2010). Studies on antibacterial, anthelmintic and antioxidant activities of a Macrolichen *Parmotrema pseudotinctorum* (des. Abb.) Hale (Parmeliaceae) from Bhadra wildlife sanctuary, Karnataka. *Int J Pharm Tech Res* 2:1207–14.
- Rabelo TK, Zeidán-Chuliá F, Vasques L, et al. (2012). Redox characterization of usnic acid and its cytotoxic effect on human neuron-like cells (SH-SY5Y). *Toxicol In Vitro* 26:304–14.
- Racine PH, Hartmann VE, Tollard D'Audiffret Y. (1980). Antioxidant properties of wax from Yugoslavian oakmoss (*Evernia prunastri*). *Int J Cosmet Sci* 2:305–13.
- Ranković B, Kosanić M, Stanojković T. (2011). Antioxidant, antimicrobial and anticancer activity of the lichens *Cladonia furcata*, *Lecanora atra* and *Lecanora muralis*. *BMC Complement Altern Med* 11:97.
- Ranković B, Kosanić M, Stanojković T, et al. (2012). Biological activities of *Toninia candida* and *Usnea barbata* together with their norstictic acid and usnic acid constituents. *Int J Mol Sci* 13: 14707–22.
- Ranković B, Ranković D, Kosanić M, Marić D. (2010a). Antioxidant and antimicrobial properties of the lichens *Anaptychia ciliaris*, *Nephroma parile*, *Ochrolechia tartarea* and *Parmelia centrifuga*. *Cent Eur J Biol* 5:649–55.
- Ranković B, Ranković D, Marić D. (2010b). Antioxidant and antimicrobial activity of some lichen species. *Microbiology* 79:809–15.
- Sawa T, Nakao M, Akaike T, et al. (1999). Alkylperoxyl radical-scavenging activity of various flavonoids and other phenolic compounds: Implications for the anti-tumor-promoter effect of vegetables. *J Agric Food Chem* 47:397–402.
- Sayre LM, Perry G, Smith MA. (2008). Oxidative stress and neurotoxicity. *Chem Res Toxicol* 21:172–88.
- Schmitt I, Barker FK. (2009). Phylogenetic methods in natural product research. *Nat Prod Rep* 26:1585–602.
- Silva JA, Bomfim RR, Estevam Cdos S, et al. (2010). Pharmacological properties of lichen *Cladonia clathrata*. *Pharm Biol* 48:745–52.
- Singh SM, Singh P, Ravindra R. (2011). Screening of antioxidant potential of Arctic lichens. *Polar Biol* 34:1775–82.
- Stojanović G, Stojanović I, Stankov-Jovanović V, et al. (2010). Reducing power and radical scavenging activity of four Parmeliaceae species. *Cent Eur J Biol* 5:808–13.
- Sundararajan R, Ahamad NH, Venkatesan K, et al. (2006). *Cytisus scoparius* link – A natural antioxidant. *BMC Complement Altern Med* 6:8.
- Takenaka Y, Tanahashi T, Nagakura N, Hamada N. (2000). Production of xanthenes with free radical scavenging properties, emodin and sclerotiorin by the cultured lichen mycobionts of *Pyrenula japonica*. *Z Naturforsch* 55c:910–14.
- Tanas S, Odabasoglu F, Halici Z, et al. (2010). Evaluation of anti-inflammatory and antioxidant activities of *Peltigera rufescens* lichen species in acute and chronic inflammation models. *J Nat Med* 64: 42–9.
- Thadhani VM, Choudhary MI, Ali S, et al. (2011). Antioxidant activity of some lichen metabolites. *Nat Prod Res* 25:1827–37.
- Toledo-Marante FJ, García Castellano A, Estevez Rosas F, et al. (2003). Identification and quantitation of allelochemicals from the lichen *Lethariella canariensis*: Phytotoxicity and antioxidative activity. *J Chem Ecol* 29:2049–71.
- Turkez H, Aydin E, Aslan A. (2011). *Xanthoria elegans* (Link) (lichen) extract counteracts DNA damage and oxidative stress of mitomycin C in human lymphocytes. *Cytotechnology* 64:679–86.
- Valencia-Islas N, Zambrano A, Rojas JL. (2007). Ozone reactivity and free radical scavenging behavior of phenolic secondary metabolites in lichens exposed to chronic oxidant air pollution from Mexico City. *J Chem Ecol* 33:1619–34.
- Verma N, Behera BC, Makhija U. (2008a). Antioxidant and hepatoprotective activity of a lichen *Usnea ghattensis* in vitro. *Appl Biochem Biotechnol* 151:167–81.
- Verma N, Behera BC, Sonone A, Makhija U. (2008b). Lipid peroxidation and tyrosinase inhibition by lichen symbionts grown in vitro. *African J Biochem Res* 2:225–31.
- Vinayaka KS, Praveen Kumar SV, Prashith Kekuda TR, et al. (2009). Proximate composition, antioxidant, anthelmintic and insecticidal activity of a macrolichen *Ramalina conduplicans* Vain. (*Ramalinaceae*). *European J App Sci* 1:40–6.
- Wang JY, Yang JY, Wang F, et al. (2013). Neuroprotective effect of pseudoginsenoside-f11 on a rat model of Parkinson's disease induced by 6-hydroxydopamine. *Evid Based Complement Alternat Med* 2013:152798.
- Wichi, HP. (1988) Safety evaluation of butylated hydroxyanisole from the perspective of effects on forestomach and oesophageal squamous epithelium. *Food Chem Toxicol* 26:717–23.
- Yücel O, Odabasoglu F, Güllüce M, et al. (2007). Antioxidant and antimicrobial properties of a lichen species *Cladonia rangiformis* growing in Turkey. *Turkish J Pharm Sci* 4:101–9.
- Zambare VP, Christopher LP. (2012). Biopharmaceutical potential of lichens. *Pharm Biol* 50:778–98.