



## Antioxidant effect of functionalized alkyl-organotellurides: a study *in vitro*

Bibiana M. Gay, Cristiane Luchese, Cristina W. Nogueira, P. Wendler, Alexandra Macedo & Alcindo A. Dos Santos

**To cite this article:** Bibiana M. Gay, Cristiane Luchese, Cristina W. Nogueira, P. Wendler, Alexandra Macedo & Alcindo A. Dos Santos (2010) Antioxidant effect of functionalized alkyl-organotellurides: a study *in vitro*, Journal of Enzyme Inhibition and Medicinal Chemistry, 25:4, 467-475, DOI: [10.3109/14756360903257892](https://doi.org/10.3109/14756360903257892)

**To link to this article:** <https://doi.org/10.3109/14756360903257892>



Published online: 17 Mar 2010.



Submit your article to this journal [↗](#)



Article views: 505



View related articles [↗](#)

ORIGINAL ARTICLE

# Antioxidant effect of functionalized alkyl-organotellurides: a study *in vitro*

Bibiana M. Gay<sup>1</sup>, Cristiane Luchese<sup>1</sup>, Cristina W. Nogueira<sup>1</sup>, P. Wendler<sup>2</sup>, Alexandra Macedo<sup>2</sup>, and Alcindo A. Dos Santos<sup>3</sup>

<sup>1</sup>Department of Chemistry, Federal University of Santa Maria, Santa Maria, RS, Brazil, <sup>2</sup>Department of Chemistry, Federal University of São Carlos, São Carlos, SP, Brazil, and <sup>3</sup>Institute of Chemistry, University of São Paulo, São Paulo, SP, Brazil

## Abstract

We evaluated the *in vitro* antioxidant effect of alkyl-organotellurides **A–D** on lipid peroxidation and protein carbonylation in rat liver homogenates. The thiol oxidase and thiol peroxidase-like activities of compounds were investigated.  $\delta$ -Aminolevulinic acid dehydratase ( $\delta$ -ALA-D) activity was determined in rat liver homogenates. Compounds **A–D** protected against lipid peroxidation induced by Fe<sup>2+</sup>/EDTA and sodium nitroprusside (SNP). According to the confidence limits of the IC<sub>50</sub> values of compounds **A–D**, the IC<sub>50</sub> values for organotellurides followed the order: **C** (0.30  $\mu$ M)  $\leq$  **B** (0.40  $\mu$ M)  $<$  **D** (0.68  $\mu$ M)  $<$  **A** (2.90  $\mu$ M), for Fe<sup>2+</sup>/EDTA, and **B** (0.21  $\mu$ M)  $\leq$  **C** (0.33  $\mu$ M)  $\leq$  **D** (0.43  $\mu$ M)  $<$  **A** (1.21  $\mu$ M) for SNP-induced lipid peroxidation. Compounds **A–D** reduced protein carbonyl content to control levels. The results demonstrated an inverse correlation between thiol oxidase and  $\delta$ -ALA-D activities. This study supports an antioxidant effect of organotellurides **A–D** on rat liver.

**Keywords:** Functionalized alkyl-organotellurides; antioxidant; tellurium; liver lipid peroxidation

## Introduction

Evidence has been provided that organochalcogenides, such as organoselenium and organotellurium compounds, are promising pharmacological agents<sup>1–4</sup>. Organotellurium compounds have been synthesized since 1840<sup>5</sup>, and it is known that these compounds are much more effective bioactive agents than their corresponding selenium and sulfur analogs, with evident chemoprotective, antioxidative, and cytotoxic properties<sup>3,6</sup>, making these compounds extremely attractive in medical therapies. However, little is known about the pharmacological and toxicological effects of these compounds<sup>7,8</sup>. Engman *et al.*<sup>1,9</sup> have investigated the possible anti-cancer properties of organotellurium compounds.

Several authors have reported that organotellurium compounds can exhibit antioxidant activity because of their glutathione peroxidase (GPx) mimetic properties<sup>10</sup> and peroxynitrite scavenger abilities<sup>11</sup>. Moreover, studies have shown an antioxidant activity *in vitro*<sup>11–13</sup> and low toxicity *ex vivo*<sup>11,13,14</sup> of organotellurium compounds.

Oxidative stress is characterized by a significantly increased concentration of intracellular oxidizing species,

such as reactive oxygen species (ROS), and is often accompanied by the simultaneous loss of antioxidant defense capacity<sup>15,16</sup>. In pathologic conditions an overproduction of ROS can occur<sup>17</sup>. Besides that, ROS overproduction has been implicated in the installation and/or progression of a variety of human diseases<sup>18,19</sup> including diabetes<sup>20</sup>, cancer<sup>21</sup>, and various neurodegenerative diseases<sup>22</sup>. Thus, interest in the synthesis of antioxidant compounds that could potentially retard the development of diseases associated with oxidative stress has grown considerably in recent decades<sup>23,24</sup>.

In this study, we evaluated the *in vitro* antioxidant effect of functionalized alkyl-organotellurides **A–D** against lipid peroxidation and protein carbonylation in rat liver homogenates. In addition, we examined the thiol peroxidase-like activity of functionalized alkyl-organotellurides **A–D** to determine the possible mechanism by which these compounds demonstrate antioxidant property. Furthermore,  $\delta$ -aminolevulinic acid dehydratase ( $\delta$ -ALA-D) and thiol oxidase activities were determined in rat liver homogenates.

Address for Correspondence: Cristina Wayne Nogueira, Centro de Ciências Naturais e Exatas, Universidade Federal de Santa Maria, Santa Maria, CEP 97105-900, RS, Brazil. Tel: 55-55-3220-8140. Fax: 55-55-3220-8978. E-mail: criswn@quimica.ufsm.br

(Received 8 May 2009; revised 17 July 2009; accepted 5 August 2009)

ISSN 1475-6366 print/ISSN 1475-6374 online © 2010 Informa UK Ltd  
DOI: 10.3109/14756360903257892

<http://www.informahealthcare.com/enz>

## Materials and methods

### Synthesis of alkyl-organotellurides

Telluro-ketone **A** was synthesized by hydrotelluration of methyl vinyl ketone in 86% isolated yield by reacting elemental tellurium with *n*-butyllithium, followed by the addition of deoxygenated water and the enone as reported previously<sup>25,26</sup>. Tellurides **B** and **C** were prepared in good yields according to our previous protocols<sup>27–31</sup> from telluride **A**. Finally, telluride **D** was prepared (65% isolated yield) from compound **A** by reacting with hydroxylamine hydrochloride<sup>32</sup>. Scheme 1 summarizes the preparation of the mentioned compounds.

### 4-(Butyltellanyl)butan-2-one (A)

In a two-necked 50 mL flask under a nitrogen atmosphere were added sequentially elemental tellurium (0.511 g, 4.0 mmol) and dry tetrahydrofuran (THF; 5 mL). To the resulting mixture under vigorous stirring, at room temperature, was added *n*-butyllithium (4.0 mmol, 1.6 mol L<sup>-1</sup> solution in hexane, 1.0 eq.). To the resulting pale yellow clean solution was added deoxygenated water (0.18 mL, 10 mmol). Methyl vinyl ketone (0.280 g, 4.0 mmol) was added in a portion at room temperature with vigorous stirring. The resulting mixture was stirred for 1 h at room temperature then quenched with water and diluted with hexane/ethyl acetate (5 mL, 1:1). The organic and aqueous phases were separated and the organic phase was washed with NH<sub>4</sub>Cl solution (3 × 15 mL) and brine (3 × 15 mL) and then dried with MgSO<sub>4</sub>. The solution was filtered and the solvent was evaporated. The residue was purified by silica gel column chromatography, eluting first with hexane and then with hexane/ethyl acetate (10:1). Yield: 0.87 g (86%). <sup>1</sup>H NMR: (300 MHz, CDCl<sub>3</sub>, ppm) δ 0.92 (t, *J* = 7.4 Hz, 3H), 1.38 (sext, *J* = 7.4 Hz, 2H), 1.73 (q, *J* = 7.4 Hz, 2H), 2.15 (s, 3H), 2.69 (m, 4H), 3.03 (t, *J* = 7.4 Hz, 2H). <sup>13</sup>C NMR: (75 MHz, CDCl<sub>3</sub>, ppm) δ 207.8, 46.4, 34.3, 29.8, 25.0, 13.3, 3.4, -7.0. <sup>125</sup>Te NMR: (157 MHz, CDCl<sub>3</sub>, ppm, (PhSe)<sub>2</sub>) δ 279.6. LRMS: *m/z* (70 eV, rel. int.): 258 (M<sup>+</sup>, 7), 71 (40), 57 (24), 55 (26), 43 (100). IR: (ZnSe, cm<sup>-1</sup>) 2955, 2925, 2865, 1715. Anal. Calc.: C, 37.56; H, 6.30. Found: C, 37.32; H, 6.02%.

### 4-(Butyltelluro)butan-2-ol (B)


To a 125 mL round-bottomed flask were added telluride **A** (1.28 g, 5 mmol) and ethanol (20 mL). To the resulting solution was added an ethanolic solution of NaBH<sub>4</sub> (0.21 g, 5.5 mmol in 6 mL ethanol). The mixture was stirred at room temperature and the progress of the reaction was monitored by thin layer chromatography (TLC). The reaction mixture was quenched with water (50 mL) and diluted with a mixture of hexane/ethyl acetate 1:1 (50 mL). The phases were separated and the aqueous phase was extracted with a mixture of hexane/ethyl acetate (1:1, 2 × 50 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, and filtered and the solvents were removed under reduced pressure. The residue was purified by silica gel column chromatography eluting with a mixture of hexane/ethyl acetate (15:1). Yield: 0.89 g (70%). <sup>1</sup>H NMR: (200 MHz, CDCl<sub>3</sub>, ppm) δ 0.85 (t, *J* = 7.2 Hz, 3H), 1.13 (d, *J* = 6.3 Hz, 3H), 1.31 (sext, *J* = 7.2 Hz, 2H), 1.65 (quint, *J* = 7.2 Hz, 2H), 1.76–1.85 (m, 2H), 2.53–2.69 (m, 4H), 3.75 (sext, *J* = 6 Hz, 1H). <sup>13</sup>C NMR: (50 MHz, CDCl<sub>3</sub>, ppm) δ 69.1, 41.1, 34.2, 25.0, 23.2, 13.4, 2.7, -2.3. <sup>125</sup>Te NMR: (157 MHz, CDCl<sub>3</sub>, ppm, (PhTe)<sub>2</sub>) δ 251.43. LRMS: *m/z* (70 eV, rel. int.) 260 (M<sup>+</sup>, 13), 258 (M<sup>+</sup>, 13), 256 (7), 255 (3), 254 (2), 215 (3), 186 (8), 72 (5), 57 (73), 55 (100), 45 (44).

### 2-(Butyltellanyl)ethyl-2-methyl-1,3-dioxolane (C)

To a 50 mL two-necked flask equipped with a Dean-Stark reflux apparatus were added telluride **A** (1.28 g, 5 mmol), benzene (10 mL), ethylene glycol (0.44 mL, 8 mmol) and Amberlist® (30 mg). The reaction mixture was refluxed under a nitrogen atmosphere for 5 h. The mixture was filtered and the residue was washed with ethyl acetate. The solvents were removed under reduced pressure. The residue was purified by silica gel column chromatography eluting with a mixture of hexane/ethyl acetate (15:1). Yield: 1.41 g (96%). <sup>1</sup>H NMR: (300 MHz, CDCl<sub>3</sub>, ppm) δ 0.92 (t, *J* = 7.3 Hz, 3H), 1.32 (s, 3H), 1.38 (sext, *J* = 7.3 Hz, 2H), 1.72 (quint, *J* = 7.3 Hz, 2H), 2.10–2.16 (m, 2H), 2.60–2.66 (m, 4H), 3.95 (m, 4H). <sup>13</sup>C NMR: (75 MHz, CDCl<sub>3</sub>, ppm) δ 110.2, 64.7, 42.1, 34.2, 25.0, 23.5, 13.3, 2.7, -5.6. <sup>125</sup>Te NMR: (157 MHz, CDCl<sub>3</sub>, ppm, (PhTe)<sub>2</sub>) δ 263.3. LRMS: *m/z* (70 eV, rel. int.) 302 (M<sup>+</sup>, 3), 99 (4), 87 (100), 55 (15), 43 (58). IR: (film, cm<sup>-1</sup>) 2979, 2958, 2927, 2874, 1457, 1378, 1246, 1212, 1040, 944. Anal. Calc.: C, 40.13; H, 6.68. Found: C, 40.35; H, 6.46%.

### 4-(Butyltellanyl)butan-2-one oxime (D)

To a 50 mL round-bottomed flask were added telluride **A** (1.28 g, 5 mmol), hydroxylamine hydrochloride (0.552 g, 7.5 mmol), molecular sieves 3 Å (1.25 g), and ethanol (20 mL). The mixture was stirred at room temperature, and the progress of the reaction was monitored by TLC. The mixture was filtered and the residue washed with hexane/ethyl acetate 1:1 (20 mL). The organic phase was sequentially washed with NaHCO<sub>3</sub> (sat. sol., 2 × 20 mL), water (2 × 20 mL), and brine (20 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, and filtered and the solvents were removed under reduced pressure. The residue was purified by silica gel column chromatography, eluting with a mixture of hexane/



**Scheme 1.** Chemical structures of functionalized alkyl-organotellurides **A**, **B**, **C**, and **D**.

ethyl acetate (15:1). Yield: 0.88 g (65%).  $^1\text{H}$  NMR: (300 MHz,  $\text{CDCl}_3$ , ppm)  $\delta$  0.91 (t,  $J = 7.2$  Hz, 3H), 1.38 (sext,  $J = 7.2$  Hz, 2H), 1.73 (quint,  $J = 7.2$  Hz, 2H), 1.90 (s, 3H), 2.61–2.76 (m, 6H).  $^{13}\text{C}$  NMR: (75 MHz,  $\text{CDCl}_3$ , ppm)  $\delta$  158.3, 79.4, 38.7, 34.1, 24.9, 13.4, 13.2, 2.8, –3.4. LRMS:  $m/z$  (70 eV, rel. int.) 271 ( $\text{M}^+$ , 11), 256 (6), 216 (44), 199 (7), 185 (5), 171 (4), 157 (2), 145 (6), 130 (8), 86 (44), 70 (16), 57 (100).

### Chemicals

The chemical purity of organotellurides (99.9%) was determined by gas chromatography/high performance liquid chromatography (GC/HPLC).  $\delta$ -Aminolevulinic acid ( $\delta$ -ALA), *p*-dimethylaminobenzaldehyde, and reduced glutathione (GSH) were purchased from Sigma (St. Louis, MO, USA). All other chemicals were of analytical grade and obtained from standard commercial suppliers. Alkyl-organotellurides were dissolved in dimethylsulfoxide (DMSO).

### Animals

Male adult Wistar rats, weighing 200–300 g, from our own breeding colony were used. The animals were kept in a separate animal room, on a 12 h light/dark cycle, at a room temperature of  $22 \pm 2^\circ\text{C}$ , with free access to food (Guabi, RS, Brazil) and water. The animals were used according to the guidelines of the Committee on Care and Use of Experimental Animal Resources, Federal University of Santa Maria, Brazil.

### Tissue preparation

Rats were euthanized and the liver tissue was rapidly dissected, placed on ice, and weighed. The liver tissue was immediately homogenized in cold 50 mM Tris-HCl, pH 7.4 (1/10, w/v). An aliquot of homogenate was used for the protein carbonyl assay. The homogenate was centrifuged at  $2400 \times g$  at  $4^\circ\text{C}$  for 10 min and a low-speed supernatant ( $\text{S}_1$ ) was used for lipid peroxidation and  $\delta$ -ALA-D assays.

Thiol peroxidase-like and thiol oxidase assays were performed in the absence of  $\text{S}_1$ .

### Lipid peroxidation

This assay was carried out to determine the *in vitro* antioxidant effect of functionalized alkyl-organotellurides **A–D**.  $\text{FeCl}_2$ /ethylenediaminetetraacetic acid (EDTA) or sodium nitroprusside (SNP) was used as the classical inductor of lipid peroxidation. Thiobarbituric acid reactive species (TBARS) were used as a measure of lipid peroxidation and determined as described by Ohkawa *et al.*<sup>33</sup>. An aliquot of 200  $\mu\text{L}$  of  $\text{S}_1$  was added to the reaction mixture containing: 100  $\mu\text{M}$   $\text{FeCl}_2$ /50  $\mu\text{M}$  EDTA or 50  $\mu\text{M}$  SNP, 50 mM Tris-HCl, pH 7.4, and alkyl-organotellurides **A–D** at different concentrations (0.1–1  $\mu\text{M}$ ). The range of concentrations used for compound **C** was 0.001–1  $\mu\text{M}$ . Compound **A** was used in a different range of concentrations in the lipid peroxidation assay induced with  $\text{FeCl}_2$ /EDTA (0.1–5  $\mu\text{M}$ ), and the range of concentrations for compound **B** was 0.01–1  $\mu\text{M}$  for lipid peroxidation induced with SNP. Vitamin E and trolox (0.1–50  $\mu\text{M}$ ) were used as positive controls. Afterward the

mixture was pre-incubated for 1 h at  $37^\circ\text{C}$ . The reaction product was determined using 500  $\mu\text{L}$  thiobarbituric acid (0.8%), 200  $\mu\text{L}$  8.1% sodium dodecyl sulfate (SDS), and 500  $\mu\text{L}$  acetic acid buffer (pH 3.4), and incubated at  $95^\circ\text{C}$  for 2 h. The absorbance of samples was determined at 532 nm. Results are reported as nmol MDA (malondialdehyde) per gram of tissue.

### Protein carbonyl content

Oxidatively modified proteins present an enhancement of carbonyl content<sup>34</sup>. This assay was utilized to determine the *in vitro* antioxidant effect of functionalized alkyl-organotellurides **A–D** against protein carbonylation induced by SNP. Carbonyl content was assayed by a method based on the reaction of protein carbonyls with 2,4-dinitrophenylhydrazine (DNPH) forming dinitrophenylhydrazone, a yellow compound<sup>35</sup>. Briefly, homogenate was diluted 1:10 (v/v) and incubated for 2 h with alkyl-organotellurides **A–D** at different concentrations (0.001–1  $\mu\text{M}$ ) and SNP (50  $\mu\text{M}$ ). Vitamin E (0.5  $\mu\text{M}$ –5 mM) and trolox (0.5–100  $\mu\text{M}$ ) were used as positive controls. Then, 1 mL aliquots were mixed with 200  $\mu\text{L}$  of 10 mM DNPH or 200  $\mu\text{L}$  2M HCl. After incubation at room temperature for 1 h in a dark environment, 500  $\mu\text{L}$  of denaturing buffer (150 mM sodium phosphate buffer, pH 6.8, containing 3% SDS), 1.5 mL of heptane (99.5%), and 1.5 mL of ethanol (99.8%) were added sequentially, and mixed with vortex agitation for 40 s and centrifuged for 15 min. Next, the protein isolated from the interface was washed twice with 1 mL of ethyl acetate/ethanol 1:1 (v/v) and suspended in 1 mL of denaturing buffer. Each DNPH sample was read at 370 nm in a spectrophotometer against the corresponding HCl sample (blank), and the total carbonylation was calculated using a molar extinction coefficient of  $22,000 \text{ M}^{-1} \text{ cm}^{-1}$  according to Levine *et al.*<sup>35</sup>. The results are expressed as nmol carbonyl per milligram of protein.

### Thiol peroxidase-like activity

The thiol peroxidase-like activity was examined to determine the possible mechanism by which functionalized alkyl-organotellurides **A–D** demonstrate antioxidant property. Free –SH groups were determined according to Ellman<sup>36</sup>. The catalytic effect of alkyl-organotellurides **A–D** on the reduction of  $\text{H}_2\text{O}_2$  by GSH was assessed using the rate of GSH oxidation. Alkyl-organotellurides **A–D** at different concentrations (0.1–100  $\mu\text{M}$ ) were incubated in the medium containing GSH (1.0 mM), with and without  $\text{H}_2\text{O}_2$  (0.3 mM). At 120 min, aliquots of the reaction mixture (200  $\mu\text{L}$ ) were checked for the amount of GSH. The results are expressed as percentage of control.

### $\delta$ -Aminolevulinic dehydratase activity

$\delta$ -ALA-D has been reported as being sensitive to pro-oxidants and heavy metals<sup>37</sup>. Since this enzyme is very sensitive to organochalcogen compounds,  $\delta$ -ALA-D activity was used as a marker of toxicity.  $\delta$ -ALA-D activity was assayed



by the method of Sassa<sup>38</sup> with some modifications. The principle of this method is based on enzyme incubation with an excess of  $\delta$ -ALA. An aliquot of 200  $\mu$ L of  $S_1$  was pre-incubated for 10 min at 37°C in the presence or absence of functionalized alkyl-organotellurides **A–D** at different concentrations (0.1–100  $\mu$ M). Enzymatic reaction was initiated by adding the substrate,  $\delta$ -ALA, to a final concentration of 2.2 mM in a medium containing 45 mM potassium phosphate buffer, pH 6.8. The incubation was carried out for 1 h at 37°C and stopped by adding trichloroacetic acid solution (10% TCA) with 10 mM  $HgCl_2$ . Porphobilinogen (PBG), formed within a fixed time, was mixed with modified Ehrlich's reagent, and the color developed was measured photometrically (555 nm) against a blank. The results are expressed as nmol PBG per milligram of protein per hour.

#### Thiol oxidase activity

The thiol oxidase activity was examined to determine the pro-oxidant property of functionalized alkyl-organotellurides **A–D**. The rate of thiol oxidation was determined in the presence of 50 mM Tris-HCl, pH 7.5, and alkyl-organotellurides **A–D** at different concentrations (0.1–100  $\mu$ M). Incubation at 37°C was initiated by the addition of 1.0 mM reduced glutathione (GSH). Aliquots of the reaction mixture (200  $\mu$ L) were checked for the amount of –SH groups at 412 nm. The rate of thiol oxidation was evaluated by measuring the disappearance of –SH groups. Free –SH groups were determined according to Ellman<sup>36</sup>. Results are expressed as percentage of control.

#### Correlation between thiol oxidase and $\delta$ -ALA-D

Since the mechanism of  $\delta$ -ALA-D inhibition may be linked to the oxidation of its sulfhydryl groups<sup>39</sup>, the correlation between  $\delta$ -ALA-D activity and thiol oxidase property of functionalized alkyl-organotellurides **A–D** was performed. Through this relationship, we wanted to verify whether the mechanism of  $\delta$ -ALA-D inhibition by organotellurides was related to the oxidation of cysteinyl residues located at the active site of the enzyme.

#### Protein determination

Protein was measured by the Coomassie blue method according to Bradford<sup>40</sup> using bovine serum albumin as standard.

#### Statistical analysis

Data were analyzed using a one-way analysis of variance (ANOVA), followed by Duncan's multiple range test when appropriate. Results were considered significantly different at values of  $p < 0.05$ . The  $IC_{50}$  values (inhibitory concentrations) were calculated by the probit method using computer software SAS/STAT, and are reported as geometric means accompanied by their 95% confidence limits. Maximal inhibition ( $I_{max}$ ) values were calculated at the most effective dose used. Pearson's correlation

coefficient was used for the estimation of correlation between parameters analyzed.

## Results

#### Effect of functionalized alkyl-organotellurides **A–D** on lipid peroxidation induced by $Fe^{2+}$ /EDTA or SNP

Compound **A** at the concentration of 1.00  $\mu$ M was effective in protecting against lipid peroxidation induced by  $Fe^{2+}$ /EDTA in the liver (Figure 1a). Compounds **D** and **B** were effective at concentrations equal to or greater than 0.50 and 0.25  $\mu$ M, respectively (Figure 1d and b). Compound **C** at the concentration of 0.01  $\mu$ M reduced lipid peroxidation induced by  $Fe^{2+}$ /EDTA (Figure 1c). According to the confidence limits of the  $IC_{50}$  values of compounds **A–D**, the  $IC_{50}$  values for functionalized alkyl-organotellurides followed the order: **C**  $\leq$  **B**  $<$  **D**  $<$  **A**. Compounds **A** ( $IC_{50} = 2.90 \mu$ M), **B** ( $IC_{50} = 0.40 \mu$ M), **C** ( $IC_{50} = 0.30 \mu$ M), and **D** ( $IC_{50} = 0.68 \mu$ M) were more effective in reducing lipid peroxidation induced by  $Fe^{2+}$ /EDTA in liver than the classical antioxidants, vitamin E ( $IC_{50} = 2.47 \mu$ M) and trolox ( $IC_{50} = 22.96 \mu$ M) (Table 1). The  $I_{max}$  values for compounds **A**, **B**, and **C** were around 70–80%, while for compound **D** the value was 50% (data not shown).

Compound **A** at the concentration equal to or greater than 0.50  $\mu$ M reduced lipid peroxidation induced by SNP in the liver (Figure 2a). Compounds **D** and **B** were effective at concentrations of 0.25 and 0.10  $\mu$ M, respectively (Figure 2d and b). Compound **C** at the concentration of 0.01  $\mu$ M reduced lipid peroxidation induced by SNP (Figure 2c). According to the confidence limits of the  $IC_{50}$  values of compounds **A–D**, the  $IC_{50}$  values for functionalized alkyl-organotellurides followed the order: **B**  $\leq$  **C**  $\leq$  **D**  $<$  **A**. Compounds **A** ( $IC_{50} = 1.21 \mu$ M), **B** ( $IC_{50} = 0.21 \mu$ M), **C** ( $IC_{50} = 0.33 \mu$ M), and **D** ( $IC_{50} = 0.43 \mu$ M) were more effective in reducing lipid peroxidation induced by SNP in liver than vitamin E ( $IC_{50} = 2.97 \mu$ M) and trolox ( $IC_{50} = 27.20 \mu$ M) (Table 1). The  $I_{max}$  values for compounds **A–D** were around 70–80% (data not shown).

#### Effect of functionalized alkyl-organotellurides **A–D** on protein carbonyl content


Compounds **A** and **C** were effective in protecting against protein carbonylation induced by SNP in liver at the concentrations of 0.50 and 0.10  $\mu$ M, respectively (Figure 3a and c). Compounds **B** and **D**, at the concentration of 0.25  $\mu$ M, reduced protein carbonyl content (Figure 3b and d). According to the confidence limits of the  $IC_{50}$  values of compounds **A–D**, the  $IC_{50}$  values for functionalized alkyl-organotellurides followed the order: **A** = **D**  $\geq$  **B** = **C**. Compounds **A** ( $IC_{50} = 0.81 \mu$ M), **B** ( $IC_{50} = 0.69 \mu$ M), **C** ( $IC_{50} = 0.71 \mu$ M), and **D** ( $IC_{50} = 0.83 \mu$ M) were more effective in reducing protein carbonyl formation in the liver than vitamin E ( $IC_{50} = 326.34 \mu$ M) and trolox ( $IC_{50} = 65.95 \mu$ M) (Table 1). The  $I_{max}$  values for compounds **A–D** were around 50% (data not shown).

**Effect of functionalized alkyl-organotellurides A-D on thiol peroxidase-like activity**

Functionalized alkyl-organotellurides **A-D** at different concentrations (0.1–100  $\mu\text{M}$ ) did not present thiol peroxidase-like activity (data not shown).

**Effect of functionalized alkyl-organotellurides A-D on  $\delta$ -ALA-D activity**

Functionalized alkyl-organotellurides **A-D**, at the concentration of 100  $\mu\text{M}$ , reduced significantly the  $\delta$ -ALA-D activity in rat liver homogenates (data not shown).




**Figure 1.** Effect of **A** (0.1–5  $\mu\text{M}$ ) (a), **B** (0.1–1  $\mu\text{M}$ ) (b), **C** (0.001–1  $\mu\text{M}$ ) (c), and **D** (0.1–1  $\mu\text{M}$ ) (d) on lipid peroxidation induced with  $\text{Fe}^{2+}/\text{EDTA}$ . Data are reported as mean  $\pm$  SD for four independent experiments, each one in duplicate. \*, Denotes  $p < 0.05$  as compared to the sample with  $\text{Fe}^{2+}/\text{EDTA}$  (one-way ANOVA/Duncan). #, Denotes  $p < 0.05$  as compared to control values (without  $\text{Fe}^{2+}/\text{EDTA}$  and compounds) (one-way ANOVA/Duncan).

**Table 1.** Calculated  $\text{IC}_{50}$  values ( $\mu\text{M}$ ) for lipid peroxidation and protein carbonyl content of functionalized alkyl-organotellurides in rat liver.

Compound	Lipid peroxidation		Protein carbonyl
	$\text{Fe}^{2+}/\text{EDTA}$	SNP	SNP
Vitamin E	2.47 (0.54–4.17)	2.97 (0.49–4.97)	326.34 (288.48–369.53)
Trolox	22.96 (19.92–26.45)	27.20 (18.76–39.55)	65.95 (60.67–71.70)
<b>A</b>	2.90 (2.60–3.20)	1.21 (1.05–1.40)	0.81 (0.72–0.92)
<b>B</b>	0.40 (0.37–0.44)	0.21 (0.19–0.22)	0.69 (0.47–1.02)
<b>C</b>	0.30 (0.24–0.54)	0.33 (0.12–0.89)	0.71 (0.58–0.89)
<b>D</b>	0.68 (0.65–0.71)	0.43 (0.36–0.51)	0.83 (0.70–0.98)

Note. 95% Confidence limit ( $\mu\text{M}$ ) presented in parentheses.



**Figure 2.** Effect of **A** (0.1–1  $\mu\text{M}$ ) (a), **B** (0.01–1  $\mu\text{M}$ ) (b), **C** (0.001–1  $\mu\text{M}$ ) (c), and **D** (0.1–1  $\mu\text{M}$ ) (d) on lipid peroxidation induced with SNP. Data are reported as mean  $\pm$  SD for four independent experiments, each one in duplicate. \*, Denotes  $p < 0.05$  as compared to sample with SNP (one-way ANOVA/Duncan). #, Denotes  $p < 0.05$  as compared to control values (without SNP and compounds) (one-way ANOVA/Duncan).

#### Effect of functionalized alkyl-organotellurides A–D on thiol oxidase activity

Functionalized alkyl-organotellurides **A–D**, at the concentration of 100  $\mu\text{M}$ , increased the thiol oxidase activity (data not shown).


#### Correlation of thiol oxidase and $\delta$ -ALA-D activities

The results demonstrated a significant inverse correlation ( $r = 0.987$ ,  $p < 0.005$ ) between thiol oxidase and  $\delta$ -ALA-D activity for compound **A** (Figure 4). The  $r$  values for alkyl-organotellurides **B**, **C**, and **D**, which also presented significant inverse correlation between these parameters, were  $r = 0.916$ ,  $r = 0.970$ , and  $r = 0.972$ , respectively (data not shown).


## Discussion

The results of the present study showed that functionalized alkyl-organotellurides **A–D**, at low  $\mu\text{M}$  range, displayed an *in vitro* antioxidant activity in rat liver homogenates. Alkyl-organotellurides were able to prevent lipid peroxidation

induced by  $\text{Fe}^{2+}$ /EDTA and SNP in rat liver homogenates. It is known that iron complexes, such as  $\text{Fe}^{2+}$ /EDTA, stimulate lipid peroxidation in cells<sup>41</sup>. SNP has been suggested to cause cytotoxicity through the release of cyanide and/or nitric oxide<sup>42</sup>. In the present study, we demonstrated that low concentrations of functionalized alkyl-organotellurides were required to reduce lipid peroxidation induced by  $\text{Fe}^{2+}$ /EDTA or SNP in rat liver homogenates. The  $\text{IC}_{50}$  values for functionalized alkyl-organotellurides **A–D** of lipid peroxidation induced by  $\text{Fe}^{2+}$ /EDTA or SNP were close. In this way, we can infer that the antioxidant property of the studied alkyl-organotellurides did not depend on the inductor used. Moreover, compounds **B**, **C**, and **D** demonstrated higher efficiency in reducing lipid peroxidation induced by both inductors than compound **A**, as can be seen by the  $\text{IC}_{50}$  values. The  $\text{IC}_{50}$  values for functionalized alkyl-organotellurides followed the order **C**  $\leq$  **B**  $<$  **D**  $<$  **A** for lipid peroxidation induced by  $\text{Fe}^{2+}$ /EDTA or SNP. However, it is difficult to clearly explain why compound **A** displayed the lowest efficiency against lipid peroxidation. Based on chemical structures of alkyl-organotellurides **A–D**



**Figure 3.** Effect of **A** (0.1–1  $\mu\text{M}$ ) (a), **B** (0.1–1  $\mu\text{M}$ ) (b), **C** (0.001–1  $\mu\text{M}$ ) (c), and **D** (0.1–1  $\mu\text{M}$ ) (d) on protein carbonyl content. Data are reported as mean  $\pm$  SD for four independent experiments, each one in duplicate. \*, Denotes  $p < 0.05$  as compared to the sample with SNP (one-way ANOVA/Duncan). #, Denotes  $p < 0.05$  as compared to control values (without SNP and compounds) (one-way ANOVA/Duncan).



**Figure 4.** Correlation between thiol oxidase property and  $\delta$ -ALA-D activity of alkyl-organotelluride **A**.

there is not an obvious explanation for the lowest antioxidant activity of compound **A**. Conversely, the literature data indicate that the chemical structure of organochalcogens has an important role in establishing their application as biological antioxidants<sup>43,44</sup>.

Different from the results for lipid peroxidation, alkyl-organotellurides **A**, **B**, **C**, and **D** demonstrated similar effects on protein carbonyl content, suggesting that structural differences did not affect the antioxidant effect on proteins. Furthermore, carbonyl protein is the most used biomarker for protein oxidative damage, and it reflects the cellular damage induced by multiple forms of ROS<sup>45</sup>. Taken together, the results of the current study demonstrated that functionalized alkyl-organotellurides, at low concentrations, displayed an antioxidant effect on lipid and protein oxidation.

Some authors have demonstrated that tellurium compounds are effective antioxidant agents, even more active than their selenium and sulfur analogs<sup>1,46</sup>. The antioxidant activity of diorganoyl tellurides and diselenides<sup>8</sup> has been



attributed, at least in part, to the glutathione peroxidase-like activity of these compounds<sup>10,43</sup>. In contrast, functionalized alkyl-organotellurides **A–D** did not display glutathione peroxidase-like activity, discarding that the antioxidant activity of these compounds could be related to this mechanism. However, other mechanisms could be involved in the antioxidant property of the studied functionalized alkyl-organotellurides<sup>11</sup>. In fact, more studies are necessary to elucidate the exact mechanism involved in the antioxidant effect caused by alkyl-organotellurides **A–D**.

Although functionalized alkyl-organotellurides presented an antioxidant effect, the literature data report that they induce toxicity<sup>7</sup>. In this way, previous works have indicated that alkyl-organotellurides inhibit cysteinyl-containing enzymes, such as Na<sup>+</sup>/K<sup>+</sup> ATPase<sup>47</sup>,  $\delta$ -ALA-D<sup>37,39</sup>, and squalene monooxygenase<sup>48</sup>. Thus, the toxicity of organotellurium compounds can be related to interaction with the thiol groups of important biomolecules<sup>8</sup>, the replacement of selenium in selenoproteins, such as thioredoxin<sup>1</sup>, and the capacity of tellurium compounds to induce the formation of ROS<sup>49</sup>.

Surprisingly, the present study demonstrated that functionalized alkyl-organotellurides **a–d** were good antioxidants, and affected  $\delta$ -ALA-D activity exclusively at high concentrations. In fact,  $\delta$ -ALA-D is a group of sulfhydryl-containing enzymes<sup>50</sup>, being extremely sensitive to the presence of pro-oxidant elements, and numerous compounds that oxidize sulfhydryl groups, such as selenium and tellurium compounds<sup>39,51,52</sup>, modify its activity. Particularly, organotellurium compounds may interact with thiol groups of this enzyme, oxidizing sulfhydryl groups of the active site, changing the configuration of the enzyme and impairing its activity. A previous study from our research group reported that organochalcogen compounds, at low concentrations, reduced  $\delta$ -ALA-D activity<sup>37,39,51</sup>. Conversely, we demonstrated here that functionalized alkyl-organotellurides **a–d** did not inhibit  $\delta$ -ALA-D activity at concentrations for which they displayed antioxidant activity. In fact, these compounds inhibited enzyme activity at the concentration of 100  $\mu$ M, a concentration 100 times higher than those at which alkyl-organotellurides presented the antioxidant effect. This result highlights this class of compounds as good antioxidant agents with low potential toxicity, demonstrated in rat liver homogenates *in vitro*. The thiol oxidase activity is an estimation of the cytotoxicity of molecules<sup>8</sup>. Organotellurides **A–D** showed thiol oxidase activity only at the concentration of 100  $\mu$ M, indicating an advantage regarding the possible toxicological properties of these compounds. Recently, our research group reported that the mechanism involved in  $\delta$ -ALA-D inhibition caused by telluroacetylenes *in vitro* is related to their ability to oxidize sulfhydryl groups from the enzyme structure<sup>52</sup>. In conformity, our results demonstrated an inverse correlation between thiol oxidase and  $\delta$ -ALA-D activities, meaning that  $\delta$ -ALA-D activity decreases as the compound's ability to oxidize sulfhydryl groups increases. Therefore, at the concentration for which compounds have the thiol oxidase property, they also inhibit  $\delta$ -ALA-D activity, suggesting that oxidation of

enzyme sulfhydryl groups could be involved in the inhibition caused by functionalized alkyl-organotellurides. Thus, similar to other organochalcogens<sup>7,39,51–53</sup>, functionalized alkyl-organotellurides oxidize thiol groups to disulfides<sup>8</sup>, clearly demonstrated by the thiol oxidase activity and the results obtained with  $\delta$ -ALA-D activity in rat liver homogenates *in vitro*.

In conclusion, the results of the present investigation support the *in vitro* antioxidant effect of functionalized alkyl-organotellurides **A–D** in rat liver homogenates. Of particular importance, functionalized alkyl-organotellurides **A–D** inhibited  $\delta$ -ALA-D activity at a concentration 100 times higher than those for which they displayed the antioxidant effect. Additional studies are necessary to investigate the possible antioxidant effect of these compounds in an *in vivo* model of oxidative damage.

## Declaration of interest

The financial support by UFSM (FIPE), FAPERGS, CAPES, FAPESP, and CNPq is gratefully acknowledged.

## References

- Engman L, Kanda T, Gallegos A. Water-soluble organotellurium compounds inhibit thioredoxin reductase and the growth of human cancer cells. *Anticancer Drug Des* 2000;15:323–30.
- Nogueira CW, Quinhones EB, Jung EAC, Zeni G, Rocha JBT. Anti-inflammatory and antinociceptive activity of diphenyl diselenide. *Inflamm Res* 2003;63:52–6.
- Cunha LOR, Urano ME, Chagas JR, Almeida PC, Bincoletto C, Tersariol ILS, et al. Tellurium-based cysteine protease inhibitors: evaluation of novel organotellurium (IV) compounds as inhibitors of human cathepsin B. *Bioorg Med Chem Lett* 2005;15:755–60.
- Savegnago L, Jesse CR, Pinto LG, Rocha JBT, Barancelli DA, Nogueira CW. Diphenyl diselenide exerts antidepressant-like and anxiolytic-like effects in mice: involvement of l-arginine-nitric oxide-soluble guanylate cyclase pathway in its antidepressant-like action. *Pharmacol Biochem Behav* 2008;88:418–26.
- Wöhler F. *Ann Chem* 1840;35:111.
- Rooseboom M, Vermeulen NPE, Durgut F. Comparative study on the bioactivation mechanisms and cytotoxicity of Te-phenyl-L-tellurocysteine, Se-phenyl-L-selenocysteine, and S-phenyl-L-cysteine. *Chem Res Toxicol* 2002;15:1610–18.
- Meotti FC, Borges VC, Zeni JBT, Nogueira CW. Potential renal and hepatic toxicity of diphenyl diselenide, diphenyl ditelluride and Ebselen for rats and mice. *Toxicol Lett* 2003;143:9–16.
- Nogueira CW, Zeni G, Rocha JBT. Organoselenium and organotellurium compounds: toxicology and pharmacology. *Chem Rev* 2004;104:6255–85.
- Engman L, Al-Maharik N, McNaughton M. Thioredoxin reductase and cancer cell growth inhibition by organotellurium compounds that could be selectively incorporated into tumor cells. *Bioorg Med Chem* 2003;11:5091–100.
- Ren X, Xue Y, Zhang K, Liu J, Luo G, Zheng J, et al. A novel dicyclodextrinyl ditelluride compound with antioxidant activity. *FEBS Lett* 2001;507:377–80.
- Kanski J, Drake J, Aksenova M. Antioxidant activity of the organotellurium compound 3-[4-(N,N-dimethylamino)benzenetellurenyl]propanesulfonic acid against oxidative stress in synaptosomal membrane systems and neuronal cultures. *Brain Res* 2001;911:12–21.
- Ávila DS, Beque MC, Folmer V, Braga AL, Zeni G, Nogueira CW, et al. Diethyl 2-phenyl-2-tellurophenyl vinylphosphonate: an organotellurium compound with low toxicity. *Toxicology* 2006;224:100–7.
- Savegnago L, Borges VC, Alves D, Jessé CR, Rocha JBT, Nogueira CW. Evaluation of antioxidant activity and potential toxicity of 1-buthyltelurenyl-2-methylthioheptene. *Life Sci* 2006;79:1546–52.

14. Ávila DS, Gubert P, Corte CLD, Alves D, Nogueira CW, Rocha JBT, et al. A biochemical and toxicological study with diethyl 2-phenyl-2-telluorophenyl vinylphosphonate in a sub-chronic intraperitoneal treatment in mice. *Life Sci* 2007;80:1865–72.
15. Arteel GE, Sies H. The biochemistry of selenium and the glutathione system. *Environ Toxicol Pharmacol* 2001;10:153–8.
16. Lecanu L, Greeson J, Papadopoulos V. Beta-amyloid and oxidative stress jointly induce neuronal death, amyloid deposits, gliosis, and memory impairment in the rat brain. *Pharmacology* 2006;76:19–33.
17. Khan TH, Sultana S. Antioxidant and hepatoprotective potential of Aegle marmelos Correa. against CCL4-induced oxidative stress and early tumor events. *J Enzyme Inhib Med Chem* 2009;24:320–7.
18. Ercan S, Cakmak A, Kosciak M. Total oxidant and antioxidant levels and oxidative stress index levels in children with congenital heart disease. *Acta Paediatr* 2008;97:86–7.
19. Nabben M, Hoeks J. Mitochondrial uncoupling protein 3 and its role in cardiac- and skeletal muscle metabolism. *Phys Behav* 2008;94:259–69.
20. Augustyniak K, Zavadnik I, Palecz D, Szosland K, Bryszewska M. The effect of oxidizing agent and diabetes mellitus on the human red blood cell membrane potential. *Clin Biochem* 1996;29:283–6.
21. Gey KF. The relationship of antioxidant status and the risk of cancer and cardiovascular disease: a critical evaluation of observational data. In: Nohl H, Esterbauer H, Rice-Evans C, eds. *Free Radicals in the Environment, Medicine and Toxicology*. London: Richelieu Press, 1994:181–219.
22. Wang BB, Wang J, Zhou SR. The association of mitochondrial aldehyde dehydrogenase gene (ALDH2) polymorphism with susceptibility to late-onset Alzheimer's disease in Chinese. *J Neurol Sci* 2008;268:172–5.
23. Ayhan-Kilicgil G, Kus C, Coban T, Can-Eke B, Iscan M. Antioxidant and neuroprotective effects of synthesized sintonin derivatives. *J Enzyme Inhib Med Chem* 2004;19:129–35.
24. Yang LX, Zhang LJ, Huang KX, Kun Li X, Hu LH, Wang XY, et al. Antioxidant and neuroprotective effects of synthesized sintonin derivatives. *J Enzyme Inhib Med Chem* 2009;24:425–31.
25. Dos Santos AA, Comassetto JV.  $\beta$ -Butyltellanyl carbonyl compounds: a useful source of masked metal homoenolates. *J Braz Chem Soc* 2005;16:511–13.
26. Comassetto JV, Gariani RA, Princival JL, Dos Santos AA, Zinn FK. Hydrochalcogenation of activated olefines. Synthesis of functionalized dialkylchalcogenides. *Organomet Chem* 2008;693:2929–36.
27. Princival JL, Barros SMG, Comassetto JV, Dos Santos AA.  $\beta$ -Butyltelluro-2-butanol: a route to reactive 1,4-dianion intermediates. *Tetrahedron Lett* 2005;46:4423–5.
28. Dos Santos AA, Ferrarini RS, Princival JL, Comassetto JV. Organotellurides as a source of organometallics: application in the synthesis of (+/-)-frontalin. *Tetrahedron Lett* 2006;47:8933–5.
29. Comassetto JV, Dos Santos AA. Organotellurides as precursors of reactive organometallics. *Phosphorus Sulfur Silicon Relat Elem* 2008;183:939–47.
30. Ferrarini RS, Princival JL, Comassetto JV, Dos Santos AA. A concise enantioselective synthesis of (+)-endo-brevicomine accomplished by a tellurium/metal exchange reaction. *Braz Chem Soc* 2008;19:811–12.
31. Wendler EP, Dos Santos AA. The use of butyl organotellurides in the synthesis of natural bioactive compounds. *Synlett* 2009:1034–40.
32. Bigdeli MA, Alvi Nikje MM, Jafari S, Heravi MM. Regioselective synthesis of syn-oximes using 3 Å molecular sieves in a solventless system. *J Chem Res Synop* 2002;1:20–1.
33. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351–8.
34. Yan LJ, Traber MG, Packer L. Spectrophotometric method for determination of carbonyls in oxidatively modified apolipoprotein B of human low-density lipoproteins. *Anal Biochem* 1995;228:349–51.
35. Levine RL, Garland D, Oliver CN. Determination of carbonyl content in oxidatively modified proteins. *Methods Enzymol* 1990;186:464–78.
36. Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys* 1959;82:70–7.
37. Nogueira CW, Borges VC, Zeni G, Rocha JBT. Organochalcogens effects on  $\delta$ -aminolevulinic acid dehydratase activity from human erythrocytic cells in vitro. *Toxicology* 2003;191:169–78.
38. Sassa S. Delta aminolevulinic acid dehydratase assay. *Enzyme* 1982;28:133–45.
39. Barbosa NBV, Rocha JBT, Zeni G, Emanuelli T, Beque MC, Braga AL. Effect of organic selenium on  $\delta$ -aminolevulinic acid dehydratase from liver, kidney, and brain of adult rats. *Toxicol Appl Pharmacol* 1998;149:243–53.
40. Bradford MM. Rapid and sensitive method for quantitation of microgram quantities of protein utilizing principle of protein-dye binding. *Anal Biochem* 1976;72:248–54.
41. Gogvadze V, Walter PB, Ames BN. The role of  $\text{Fe}^{2+}$ -induced lipid peroxidation in the initiation of the mitochondrial permeability transition. *Arch Biochem Biophys* 2003;414:255–60.
42. Bates JN, Baker MT, Guerra R, Harrison DG. Nitric oxide generation from nitroprusside by vascular tissue: *Biochem Pharmacol* 1990;42:S157–65.
43. Meotti FC, Stangherlin EC, Zeni G, Nogueira CW, Rocha JBT. Protective role of aryl and alkyl diselenides on lipid peroxidation. *Environ Res* 2004;94:276–82.
44. Prigol M, Wilhelm EA, Schneider CC, Nogueira CW. Protective effect of unsymmetrical dichalcogenide, a novel antioxidant agent, in vitro and an in vivo model of brain oxidative damage. *Chem Biol Interact* 2008;176:129–36.
45. Dalle-Donne I, Aldini G, Carini M, Colombo R, Rossi R, Milzani A. Protein carbonylation, cellular dysfunction, and disease progression. *J Cell Mol Med* 2006;10:389–406.
46. Wieslander E, Engman L, Svensjo E, Erlansson M, Johansson U, Linden M, et al. Antioxidative properties of organotellurium compounds in cell system. *Biochem Pharmacol* 1998;55:573–84.
47. Borges VC, Rocha JBT, Nogueira CW. Effect of diphenyl diselenide, diphenyl ditelluride and ebselen on cerebral  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in rats. *Toxicology* 2005;215:191–7.
48. Laden BP, Porter TD. Inhibition of human squalene monooxygenase by tellurium compounds: evidence of interaction with vicinal sulfhydryls. *J Lipid Res* 2001;42:235–40.
49. Chen F, Vallyathan V, Castranova V, Shi X. Cell apoptosis induced carcinogenic metals. *Mol Cell Biochem* 2001;221:183–8.
50. Gibson KD, Neuberger A, Scott JJ. The purification and properties of delta-aminolevulinic acid dehydratase. *Biochem J* 1955;61:618–28.
51. Luchese C, Zeni G, Rocha JBT, Nogueira CW, Santos FW. Cadmium inhibits  $\delta$ -aminolevulinic acid dehydratase from rat lung in vitro: interaction with chelating and antioxidant agents. *Chem Biol Interact* 2007;165:127–37.
52. Souza ACG, Luchese C, Neto JSS, Nogueira CW. Antioxidant effect of a novel class of telluroacetylene compounds: studies in vitro and in vivo. *Life Sci* 2009;84: 351–7.
53. Maciel EN, Bolzan RC, Braga AL, Rocha JBT. Diphenyl diselenide and diphenyl ditelluride differentially affect delta-aminolevulinic acid dehydratase from liver, kidney, and brain of mice. *J Biochem Mol Toxicol* 2000;310: 319–14.