



Synthesis, characterization, cytotoxicity, and DNA binding of some new platinum(II) and platinum(IV) complexes with benzimidazole ligands

Semra Utku, Fatma Gumus, Seda Tezcan, Mehmet Sami Serin & Aykut Ozkul

To cite this article: Semra Utku, Fatma Gumus, Seda Tezcan, Mehmet Sami Serin & Aykut Ozkul (2010) Synthesis, characterization, cytotoxicity, and DNA binding of some new platinum(II) and platinum(IV) complexes with benzimidazole ligands, Journal of Enzyme Inhibition and Medicinal Chemistry, 25:4, 502-508, DOI: [10.3109/14756360903282858](https://doi.org/10.3109/14756360903282858)

To link to this article: <https://doi.org/10.3109/14756360903282858>



Published online: 16 Mar 2010.



Submit your article to this journal [↗](#)



Article views: 1041



View related articles [↗](#)



Citing articles: 5 View citing articles [↗](#)

ORIGINAL ARTICLE

Synthesis, characterization, cytotoxicity, and DNA binding of some new platinum(II) and platinum(IV) complexes with benzimidazole ligands

Semra Utku¹, Fatma Gumus², Seda Tezcan¹, Mehmet Sami Serin⁴, and Aykut Ozkul⁵

¹Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Mersin, Mersin, Turkey, ²Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Gazi, Ankara, Turkey, ³Department of Medical Microbiology, Faculty of Medicine, University of Mersin, Mersin, Turkey, ⁴Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Mersin, Mersin, Turkey, and ⁵Department of Virology, Faculty of Veterinary Medicine, University of Ankara, Ankara, Turkey.

Abstract

In this study, two Pt(II) and three Pt(IV) complexes with the structures of [PtL₂Cl₂] (1), [PtL₂I₂] (2), [PtL₂Cl₂(OH)₂] (3), [PtL₂Cl₂(OCOCH₃)₂] (4), and [PtL₂Cl₄] (5) (L = benzimidazole as carrier ligand) were synthesized and evaluated for their *in vitro* antiproliferative activities against the human MCF-7, HeLa, and HEP-2 cancer cell lines. The influence of compounds 1–5 on the tertiary structure of DNA was determined by their ability to modify the electrophoretic mobility of the form I and II bands of pBR322 plasmid DNA. The inhibition of BamH1 restriction enzyme activity of compounds 1–5 was also determined. In general, it was found that compounds 1–5 were less active than cisplatin and carboplatin against MCF-7 and HeLa cell lines (except for 1, which was found to be more active than carboplatin against the MCF-7 cell line). Compounds 1 and 3 were found to be significantly more active than cisplatin and carboplatin against the HEP-2 cell line.

Keywords: Benzimidazole; cytotoxic activity; DNA binding; gel electrophoresis; platinum complexes

Introduction

Platinum-based drugs such as cisplatin, carboplatin, and oxaliplatin are widely used against various solid tumors including genitourinary, colorectal, and non-small cell lung cancers¹. In general, the platinum-based drugs suffer from two main disadvantages: chemoresistance to the drugs can occur and, second, they are non-selective toward cancer cells, which lead to severe toxic side effects, primarily kidney toxicity and neurotoxicity².

Although there is some evidence to suggest that other biological targets may be important in the mechanism, it is generally accepted that DNA is the primary biological target of the platinum drugs^{3,4}. Numerous studies show that, in DNA, cisplatin forms ~90% intrastrand cross-links (CLs) between neighboring purine bases (1,2-GG or 1,2-AG intrastrand CLs), and remaining lesions are intrastrand CLs between purine bases separated by a third base, interstrand CLs, and monofunctional adducts⁵.

Cellular resistance to cisplatin is largely attributed to up-regulation of DNA repair and damage tolerance pathways, lowered intracellular accumulation, and inactivation by thiol-containing reductants such as glutathione and metallothionein, and alterations in proteins involved in apoptosis⁶. One strategy to overcome cisplatin resistance is to design new platinum complexes that specifically deal with some or even all of the resistance mechanisms⁷.

In addition to square-planar Pt(II) complexes, an attempt has also been made recently to synthesize octahedral Pt(IV) complexes, since some of these complexes are toxic to tumors that are resistant to cisplatin⁸. The antitumor activity of Pt(IV) complexes has been suggested to require *in vivo* reduction to the kinetically more labile, and therefore reactive, Pt(II) derivatives⁹. On the other hand, there are a few papers reporting that Pt(IV) complexes can bind to DNA and RNA fragments without being reduced^{10,11}.

Address for Correspondence: Fatma Gumus, Pharmaceutical Chemistry, Faculty of Pharmacy, Gazi University Ankara, Turkey, Tel. and Fax: +90 0312 2120236
E-mail: fgumus@gazi.edu.tr or gumus53@gmail.com

(Received 07 May 2009; revised 19 August 2009; accepted 25 August 2009)

ISSN 1475-6366 print/ISSN 1475-6374 online © 2010 Informa UK Ltd
DOI: 10.3109/14756360903282858

<http://www.informahealthcare.com/enz>

In the literature it is reported that the reduction potentials of diam(m)ine Pt(IV) complexes are dependent on the nature of the axial and equatorial ligands, but axial ligands generally exert the stronger influence^{12,13}. The axial ligands of Pt(IV) complexes offer a unique ability in drug design to modify the pharmacokinetic parameters of a prodrug, including the rate of reduction, the lipophilicity, molecular targeting, and microenvironmental targeting, in order to reduce side effects, increase activity, and target tumor sites, or the axial ligands themselves can confer additional cytotoxicity upon release^{12,14}.

Although no platinum(IV) complex is an approved drug yet, some Pt(IV) complexes have shown sufficient promise to enter clinical trials: iproplatin, [*cis,trans,cis*-[PtCl₂(OH)₂(isopropylamine)₂], tetraplatin, [PtCl₄(cyclohexane-1,2-diamine)], and satraplatin, [*cis,trans*-[PtCl₂(OAc)₂(NH₃)₂ cyclohexylamine] have been tested in clinical trials¹⁵. At present, the most interesting candidate is satraplatin, (formerly JM216), which is in advanced phase III clinical trials showing activity in hormone-refractory prostate cancer in combination with prednisone as second-line chemotherapy¹⁶.

In previous studies, taking into consideration the fact that variations in the chemical structure of the ammine groups of the cisplatin can have significant effects on the cytotoxic activity and toxicity of platinum complexes, we synthesized some Pt(II) and also Pt(IV) complexes with 2-substituted benzimidazole ligands^{17–24}. It was determined that some of these platinum complexes have *in vitro* cytotoxic activities on RD¹⁹, HeLa^{21,22,24}, HEP-2²³, and MCF-7^{20–22,24} cell lines.

In the present study, with the aim of determining the effect of axial and equatorial ligand variation on the cytotoxic activities of the Pt(II) and Pt(IV) complexes, we synthesized some Pt(II) and Pt(IV) complexes with the structures of [PtL₂Cl₂] (**1**), [PtL₂I₂] (**2**), [PtL₂Cl₂(OH)₂] (**3**), [PtL₂Cl₂(OCOCH₃)₂] (**4**), and [PtL₂Cl₄] (**5**) (L = benzimidazole as carrier ligand) and tested for their preliminary *in vitro* antiproliferative activities against the human MCF-7 breast, HeLa cervix, and HEP-2 larynx carcinoma cell lines. The plasmid DNA interaction and the inhibition of BamHI restriction enzyme activity of the compounds **1–5** were also studied.

Experimental

Materials and instruments

All chemicals and solvents used in the synthesis were purchased from Merck or Aldrich Chemical Co. Cisplatin and carboplatin used in the cytotoxicity test were purchased from Sigma Co. Infrared (IR) spectra were recorded on KBr pellets and Nujol mulls on a Mattson 1000 FTIR spectrometer in the range of 4000–200 cm⁻¹. For the region 400–200 cm⁻¹ the samples were prepared as Nujol mulls on CsI windows. Elemental analyses were performed with a Leco 932 CHNS analyser, and ¹H nuclear magnetic resonance (NMR) spectra were recorded in dimethylsulfoxide (DMSO)-d₆ (Merck) on a Varian Mercury 400 MHz FT NMR spectrometer at the Central Laboratory of the Faculty of Pharmacy, Ankara

University (Ankara, Turkey). Thin-layer chromatography (TLC) was performed on aluminum plates pre-coated with Merck silica gel 60 F₂₅₄. Plates were visualized by ultraviolet (UV) light, Dragendorff reagent, or iodine vapor.

Synthesis of ligand, Pt(II) and Pt(IV) complexes

Benzimidazole (L)

L, used as carrier ligand in the structure of **1–5**, was prepared according to the published procedure²⁵. Mp 170°C, (169–170°C²⁶); yield 59%. ¹H NMR (400 MHz, DMSO-d₆) δ 12.40 (broad s, 1H, N-H, exchangeable with D₂O), 8.22 (s, 1H, ArH), 7.58–7.60 (dd, *J* = 9.2 and 2.8 Hz, 2H, ArH), 7.17–7.20 (dd, *J* = 9.2 and 2.8 Hz, 2H, ArH). IR (KBr, cm⁻¹) ν 3300–2300 (N-H) cm⁻¹.

[Dichloro-bis(benzimidazole)platinum(II)] [PtL₂Cl₂] (**1**)

To a stirred aqueous solution of K₂PtCl₄ (0.830 g, 2.00 mmol) was added a solution of benzimidazole (**L**) (0.472 g, 4.00 mmol) in an ethanol–water mixture (8:12 mL) dropwise over 2 h at room temperature. The pH was adjusted to ~7 and kept constant with the addition of 0.1 M NaHCO₃. The reaction mixture, protected from light, was heated at 60°C for 2 days. After that time the mixture was cooled to room temperature. The resulting crude precipitate was filtered off and purified by repeated washing with small portions of water, ethanol, and diethylether, and dried *in vacuo* to yield 0.755 g (75.16%). ¹H NMR (400 MHz, DMSO-d₆) δ 13.44 (s, 2H, 2 × N-H, exchangeable with D₂O), 8.84 (s, 2H, 2 × ArH), 7.80–7.79 (d, *J* = 7.2 Hz, 2H, ArH), 7.50–7.48 (d, *J* = 7.2 Hz, 2H, ArH), 7.25–7.18 (m, 4H, 2 × ArH). IR (KBr, cm⁻¹) ν 3287 (N-H), 326 (Pt-Cl) cm⁻¹. Anal. Calcd. for C₁₄H₁₂Cl₂N₄Pt: C, 33.48; H, 2.41; N, 11.15. Found: C, 33.46; H, 2.64; N, 11.09%.

[Diiodo-bis(benzimidazole)platinum(II)] [PtL₂I₂] (**2**)

K₂PtCl₄ (0.622 g, 1.50 mmol) and KI (0.996 g, 6.00 mmol) were dissolved in water (15 mL) and stirred at 60°C for 45 min. Then a solution of benzimidazole (**L**) (0.331 g, 2.80 mmol) in an ethanol–water mixture (6:10 mL) was added dropwise over 2 h at room temperature to the resulting K₂PtI₄. The reaction mixture, protected from light, was heated at 60°C for 2 days. The resulting crude yellowish precipitate was filtered off and purified by repeated washing with small portions of water, ethanol, and diethylether, and dried *in vacuo* to yield 0.410 g (39.92%). ¹H NMR (400 MHz, DMSO-d₆) δ 13.38 (broad s, 2H, 2 × N-H, exchangeable with D₂O), 8.98 (s, 2H, 2 × ArH), 8.05–8.03 (m, 2H, 2 × ArH), 7.67–7.64 (m, 2H, 2 × ArH), 7.38–7.20 (m, 4H, 2 × ArH). IR (KBr, cm⁻¹) ν 3310 (N-H) cm⁻¹. Anal. Calcd. for C₁₄H₁₂I₂N₄Pt: C, 24.54; H, 1.77; N, 8.18. Found: C, 24.13; H, 2.00; N, 8.41%.

[Dichloro-dihydroxy-bis(benzimidazole)platinum(IV)] [PtL₂Cl₂(OH)₂] (**3**)

To a suspension of **1** (0.115 g, 0.23 mmol) in water (15 mL) was added an aqueous solution of 30% H₂O₂ (1 mL), and the reaction mixture, protected from light, was stirred at 60°C for 5 days. The resulting crude yellow precipitate was filtered off and purified by repeated washing with small portions of

water, ethanol, acetone, and diethylether, and dried *in vacuo* to yield 0.049 g (39.72%). ^1H NMR (400 MHz, DMSO- d_6) δ 13.50 (s, 2H, 2 \times N-H, exchangeable with D_2O), 8.92 (s, 1H, 2 \times ArH), 8.70 (s, 1H, 2 \times ArH), 7.90–7.84 (m, 1H, 2 \times ArH), 7.68–7.58 (m, 2H, 2 \times ArH), 7.33–7.28 (m, 4H, 2 \times ArH), 7.13–7.09 (m, 1H, ArH), 2.59 (s, 2H, Pt-OH, exchangeable with D_2O). IR (KBr, cm^{-1}) ν 3500 (N-H), 3445 (PtO-H), 540 (Pt-O), and 348 (Pt-Cl) cm^{-1} . Anal. Calcd. for $\text{C}_{14}\text{H}_{14}\text{Cl}_2\text{N}_4\text{O}_2\text{Pt}$: C, 31.36; H, 2.63; N, 10.45. Found: C, 31.30; H, 2.29; N, 10.03%.

[Diacetato-dichloro-bis(benzimidazole)platinum(IV)]
[PtL₂Cl₂(OCOCH₃)₂] (4)

1 (0.452 g, 0.90 mmol) was dissolved in 1.37 mL acetic acid, 0.23 mL acetic acid anhydride, and 0.12 mL of 30% H_2O_2 . The reaction mixture, protected from light, was stirred at room temperature for 6 days. The resulting crude precipitate was filtered off and purified by repeated washing with small portions of water, ethanol, acetone, and diethylether, and dried *in vacuo* to yield 0.130 g (23.28%). ^1H NMR (400 MHz, DMSO- d_6) δ 13.90 (s, 2H, 2 \times N-H, exchangeable with D_2O), 8.77 (s, 2H, 2 \times ArH), 7.61–7.59 (d, J = 8.11 Hz, 2H, 2 \times ArH), 7.39–7.37 (d, J = 8.41 Hz, 2H, 2 \times ArH), 7.27–7.23 (m, 2H, 2 \times ArH), 7.07–7.03 (m, 2H, 2 \times ArH), 1.89 (s, 6H, 2 \times OCOCH₃). IR (KBr, cm^{-1}) ν 3557 (N-H), 1664 (C=O), and 340 (Pt-Cl) cm^{-1} . Anal. Calcd. for $\text{C}_{18}\text{H}_{18}\text{Cl}_2\text{N}_4\text{O}_4\text{Pt}$: C, 34.85; H, 2.92; N, 9.03. Found: C, 34.49; H, 2.62; N, 9.20%.

[Tetrachloro-bis(benzimidazole)platinum(IV)] [PtL₂Cl₄] (5)

4 (0.062 g, 0.10 mmol) was dissolved in aqueous 15% HCl (5 mL). The reaction mixture, protected from light, was stirred at room temperature for 3 days. The resulting crude yellow precipitate was filtered off and purified by repeated washing with small portions of water, ethanol, acetone, and diethylether, and dried *in vacuo* to yield 0.035 g (61.07%). ^1H NMR (400 MHz, DMSO- d_6) δ 13.80 (s, 2H, 2 \times N-H, exchangeable with D_2O), 8.88 (s, 2H, 2 \times ArH), 7.73–7.66 (m, 2H, 2 \times ArH), 7.43–7.23 (m, 4H, 2 \times ArH), 7.18–7.06 (m, 2H, 2 \times ArH). IR (KBr, cm^{-1}) ν 3510 (N-H), 335 (Pt-Cl) cm^{-1} . Anal. Calcd. for $\text{C}_{14}\text{H}_{12}\text{Cl}_4\text{N}_4\text{Pt}$: C, 29.34; H, 2.11; N, 9.77. Found: C, 29.29; H, 2.26; N, 9.52%.

Preliminary cytotoxicity test

Cell lines and growth conditions

Cisplatin and carboplatin were obtained from Sigma with a purity of more than 99.9%. The human MCF-7 breast, HeLa cervix, and HEp-2 larynx cancer cell lines used in this study were obtained from the Department of Virology, Faculty of Veterinary Medicine, University of Ankara. The cells were grown in Dulbecco's (Seromed, Germany) minimal essential medium (DMEM) enriched with 10% fetal calf serum (FCS) (Biochrom, Germany), 100 mg/mL streptomycin, and 100 IU/mL penicillin in a humidified atmosphere of 5% CO_2 at 37°C. The cells were harvested using trypsin (Bibco Life Technologies, UK)/Versene (0.05:0.02%) solution. *Mycoplasma* contamination was routinely monitored, and only *Mycoplasma*-free cultures were used.

In vitro chemosensitivity assay

The preliminary *in vitro* testing of compounds **1–5** on antitumor activity was carried out on human MCF-7 breast, HeLa cervix, and HEp-2 larynx cancer cells according to a previously published microtiter test²⁷. Briefly, the cells were seeded into 96-well plates (Greiner GmbH, Germany) in a volume of 100 μL to give 18–22 cells/microscopic area. After attachment to the culture surface, the cells were incubated in an atmosphere containing 5% CO_2 at 37°C for 24 h. At the end of this period the growth medium was carefully removed by suction, and 100 μL of fresh medium was added into each well. The medium used contained an adequate volume of a stock solution of the respective compound in order to obtain the desired test concentration (1, 5, 10, 20, 40, and 80 μM ; solvent, dimethylformamide (DMF); the complexes tested were added to the culture medium such that final DMF was 0.1% (v/v)). Sixteen wells were used for each complex (**1–5**, and reference compounds cisplatin and carboplatin) tested at individual concentrations, while 16 wells were reserved for the cell culture control, which contained the corresponding amount of DMF. After 72 h of incubation at 37°C, the medium was removed and the cells were fixed with 100 μL of 1% glutardialdehyde in phosphate-buffered saline (PBS) per well for 25 min. The fixative was replaced by 150 μL PBS/well and the plates were stored in a refrigerator (4°C). Cell biomass was determined by a crystal violet staining technique²⁸. Absorbance was measured at 492 nm using a Titertek Multiscan plus MKII Autoreader. The results correspond to three independent experiments.

IC_{50} values of the complexes **1–5** and reference compounds cisplatin and carboplatin were calculated (using Prism4 GraphPad software) from the dose–survival curves for the growth inhibition of all cell lines.

All cytotoxicity tests were performed as three independent experiments.

Interaction with pBR322 plasmid DNA

Cisplatin, ethidium bromide, and agarose, and plasmid DNA pBR322 and enzyme BamH1, were purchased from Sigma and Promega, respectively.

The interaction of cisplatin and synthesized compounds **1–5** with pBR322 plasmid DNA (Promega, cat. no: D1511) was studied by agarose gel electrophoresis based on a method described by Stellwagen²⁹. Stock solutions of the tested substances in DMF were prepared and used within 1 h. The final amount of DMF never exceeded 0.1%. pBR322 plasmid DNA (0.15 $\mu\text{g}/\text{mL}$) was incubated in the presence of increasing concentrations of **1–5** and cisplatin ranging from 0.625 to 160 μM and 0.625 to 10 μM , respectively, in a shaking water bath at 37°C for 4 h. Five microliters of drug–plasmid DNA mixture was loaded onto 1.5% agarose gel and electrophoresis was carried out under TAE (Tris base, acetic acid, and ethylenediaminetetraacetic acid (EDTA)) buffer for 2 h at 5 V/cm. At the end of electrophoresis, the gel was stained in the same buffer containing ethidium bromide (0.5 mg/mL) and visualized under UV transillumination using a gel documentation system (Vilber Lourmat, Marne la Vallée, France). The experiments were repeated three times.

Restriction endonuclease reaction with BamHI enzyme

In this series of experiments, a set of synthesized compounds **1–5** and cisplatin–plasmid DNA mixtures the same as that described previously was first incubated for 4 h in a shaking water bath at 37°C and then subjected to enzyme BamHI digestion. To each 10 μ L **1–5**/cisplatin–plasmid DNA mixture was added 3 μ L of 10 \times RE buffer followed by the addition of 0.3 μ L BamHI (3U). The mixtures were left in a shaking water bath at 37°C for 1.5 h, at the end of which the reaction was terminated by rapid cooling. Following the digestion reaction, products were separated on 1.5% agarose gel, and then the gel was stained with ethidium bromide and visualized under UV transillumination using a gel documentation system (Vilber Lourmat) to detect digestion.

Result and discussion

Chemistry

The carrier ligand benzimidazole (**L**) was prepared according to the Phillips method²⁵ as shown in Scheme 1, and its melting point was in accordance with the literature²⁶.

The Pt(II) and Pt(IV) compounds **1–5** were synthesized and characterized by elemental analysis, IR, and ¹H NMR spectra (Scheme 1). Among the compounds synthesized, **1**, which was reported previously by us²⁰, was synthesized in this study as reported previously. The elemental analysis data for each complex were in good agreement with the empirical formula proposed.

The IR spectrum of **L** showed a very broad band in the region 3300–2300 cm^{-1} due to the imidazole N-H. The Pt(II) complexes exhibited N-H stretching bands centered at 3310 and 3287 cm^{-1} sharper than that of the free ligand, due to the breaking of tautomerism, indicating that the N-H group was not involved in the coordination^{26,30}. According to the kinetic *trans* effect³¹, the synthesis method used is expected to yield complexes with *cis* geometry. In the far-IR region of the complexes **1, 3–5** spectra, a new broad band with a half-width


of about 30 cm^{-1} appeared, assigned to ν (Pt-Cl) centered at $\sim 337 \text{ cm}^{-1}$. It is well known that *cis*-dichloro complexes should show two bands of medium intensity because the vibrations are additive, but in a lot of cases the second band is only a shoulder³². Although no shoulder was apparently observable at the Pt-Cl stretching band of compounds **1, 3–5** the broad nature of the band suggested the presence of overlapping bands in this domain. The Pt-I band of the diiodo complex **2** should show at 195–183 cm^{-1} in the far-IR region of the complex's spectrum³³, but the ν (Pt-I) stretching band for **2** could not be measured on the spectrophotometer used.

The IR spectra of the Pt(IV) complexes showed some characteristic changes when compared to those of the Pt(II) complexes and the free ligand spectrum. Pt-Cl stretching vibrations were shifted ($\sim 15 \text{ cm}^{-1}$) toward higher frequencies in the spectra of Pt(IV) complexes **3–5** upon oxidation³⁴. Pt(IV) complexes **3–5** exhibited only one ν (Pt-Cl) absorption, which is consistent with the literature³⁵. In the IR spectra of **3–5**, N-H stretching vibrations appeared at the highest frequencies from 3557 to 3500 cm^{-1} . Being influenced by the oxidation states of the central metal, the N-H vibrations in the Pt(IV) compounds were broad in comparison with that in the Pt(II) compound **1**^{22,36}. The hydroxyplatinum(IV) complex **3**, which was obtained by the oxidation reaction of **1** with hydrogen peroxide, was readily identified by its characteristic PtO-H stretching band at 3445 cm^{-1} and Pt-O stretching band at 540 cm^{-1} ^{22,37,38}.

The insolubility of the complexes in the other organic solvents made it necessary to record ¹H NMR spectra in DMSO- d_6 . All ¹H NMR measurements were recorded immediately, in order to avoid the ligand exchange reaction between the platinum complexes synthesized and DMSO- d_6 . The ¹H-NMR spectral data of the ligand and the complexes are presented above in the "Experimental" section. The ¹H NMR spectra of all complexes were consistent with their corresponding protons, both in the chemical shifts and in the number of hydrogens³⁹. The spectra of the complexes compared to those of the free ligand showed considerable differences. The large downfield shifts in the imidazole N-H signal in the spectra of all complexes with respect to their ligands are a result of an increase in the N-H acid character after platinum binding. In all the complexes the aromatic proton resonances were observed in two main groups of lines. Two protons of the lower field pair were shifted downfield considerably, while the others were shifted slightly. It was assumed that the protons nearest to the tertiary nitrogen of the imidazole ring were the most deshielded, and would be observed at a lower field due to coordination of the benzimidazole ring to platinum through the -N= group.

Biological evaluation

The preliminary *in vitro* antiproliferative activities of compounds **1–5** and cisplatin and carboplatin used as reference compounds were determined on the human MCF-7, HeLa, and HEp-2 cell lines, according to a previously published microtiter assay^{27,28}, and the results are shown in Table 1. The Pt(II) complex **1**, bearing chloro ligands as leaving groups, was found to be more active than **2**, bearing iodo ligands as



Scheme 1. Synthesis of carrier ligand and Pt(II) and Pt(IV) complexes.

Table 1. IC₅₀ (μM) values of Pt(II) and Pt(IV) complexes on selected human tumor cell lines.

Compound	IC ₅₀ (μM)		
	MCF-7	HeLa	HEp-2
1	18.8 ± 3.4	39.5 ± 1.7	1.3 ± 0.4
2	>100	41.8 ± 2.0	>100
3	>100	>100	7.2 ± 1.9
4	41.9 ± 4.0	22.5 ± 3.0	14.6 ± 3.5
5	>100	20.7 ± 3.8	>100
Cisplatin	3.5 ± 1.1	1.3 ± 0.3	15.8 ± 2.0
Carboplatin	27.5 ± 2.6	14.8 ± 4.4	>100

leaving groups, on all cell lines used. These findings are consistent with data reported in the literature that the cisplatin analogs, containing chloro ligands, exhibit much stronger therapeutic features than analogs in which the ligands are azides, cyanides, iodides, or rhodanates⁴⁰. Moreover, other data in the literature report that dibromo and diiodo analogs of cisplatin hydrolyze at progressively slower rates compared to the dichloro complex, and this reactivity is reflected in the order of antitumor potency: I⁻ < Br⁻ < Cl⁻. It was reported that this could be attributed to an increase in kinetic reactivity toward water in going from the iodo to the chloro complex. This order implies that both electronic and steric factors influence the activity⁴¹. It is also reported that the binding to DNA of these drugs with halogeno leaving groups is almost certainly preceded by aquation, with the loss of one or more of the leaving groups⁴², and also iodo is generally a poorer leaving group than chloro in the structure of Pt(II) complexes⁴³. In the test on the HEp-2 cell line, compound **1**, which was found to be almost 12-fold more active than cisplatin, caused a significant reduction of cell growth.

In the case of the Pt(IV) complexes, the compounds **3** bearing hydroxo and **4** bearing acetato groups as axial ligands were found to be two-fold more active than cisplatin and as active as cisplatin, respectively, against the HEp-2 cancer cell line. It is interesting to note that **3** did not show any significant cytotoxicity against the other two cancer cell lines used. Among the complexes synthesized, Pt(IV) complex **5** bearing chloro axial ligands had no activity against MCF-7 and HEp-2 cell lines at the concentration range tested.

In order to detect whether synthesized compounds **1–5** induce conformational changes in the DNA helix and whether there is a relationship between the plasmid-DNA binding affinity and the cytotoxicity of the compounds, the effect of the binding of compounds **1–5** to pBR322 plasmid DNA was determined by the ability of these compounds to alter the electrophoretic mobility and intensity of the covalently closed circular form I and open circular form II bands (Figure 1). When pBR322 plasmid DNA was allowed to interact with increasing concentrations of cisplatin (Figure 1a), the mobility of both form I and form II bands increased essentially at the same rate, so that the bands remained parallel at all concentrations of cisplatin. The behavior of the gel electrophoretic mobility of both forms of pBR322 plasmid DNA-cisplatin adducts is consistent with previous reports^{44,45}.

However, the increase in mobility of the form II band of the plasmid DNA was more pronounced in the case of **2** than **1**;




Figure 1. Electrophoretograms relating to the interaction of pBR322 plasmid DNA with increasing concentration of cisplatin and **1–5**. Lane 1 in both electrophoretograms contains untreated pBR322 plasmid DNA to serve as a control. For cisplatin, lane 2: 0.625 μM, lane 3: 1.25 μM, lane 4: 2.5 μM, lane 5: 5 μM, lane 6: 10 μM. For **1–5**, lane 2: 0.625 μM, lane 3: 1.25 μM, lane 4: 2.5 μM, lane 5: 5 μM, lane 6: 10 μM, lane 7: 20 μM, lane 8: 40 μM, lane 9: 80 μM, lane 10: 160 μM. Roman numerals I and II indicate form I (covalently closed circular) and form II (open circular) plasmids, respectively.

for **1** and **2** the increase in mobility of the form I and form II bands was smaller than that induced by cisplatin. The lower activity of **1** and **2** as compared to cisplatin may be interpreted by their reduced ability to bind with DNA because of a greater steric constraint introduced by bulkier benzimidazole carrier ligands. The change in mobility of pBR322 plasmid DNA bands as a result of the interaction with cisplatin is believed to be due to intrastrand bifunctional Pt(GG) and monofunctional Pt(G) adducts^{5,46}. This may be interpreted in that the increase in mobility of form I and form II bands of pBR322 plasmid DNA with increasing concentration of **1** and **2** is indicative of the conformational change that occurs on the formation of bifunctional and monofunctional platinum adducts. For Pt(IV) complexes **3–5**, no significant changes in mobility of the forms I and II bands of plasmid DNA were observed at the concentration range tested for 4 h incubation time.

In order to assess whether the benzimidazole-platinum complexes show affinity toward the guanine-guanine (GG) region, we carried out restriction endonuclease analysis of the compound-pBR322 plasmid DNA adducts digested by BamH1 enzyme. BamH1 is a restriction enzyme that recognizes G/GATCC and hydrolyzes the phosphodiester bond between adjacent guanine sites. pBR322 plasmid DNA has only one binding site for BamH1 that converts form I and form II DNA to linear form III DNA^{44,47}.

As the concentration of the compounds tested was increased it was seen that BamH1 digestion was increasingly prevented (Figure 2). This is probably due to conformational




Figure 2. Electrophoretograms relating to the incubated mixtures of pBR322 plasmid DNA and increasing concentration of cisplatin and **1–5** followed by BamH1 digestion. For both cisplatin and **1–5**, lane 1 is untreated and undigested with BamH1, lane 2 is untreated pBR322 plasmid DNA but digested with BamH1. For cisplatin, lane 3: 0.625 μ M, lane 4: 1.25 μ M, lane 5: 2.5 μ M, lane 6: 5 μ M, lane 7: 10 μ M, lane 8: 20 μ M. For **1–5**, lane 3: 0.625 μ M, lane 4: 1.25 μ M, lane 5: 2.5 μ M, lane 6: 5 μ M, lane 7: 10 μ M, lane 8: 20 μ M, lane 9: 40 μ M, lane 10: 80 μ M, lane 11: 160 μ M. Roman numerals I, II, and III indicate form I (covalently closed circular), form II (open circular), and form III (linear) plasmids, respectively.

change in the DNA brought about by covalent binding of the compounds with the GG region of the plasmid DNA. Arranging in the order of decreasing prevention of BamH1 digestion by the compounds gives: cisplatin > **1, 2** > **5** > **4** > **3**. These results are consistent with the data reported in literature that the reduction rates and potentials of some diam(m)ine Pt(IV) complexes are dependent on the nature of the axial and equatorial ligands, but the axial ligands generally exert the stronger influence. It has also been reported that reduction occurs most readily when the axial ligands are chloro, least readily when they are hydroxo, and is intermediate when they are carboxylato¹³.

In the literature, it has been reported that steric constraint, hindering access to DNA for platinum complexes, would be greater in Pt(IV) complexes than in Pt(II) complexes, and Pt(IV)–DNA adducts are formed slowly as compared with Pt(II)–DNA adducts^{10,48}. Furthermore, small changes in mobility of the form I band of pBR322 plasmid DNA treated with some Pt(II) and Pd(II) complexes containing sterically demanding carrier-ligands were attributed to monofunctional binding⁴⁹. Considering the data reported in the literature and the preliminary results obtained in this study, it may be concluded that although Pt(IV) complexes **3–5** did not modify the electrophoretic mobility of plasmid DNA significantly, the formation of adequate adducts, presumably monofunctional adducts, may be sufficient to cause enough conformational change in pBR322 plasmid DNA treated with these compounds such that BamH1 digestion is

prevented below their concentration, 10 μ M. The conclusion drawn for Pt(IV) complexes **3–5** that their binding to plasmid DNA may be monofunctional for the incubation time of 4 h is consistent with the data obtained in the literature that monofunctional binding of octahedral Pt(IV) to DNA through the N7 guanine base is feasible¹¹.

In summary, it was found that there is no clear correlation between the axial and equatorial ligand variation and plasmid DNA binding affinity and the *in vitro* antiproliferative activity of the platinum(II) and platinum(IV) complexes synthesized, which modified the tertiary structure of pBR322 plasmid DNA. It can also be concluded that the *in vitro* antiproliferative activity of compounds **1, 3**, and **4** against the HEP-2 cell line is noteworthy, and must be taken into consideration in future studies.

Acknowledgements

We would like to thank Prof. Dr. İbrahim Burgu (Faculty of Veterinary Medicine, Ankara University) for his valuable support.

Declaration of interest

Financial support of part of this work (ECZ-F-EMB(SU) 2006-1) by the Research Foundation of Mersin University is gratefully acknowledged.

References

- Jung Y, Lippard SJ. Direct cellular responses to platinum-induced DNA damage. *Chem Rev* 2007;107:1387–407.
- Rabik CA, Dolan ME. Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat Rev* 2007;33:9–23.
- Jamieson ER, Lippard SJ. Structure, recognition, and processing of cisplatin–DNA adducts. *Chem Rev* 1999;99:2467–98.
- Chválová K, Brabec V, Kašpárková J. Mechanism of the formation of DNA–protein cross-links by antitumor cisplatin. *Nucleic Acids Res* 2007;35:1812–21.
- Fichtinger-Schepman AMJ, Van der Veer JL, den Hartog JHJ, Lohman PHM, Reedijk J. Adducts of the antitumor drug cis-diamminedichloroplatinum(II) with DNA: formation, identification, and quantitation. *Biochemistry* 1985;24:707–13.
- Fuertes MA, Alonso C, Perez JM. Biochemical modulation of cisplatin mechanisms of action: enhancement of antitumor activity and circumvention of drug resistance. *Chem Rev* 2003;103:645–62.
- Brabec V, Kasparkova J. Modification of DNA by platinum complexes. Relation to resistance of tumors to platinum antitumor drugs. *Drug Resist Update* 2005;8:131–46.
- Reedijk J. Improved understanding in platinum antitumor chemistry. *Chem Commun* 1996;7:801–6.
- Wong E, Giandomenico CM. Current status of platinum-based antitumor drugs. *Chem Rev* 1999;99:2451–66.
- Nováková O, Vrána O, Kiseleva VI, Brabec V. DNA interactions of antitumor platinum(IV) complexes. *Eur J Biochem* 1995;228:616–24.
- Ellis LT, Er HM, Hambley TW. The influence of the axial ligands of a series of platinum(IV) anti-cancer complexes on their reduction to platinum(II) and reaction with DNA. *Aust J Chem* 1995;48:793–806.
- Hall MD, Hambley TW. Platinum(IV) antitumor compounds: their bioinorganic chemistry. *Coord Chem Rev* 2002;232:49–67.
- Choi S, Filotto C, Bisanzo M, Delaney S, Lagasee D, Whitworth JL, et al. Reduction and anticancer activity of platinum(IV) complexes. *Inorg Chem* 1998;37:2500–4.
- Hall MD, Mellor HR, Callaghan R, Hambley TW. Basis for design and development of platinum(IV) anticancer complexes. *J Med Chem* 2007;26:3403–11.

15. Abu-Surrah AS, Kettunen M. Platinum group antitumor chemistry: design and development of new anticancer drugs complementary to cisplatin. *Curr Med Chem* 2006;13:1337-57.
16. Kerr C. Satraplatin for hormone-refractory prostate cancer. *Lancet Oncol* 2007;8:290.
17. Gümüş F, İzgü F, Algül Ö. Synthesis and structural characterization of some 5(6)-substituted-2-hydroxymethylbenzimidazole derivatives and their platinum(II) complexes and determination of their in vitro antitumor activities by "Rec-Assay" test. *FABAD J Pharm Sci* 1996;21:7-15.
18. Gümüş F, Algül Ö. DNA binding studies with cis-dichlorobis(5(6)-non/chlorosubstituted-2-hydroxymethyl-benzimidazole)platinum(II) complexes. *J Inorg Biochem* 1997;68:71-4.
19. Gümüş F, Pamuk İ, Özden T, Yıldız S, Diril N, Öksüzöğlü E, et al. Synthesis, characterization and in vitro cytotoxic, mutagenic and antimicrobial activity of platinum(II) complexes with substituted benzimidazole ligands. *J Inorg Biochem* 2003;94:255-62.
20. Gümüş F, Algül Ö, Eren G, Eroğlu H, Diril N, Gür S, et al. Synthesis, cytotoxic activity on MCF-7 cell line and mutagenic activity of platinum(II) complexes with 2-substituted benzimidazole ligands. *Eur J Med Chem* 2003;38:473-80.
21. Gökçe M, Utku S, Gür S, Özkul A, Gümüş F. Synthesis, in vitro cytotoxic and antiviral activity of cis-[Pt(R(-) and S(+)-2- α -hydroxybenzylbenzimidazole)₂Cl₂] complexes. *Eur J Med Chem* 2005;40:135-41.
22. Utku S, Gümüş F, Gür S, Özkul A. Synthesis and cytotoxic activity of platinum(II) and platinum(IV) complexes with 2-hydroxymethylbenzimidazole or 5(6)-chloro-2-hydroxymethylbenzimidazole ligands against MCF-7 and HeLa cell lines. *Türk J Chem* 2007;31:503-14.
23. Utku S, Gümüş F, Karaoğlu T, Özkul A. Cytotoxic activity of platinum(II) and platinum(IV) complexes bearing 5(6)-non/chlorosubstituted-2-hydroxymethylbenzimidazole ligands against Hep-2 cell line. *Ankara Ecz Fak Derg* 2007;36:21-30.
24. Gümüş F, Eren G, Açıık L, Çelebi A, Öztürk F, Yılmaz Ş, et al. Synthesis, cytotoxic, and DNA interactions of new cisplatin analogues containing substituted benzimidazole ligands. *J Med Chem* 2009;52:1345-57.
25. Preston PN. Synthesis, reactions, and spectroscopic properties of benzimidazoles. *Chem Rev* 1974;74:279-314.
26. Rabiger DJ, Joullie MM. The ionization constant, ultraviolet and infrared spectra of some substituted benzimidazoles. *J Org Chem* 1964;29:476-82.
27. Bernhard G, Reile H, Birnböck H, Spruss T, Schönenberger H. Standardized kinetic microassay to quantify differential chemosensitivity on the basis of proliferative activity. *J Cancer Res Clin Oncol* 1992;118:35-43.
28. Reile H, Birnböck H, Bernhard G, Spruss T, Schönenberger H. Computerized determination of growth kinetic curves and doubling times from cells in microculture. *Anal Biochem* 1990;187:262-7.
29. Stellwagen N. *Nucleic Acid Electrophoresis*. Berlin: Springer, 1998.
30. Maurya RC, Mishra DD. Studies on some novel mixed-ligand cyanonitrosyl hetero complexes of chromium(I) with potentially bi- and tri-dentate benzimidazole derivatives. *Syn React Inorg Met Org Chem* 1990;20:1013-24.
31. Tunalı NK, Özkul S. *Anorganik Tepkime Mekanizmaları*. Anorganik Kimya. Ankara: Gazi Publishing, 1999:373-402.
32. Mylonas S, Valavanidis A, Voukoulidis V, Polyssiou M. Platinum(II) and palladium(II) complexes with amino acid derivatives. Synthesis, interpretation of IR and ¹H NMR spectra and conformational implications. *Inorg Chim Acta* 1981;55:125-8.
33. Dodoff N, Grancharov K, Gugova R, Spassovska N. Platinum(II) complexes of benzoic- and 3-methoxybenzoic acid hydrazides. Synthesis, characterization, and cytotoxic effect. *J Inorg Biochem* 1994;54:221-33.
34. Giandomenico CM, Abrams MJ, Murrer BA, Vollano JF, Rheinheimer MI, Wyer SB, et al. Carboxylation of kinetically inert platinum(IV) hydroxy complexes. An entree into orally active platinum(IV) antitumor agents. *Inorg Chem* 1995;34:1015-21.
35. Miller B, Altman J, Beck W. Platinum(IV) complexes of vicinal-1,2-diamines, and bis(vicinal-1,2-diamines) with an acylamino function. Evidence for a platinum hydroperoxide intermediate upon oxidation. *Inorg Chim Acta* 1997;264:101-8.
36. Khokhar AR, Deng Y, Al-Baker S, Yoshida M, Siddik ZH. Synthesis and antitumor activity of ammine/amine platinum(II) and (IV) complexes. *J Inorg Biochem* 1993;51:677-87.
37. Lee EJ, Jun MJ, Sohn YS. Synthesis and properties of novel platinum(IV) complexes involving asymmetric chiral diamines as carrier ligands. *Bull Korean Chem Soc* 1999;20:1295-8.
38. Hambley TW, Battle AR, Deacon GB, Lawrenz ET, Fallon GD, Gatehouse BM, et al. Modifying the properties of platinum(IV) complexes in order to increase biological effectiveness. *Inorg Biochem* 1999;77:3-12.
39. Lippert B. Platinum nucleobase chemistry. *Prog Inorg Chem* 1989;37:1-97.
40. Sawicka M, Kalinowska M, Skierska J, Lewandowski W. A review of selected anti-tumour therapeutic agents and reasons for multidrug resistance occurrence. *J Pharm Pharmacol* 2004;56:1067-81.
41. Kushev D, Grünert R, Spassovska N, Golovinsky E, Bednarski PJ. Unusual reactivity of cytotoxic cis-dihydrazide Pt(II) complexes in aqueous solution. *J Inorg Biochem* 2003;96:469-77.
42. Reedijk J. The relevance of hydrogen bonding in the mechanism of action of platinum antitumor compounds. *Inorg Chim Acta* 1992;198-200:873-81.
43. Whittaker J, McFadyen WD, Wickham G, Wakelin LPG, Murray V. The interaction of DNA-targeted platinum phenanthridinium complexes with DNA. *Nucleic Acids Res* 1998;26:3933-9.
44. Ushay HM, Tullius TD, Lippard SJ. Inhibition of the BamHI cleavage and unwinding of pBR322 deoxyribonucleic acid by the antitumor drug cis-dichlorodiammineplatinum(II). *Biochemistry* 1981;20:3744-8.
45. Cheng H, Huq F, Beale P, Fisher K. Synthesis, characterization, activities, cell uptake and DNA binding of trinuclear complex: [{trans-PtCl(NH₃)₂}]₂μ-{trans-Pd(NH₃)₂(2-hydroxypyridine)-(H₂N(CH₂)₆NH₂)₂]Cl₄. *Eur J Chem* 2006;41:896-903.
46. Keck MV, Lippard SJ. Unwinding of supercoiled DNA by platinum-ethidium and related complexes. *J Am Chem Soc* 1992;114:3386-90.
47. Bowler BE, Hollis LS, Lippard SJ. Synthesis and DNA binding and photophysical properties of acridine orange linked by a polymethylene tether to (1,2-diaminoethane)dichloroplatinum(II). *J Am Chem Soc* 1984;106:6102-4.
48. Perez JM, Camazon M, Alvarez-Valdes A, Quiroga AG, Kelland LR, Alonso C, et al. Synthesis, characterization and DNA modification induced by a novel Pt(IV)-bis(monoglutarate) complex which induced apoptosis in glioma cells. *Chem-Biol Interact* 1999;117:99-115.
49. Nepelchova K, Kasparkova J, Vrana O, Novakova O, Habtemariam A, Watchman B, et al. DNA interactions of new antitumor aminophosphine platinum(II) complexes. *Mol Pharmacol* 1999;56:20-30.