



Antiproliferative activity of some 1,4-dimethylcarbazoles on cells that express estrogen receptors: part I

Anna Caruso, Adele Chimento, Hussein El-Kashef, Jean-Charles Lancelot, Antonella Panno, Vincenzo Pezzi, Carmela Saturnino, Maria Stefania Sinicropi, Rosa Sirianni & Sylvain Rault

To cite this article: Anna Caruso, Adele Chimento, Hussein El-Kashef, Jean-Charles Lancelot, Antonella Panno, Vincenzo Pezzi, Carmela Saturnino, Maria Stefania Sinicropi, Rosa Sirianni & Sylvain Rault (2012) Antiproliferative activity of some 1,4-dimethylcarbazoles on cells that express estrogen receptors: part I, Journal of Enzyme Inhibition and Medicinal Chemistry, 27:4, 609-613, DOI: [10.3109/14756366.2011.603132](https://doi.org/10.3109/14756366.2011.603132)

To link to this article: <https://doi.org/10.3109/14756366.2011.603132>



Published online: 02 Sep 2011.



Submit your article to this journal [↗](#)



Article views: 672



View related articles [↗](#)



Citing articles: 5 View citing articles [↗](#)

SHORT COMMUNICATION

Antiproliferative activity of some 1,4-dimethylcarbazoles on cells that express estrogen receptors: part I

Anna Caruso¹, Adele Chimento², Hussein El-Kashef³, Jean-Charles Lancelot⁴, Antonella Panno¹, Vincenzo Pezzi², Carmela Saturnino⁵, Maria Stefania Sinicropi¹, Rosa Sirianni², and Sylvain Rault⁴

¹Department of Pharmaceutical Sciences, University of Calabria, Arcavacata di Rende (CS), Italy, ²Department of Pharmaco-Biology, University of Calabria, Arcavacata di Rende (CS), Italy, ³Chemistry Department, Faculty of Science, Assiut University, Assiut, Egypt, ⁴Université de Caen Basse-Normandie, U.F.R. des Sciences Pharmaceutiques, Centre d'Etudes et de Recherche sur le Médicament de Normandie, Boulevard Becquerel, Caen Cedex, France, and ⁵Department of Pharmaceutical and Biomedical Sciences, University of Salerno, Fisciano (SA), Italy

Abstract

Several 9H-carbazole derivatives are used for various pharmacological applications. Many of these compounds demonstrated cytotoxic and anticancer activities. In this work, we have investigated the cytotoxic activity of some substituted carbazoles against cancer cell lines (MCF-7, and ISK). The derivative **2a** showed the highest inhibitory activity against both cell lines.

Keywords: Carbazoles, anticancer, MCF-7 cell line, Ishikawa cell line

Introduction

The indole nucleus is often found embedded in compounds with a wide range of biological activities¹. Carbazoles, whether synthetic or naturally occurring, are important members of indole-containing heterocycles^{1,2}. Several 9H-carbazoles possess various pharmacological applications and also they are used in dyes-, insecticide-, and plastic industries. In addition, several derivatives of 1,2,3,4-tetrahydro-9H-carbazole and 9-methyl-9H-carbazole have been known by their medicinal use as antitubercular, antifungal, antibacterial, β_3 -adrenoceptor agonists and anticancer activities^{2–4}.

The precise mechanisms of their anti-neoplastic activity have not yet been explained. Many of these carbazoles are cytostatic and their activity results from alternative cytotoxic effects. It was suggested that the prevalent mechanisms of the antitumor, mutagenic and cytotoxic activities are their intercalation into DNA and inhibition of DNA-topoisomerase II activity^{5,6}.

Based on this, the aim of this work is to investigate the cytotoxic activity of the prepared carbazoles (Figure 1)

against a cancer cell line. We have used MCF-7 cell line derived from a pleural effusion of a postmenopausal 69-year-old patient with metastatic breast cancer in 1970⁷. This cell line was grown in Dulbecco's modified Eagle's medium (DMEM), without phenol red, containing 10% foetal calf serum (Invitrogen, Milan, Italy) and 1 mg/mL penicillin-streptomycin. The MCF-7 cell line is the most widely used and best characterized of all the human breast cancer cell lines⁸. MCF-7 cells respond to estrogens and anti-estrogens, have differential expression of estrogen receptors (ER), progesterone receptor, and have high proliferation rates⁹. MCF-7 cells are also a perfect model to study the pathway of malignant progression as they can be subjected to appropriate endocrinologic and physiologic selective pressures for the derivation of variants with more progressed phenotypes⁸.

Further in this work, some tests were performed on Ishikawa cell line (ISK) derived from an endometrial adenocarcinoma of a 39-year-old Asian woman¹⁰. This cell line was maintained in Dulbecco's modified Eagle's

Address for Correspondence: Prof. Maria Stefania Sinicropi, Department of Pharmaceutical Sciences, University of Calabria, 87036 Arcavacata di Rende (CS), Italy. E-mail: s.sinicropi@unical.it

(Received 09 June 2011; revised 29 June 2011; accepted 29 June 2011)




Figure 1. Carbazole derivatives (1a-g).

medium-F12 (DMEM/F12), without phenol red, supplemented with 10% foetal bovine serum (Invitrogen, Milan, Italy) and 1 mg/mL penicillin-streptomycin.

Ishikawa cells express estrogen¹¹, progesterone¹², androgen¹³ and aryl hydrocarbons receptors¹⁴. There is a big evidence for the hormonal responsiveness of Ishikawa cells. This evidence is based mainly on hormonal modification of proliferation, cellular functions or gene expression¹⁵. In particular, these cells are




Figure 2. Effect of carbazole derivatives (1a-e) against MCF-7 and ISK cell proliferation. MCF-7 (A) and ISK (B) cells were treated for 96 or 24 h, respectively, after 24-h starvation, with the indicated concentrations of substances 1a, 1b, 1c, 1d and 1e. Cells proliferation was evaluated by MTT assay. Statistically significant differences are indicated. Histograms; mean of three independent experiments each performed with triplicate samples expressed as percent of basal; bars, SE (* $P < 0.01$ compared with basal).




Figure 3. Effect of carbazole derivatives (**1f-g**) against MCF-7 and ISK cell proliferation. MCF-7 (**A**) and ISK (**B**) cells were treated for 96 or 24 h, respectively, after 24-h starvation, with the indicated concentrations of substances **1f** and **1g**. Cells proliferation was evaluated by MTT assay. Statistically significant differences are indicated. Histograms; mean of three independent experiments each performed with triplicate samples expressed as percent of basal; bars, SE (* $P < 0.01$ compared with basal).

used *in vitro* for the elucidation of molecular mechanisms of hormone action (e.g., in drug and discovery processes), testing of potential agonistic functions of anti-estrogens or selective estrogen receptor modulators in an endometrial-derived model¹⁶, in studies of ligand independent activation of the estrogen receptor and anchorage independent tumour growth¹⁷. In addition, they are used for studies on factors controlling hormonal receptivity¹⁸ and on environmental toxicology on the function of phytoestrogens¹⁹ and endocrine-disrupting chemicals²⁰ in an endometrial model; paracrine cell/cell-interaction²¹, signalling cross-talk¹⁴ and others²². Because of their hormonal responsiveness *in vitro*, Ishikawa endometrial adenocarcinoma cells have also been developed as an estrogen sensitive *in vivo* tumour model, which is particularly suitable for the study of hormonal growth control²³.

Results and discussion

Carbazoles **1a–g** were synthesized^{24–27} and characterized^{28–34} using earlier reported methods including ours.

In particular, the 9H-carbazoles **1a–d** were prepared by the reaction of appropriate indoles with hexane-2,4-dione in the presence of *p*-toluenesulphonic acid. The 9H-carbazole **1e** was instead prepared by reaction of **1a** with ethyl chloroformate under standard conditions³⁵. Furthermore, we have used **1a** as starting compound for the preparation of **1f–g**. In fact, **1a** was *N*-methylated by iodomethane and in the presence of sodium hydride to give compound **1f**. The carbazole NH could be *N*-protected by (Boc)₂O to give **1g** (Figure 1).

In order to evaluate the cytotoxic activity of prepared compounds (**1a–g**) against MCF-7 and ISK cell lines, an MTT assay was performed³⁶ using increasing concentrations (0.1, 1, 10 µM) of the compounds under testing. Compound **1a** showed a significant inhibition at 1 and 10 µM, against both MCF-7 and ISK cell lines (Figure 2A, and B). Concerning the MCF-7 cells, a 30% inhibition was observed by the lower concentrations of 0.1 and 1 µM of **1b** and **1c**, respectively (Figure 2A). However, with ISK cells, compound **1b** displayed a significant inhibitory effect on cell proliferation only at 10 µM concentration, while the inhibitory effect of **1c** was observed at a concentration of 0.1 and 10 µM (Figure 2B). On the other hand, **1d** showed an inhibitory effect against MCF-7 cells at 1 µM concentration (Figure 2A), and the same compound, however, displayed more inhibition against ISK cells at a concentration of 1 and 10 µM (Figure 2B). Compound **1e** significantly reduced cell proliferation of both MCF-7 and ISK cells at a concentration of 0.1 µM and 1 µM (Figure 2A, and B).

Based on the above results indicating the ability of compound **1a** to inhibit proliferation of both cell lines (Figure 2A, and B), structural modifications of **1a** were made in order to find out whether the introduction of a substituent such as a methyl- (**1f**) or a Boc substituent

(**1g**), at the 9 position of the carbazole nucleus could affect the antiproliferative activity.

Figure 3 showed that the introduction of a methyl group (an electron-donating small substituent), improved the inhibitory activity, whereas the introduction of Boc group did not show any improvement of activity. The derivative **1f** showed the highest inhibitory activity against both cell lines (at low concentration of 0.1 µM) (Figure 3A, and B), the matter that encourages its use as a possible antitumor agent. However, the observation that higher concentrations (1, 10 µM) of **1f** do not induce dose-response inhibitory effect on both MCF-7 and ISK cell viability suggests a possible role of **1f** in blocking a survival pathway binding to receptor desensitized at high concentration. This hypothesis is currently under investigation in our laboratories.

Acknowledgements

We are thankful to Dr. Ewa Surmacz (Sbarro Institute for Cancer Research and Molecular Medicine, Philadelphia, PA, USA) and to Dr. D. Picard (University of Geneva, Geneva, Switzerland) for their kind offer of MCF-7 and Ishikawa cell lines, respectively.

Declaration of interest

The authors report no conflicts of interest.

References

- De Koning CB, Michael JP, Rousseau AL. A versatile and convenient method for the synthesis of substituted benzo[a]carbazoles and pyrido[2,3-a]carbazoles. *J Chem Soc Perkin Trans* 2000;1:1705–1713.
- Caruso A, Lancelot JC, El-Kashef H, Sinicropi MS, Legay R, Lesnard A, Rault S. A rapid and versatile synthesis of novel pyrimido[5,4-b]carbazoles. *Tetrahedron* 2009;65:10400–10405.
- Caruso A, Voisin-Chiret AS, Lancelot JC, Sinicropi MS, Garofalo A, Rault S. Efficient and simple synthesis of 6-aryl-1,4-dimethyl-9H-carbazoles. *Molecules* 2008;13:1312–1320.
- Waldau D, Mikolasch A, Lalk M, Schauer F. Derivatization of bioactive carbazoles by the biphenyl-degrading bacterium *Ralstonia* sp. strain SBUG 290. *Appl Microbiol Biotechnol* 2009;83:67–75.
- Auclair C. Multimodal action of antitumor agents on DNA: The ellipticine series. *Arch Biochem Biophys* 1987;259:1–14.
- Stiborova M, Rupertova M, Schmeiser HH, Frei E. Molecular mechanisms of antineoplastic action of an anticancer drug ellipticine. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2006;150:13–23.
- Soule HD, Vazquez J, Long A, Albert S, Brennan M. A human cell line from a pleural effusion derived from a breast carcinoma. *J Natl Cancer Inst* 1973;51:1409–1416.
- Bullinger D, Neubauer H, Fehm T, Laufer S, Gleiter CH, Kammerer B. Metabolic signature of breast cancer cell line MCF-7: Profiling of modified nucleosides via LC-IT MS coupling. *BMC Biochem* 2007;8–25.
- Matthews J, Gustafsson JA. Estrogen signaling: A subtle balance between ER α and ER β . *Mol Interv* 2003;3:281–292.
- Nishida M, Kasahara K, Kaneko M, Iwasaki H, Hayashi K. Establishment of a new human endometrial adenocarcinoma cell line, Ishikawa cells, containing estrogen and progesterone receptors. *Nippon Sanka Fujinka Gakkai Zasshi* 1985;37:1103–1111.

11. Holinka CF, Hata H, Kuramoto H, Gursipide E. Responses to estradiol in a human endometrial adenocarcinoma cell line (Ishikawa). *J Steroid Biochem* 1986;24:85–89.
12. Lessey BA, Ilesanmi AO, Castelbaum AJ, Yuan L, Somkuti SG, Chwalisz K et al. Characterization of the functional progesterone receptor in an endometrial adenocarcinoma cell line (Ishikawa): Progesterone-induced expression of the $\alpha 1$ integrin. *J Steroid Biochem Mol Biol* 1996;59:31–39.
13. Lovely LP, Appa Rao KB, Gui Y, Lessey BA. Characterization of androgen receptors in a well-differentiated endometrial adenocarcinoma cell line (Ishikawa). *J Steroid Biochem Mol Biol* 2000;74:235–241.
14. Wormke M, Castro-Rivera E, Chen I, Safe S. Estrogen and aryl hydrocarbon receptor expression and crosstalk in human Ishikawa endometrial cancer cells. *J Steroid Biochem Mol Biol* 2000;72:197–207.
15. Vollmer G. Endometrial cancer: Experimental models useful for studies on molecular aspects of endometrial cancer and carcinogenesis. *Endocr Relat Cancer* 2003;10:23–42.
16. Labrie F, Labrie C, Bélanger A, Simard J, Giguère V, Tremblay A et al. EM-652 (SCH57068), a pure SERM having complete antiestrogenic activity in the mammary gland and endometrium. *J Steroid Biochem Mol Biol* 2001;79:213–225.
17. Holinka CF, Anzai Y, Hata H, Kimmel N, Kuramoto H, Gursipide E. Proliferation and responsiveness to estrogen of human endometrial cancer cells under serum-free culture conditions. *Cancer Res* 1989;49:3297–3301.
18. Apparao KB, Murray MJ, Fritz MA, Meyer WR, Chambers AF, Truong PR et al. Osteopontin and its receptor $\alpha \beta (3)$ integrin are coexpressed in the human endometrium during the menstrual cycle but regulated differentially. *J Clin Endocrinol Metab* 2001;86:4991–5000.
19. Frigo DE, Duong BN, Melnik LI, Schief LS, Collins-Burow BM, Pace DK et al. Flavonoid phytochemicals regulate activator protein-1 signal transduction pathways in endometrial and kidney stable cell lines. *J Nutr* 2002;132:1848–1853.
20. Bergeron RM, Thompson TB, Leonard LS, Pluta L, Gaido KW. Estrogenicity of bisphenol A in a human endometrial carcinoma cell line. *Mol Cell Endocrinol* 1999;150:179–187.
21. Arnold JT, Lessey BA, Seppälä M, Kaufman DG. Effect of normal endometrial stroma on growth and differentiation in Ishikawa endometrial adenocarcinoma cells. *Cancer Res* 2002;62:79–88.
22. Kanishi Y, Kobayashi Y, Noda S, Ishizuka B, Saito K. Differential growth inhibitory effect of melatonin on two endometrial cancer cell lines. *J Pineal Res* 2000;28:227–233.
23. Nishida M, Kasahara K, Tsuji T, Kaneko M, Iwasaki H. Mechanisms of the antitumor action of gestagens on endometrial cancer. *Nippon Sanka Fujinka Gakkai Zasshi* 1986;38:2214–2222.
24. (a) Cranwell A, Saxton JE. A synthesis of ellipticine. *J Chem Soc* 1962;3482–3487; (b) Lancelot JC, Rault S, Robba M, Nguyen HD. Efficient synthesis of 6H-pyrido[3,2-b]carbazole derivatives from 3-amino-1,4-dimethylcarbazole. *Chem Pharm Bull* 1987;35:425–428.
25. (a) Tabka T, Letois B, Lancelot JC, Gauduchon P, Heron JF, Le Talaer JY, Rault S, Robba M. Etude de la cytotoxicité *in vitro* de dérivés du carbazole II. Nitro et amino halogéno-6 diméthyl-1,4 9H-carbazoles. *Eur J Med Chem*, 1989;24:605–610; (b) Testard A, Picot L, Fruitier-Arnaudin I, Piot JM, Chabane H, Domon L, Thiery V, Besson T. Microwave-assisted synthesis of novel thiazolocarbazoles and evaluation as potential anticancer agents. Part III. *J Enzym Inhib Med Chem*, 2004;19:467–473.
26. Moinet-Hedin V, Tabka T, Sichel F, Gauduchon P, Le Talaer JY, Saturnino C, Letois B, Lancelot JC, Robba M. *In vitro* cytotoxicity of carbazole derivatives. V. 9-Halogeno-substituted 5,11-dimethyl-6H-pyrido[3,2-b]carbazoles. *Eur J Med Chem* 1997;32:113–122.
27. Caruso A, Voisin-Chiret AS, Lancelot JC, Sinicropi MS, Garofalo A, Rault S. Novel and efficient synthesis of 5,8-dimethyl-9H-carbazol-3-ol via a hydroxydeboronation reaction. *Heterocycles* 2007;71(10):2203–2210.
28. (1a) White solid, mp 174°C. IR (KBr) (cm⁻¹): 3517, 3415, 1461, 1165, 847, 809, 543. ¹H NMR (DMSO-d₆): δ 2.50 (s, 3H, CH₃), 2.61 (s, 3H, CH₃); 6.83 (d, J = 7.36 Hz, 1H, Ar); 6.95 (d, J = 8.56 Hz, 1H, Ar); 7.08 (d, J = 7.36 Hz, 1H, Ar); 7.40 (d, J = 8.56 Hz, 1H, Ar); 7.53 (s, 1H, Ar); 9.12 (br, 1H, OH); 10.88 (s, 1H, NH). MS (ESI⁺): 212 (M⁺ + 1).
29. (1b) White solid, mp 150°C. IR (KBr) (cm⁻¹): 3406, 2959, 1481, 1210, 1045, 812, 545. ¹H NMR (DMSO-d₆): δ 2.62 (s, 3H, CH₃), 2.69 (s, 3H, CH₃), 3.90 (s, 3H, OCH₃), 6.85 (d, J = 7.08 Hz, 1H, Ar); 7.09 (d, J = 8.28 Hz, 2H, Ar), 7.48 (d, J = 8.56 Hz, 1H, Ar), 7.64 (s, 1H, Ar), 11.01 (br, 1H, NH). MS (ESI⁺): 226 (M⁺ + 1), 224 (M⁺ - 1).
30. (1c) White solid, mp 266°C. IR (KBr) (cm⁻¹): 3337, 1702, 1615, 1430, 1290, 1122, 989, 803, 764, 735, 557. ¹H NMR (DMSO-d₆): δ 2.52 (s, 3H, CH₃), 2.71 (s, 3H, CH₃), 3.84 (s, 3H, OCH₃), 6.89 (d, J = 7.84 Hz, 1H, Ar); 7.11 (d, J = 7.32 Hz, 1H, Ar), 7.55 (d, J = 8.08 Hz, 1H, Ar), 7.98 (dd, J₁ = 1.48 Hz, J₂ = 8.56 Hz, 1H, Ar), 8.67 (s, 1H, Ar), 11.62 (s, 1H, NH). MS (ESI⁺): 254 (M⁺ + 1).
31. (1d) White solid, mp 253°C. IR (KBr) (cm⁻¹): 3339, 1705, 1618, 1430, 1295, 1129, 990, 806, 765, 738, 560. ¹H NMR (DMSO-d₆): δ 1.67 (s, 3H, CH₃), 2.53 (s, 3H, CH₃), 2.80 (s, 3H, CH₂CH₃), 4.30–4.60 (m, 2H, OCH₂), 6.95 (d, J = 7.50 Hz, 1H, Ar); 7.16 (d, J = 7.20 Hz, 1H, Ar), 7.55 (d, J = 8.70 Hz, 1H, Ar), 8.04 (d, J = 8.70 Hz, 1H, Ar), 8.72 (s, 1H, Ar), 11.59 (br, 1H, NH). MS (ESI⁺): 268 (M⁺ + 1).
32. (1e) White solid, mp 130°C. IR (KBr) (cm⁻¹): 3396, 1744, 1462, 1253, 1181, 1002, 811, 780, 552. ¹H NMR (DMSO-d₆): δ 1.32–1.40 (t, 3H, CH₂CH₃), 2.57 (s, 3H, CH₃), 2.78 (s, 3H, CH₃), 4.28–4.35 (q, 2H, CH₂CH₃), 6.91 (d, 1H, J = 7.08 Hz, Ar), 7.16 (d, 1H, J = 7.32 Hz, Ar), 7.29 (dd, 1H, J₁ = 1.72 Hz, J₂ = 8.56 Hz, Ar), 7.58 (d, 1H, J = 8.56 Hz, Ar), 7.94 (s, 1H, Ar), 11.36 (s, 1H, NH). MS (ESI⁺): 284 (M⁺ + 1).
33. (1f) White solid, mp 202°C. IR (KBr) (cm⁻¹): 3415, 1587, 1460, 1165, 809, 543. ¹H NMR (CDCl₃): δ 8.23 (d, J = 1.96 Hz, 1H, Ar); 7.52 (dd, J = 8.80 Hz, 1H, Ar); 7.23 (d, J = 8.80 Hz, 1H, Ar); 7.07 (d, J = 7.32 Hz, 1H, Ar); 6.88 (d, J = 7.32 Hz, 1H, Ar); 4.07 (s, 3H, NCH₃); 2.80 (s, 3H, CH₃); 2.78 (s, 3H, CH₃). MS (ESI⁺): 226 (M⁺ + 1).
34. (1g) White solid, mp 205°C. IR (KBr) (cm⁻¹): 3410, 1701, 1449, 1329, 1157, 807. ¹H NMR (CDCl₃): δ 1.64 (s, 9H, CH₃); 2.43 (s, 3H, CH₃); 2.74 (s, 3H, CH₃); 6.91 (d, J = 8.56 Hz, 1H, Ar); 6.98 (d, J = 7.32 Hz, 1H, Ar); 7.09 (d, J = 7.32 Hz, 1H, Ar); 7.46 (s, 1H, OH); 7.63 (d, J = 8.56 Hz, 1H, Ar); 7.85 (s, 1H, Ar). MS (EI) m/z (%): 311 (M⁺, 15), 255 (48) (M⁺, -tBu), 211 (100) (M⁺, -CO₂tBu).
35. Caruso A. Ph.D. Thesis. Synthesis of novel dimethylcarbazole, dimethylpyrimidocarbazole, benzofuro quinazolinone, benzothienoquinazolinone derivatives with potential anticancer activity. University of Calabria-Italy, 17.12. 2008.
36. Sylvester PW. Optimization of the tetrazolium dye (MTT) colorimetric assay for cellular growth and viability. *Methods Mol Biol* 2011;716:157–168.