



Inhibition of mammalian carbonic anhydrase isoforms I, II and VI with thiamine and thiamine-like molecules

Zafer Ömer Özdemir, Murat Şentürk & Deniz Ekin

To cite this article: Zafer Ömer Özdemir, Murat Şentürk & Deniz Ekin (2013) Inhibition of mammalian carbonic anhydrase isoforms I, II and VI with thiamine and thiamine-like molecules, Journal of Enzyme Inhibition and Medicinal Chemistry, 28:2, 316-319, DOI: [10.3109/14756366.2011.637200](https://doi.org/10.3109/14756366.2011.637200)

To link to this article: <https://doi.org/10.3109/14756366.2011.637200>



Published online: 06 Dec 2011.



Submit your article to this journal [↗](#)



Article views: 2063



View related articles [↗](#)



Citing articles: 6 View citing articles [↗](#)

RESEARCH ARTICLE

Inhibition of mammalian carbonic anhydrase isoforms I, II and VI with thiamine and thiamine-like molecules

Zafer Ömer Özdemir¹, Murat Şentürk², and Deniz Ekinci³

¹Department of Chemistry, Art and Science Faculty, Kırklareli University, Kırklareli, Turkey, ²Department of Chemistry, Art and Science Faculty, Ağrı İbrahim Çeçen University, Ağrı, Turkey, and ³Department of Agricultural Biotechnology, Faculty of Agriculture, Ondokuz Mayıs University, Samsun, Turkey

Abstract

Here we determined the *in vitro* inhibitory effects of 5-(2-hydroxyethyl)-3,4-dimethylthiazolium iodide (**1**), 3-Benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride (**2**) and thiamine (**3**) on human erythrocyte carbonic anhydrase I, II isozymes (hCA I and hCA II) and secreted isoenzyme CA VI. *K_i* values ranged from 0.38 to 2.27 μ M for hCA I, 0.085 to 0.784 μ M for hCA II and 0.062 to 0.593 μ M for hCA VI, respectively. The compounds displayed relatively strong actions on hCA II, in the same range as the clinically used sulfonamidesethoxzolamide, zonisamide and acetazolamide.

Keywords: Carbonic anhydrase, thiamine, inhibition

Introduction

Carbonic anhydrase (EC 4.2.1.1., CA) has been a well-characterized pH regulatory enzyme in most tissues including erythrocytes¹. CA catalyzes the reversible hydration of CO₂ to HCO₃⁻ and H⁺. At least 16 CA isozymes have so far been described in mammals, the most active ones as catalysts for carbon dioxide hydration being CA II and CA IX²⁻⁵. The first one is primarily found in red blood cells but also in many other secretory tissues of the gastrointestinal tract, kidneys, lungs, eye, central nervous system etc.^{3,6}, whereas the second one is a tumor-associated isoform³⁻⁷.

Carbonic anhydrase inhibitors are a class of pharmaceuticals that suppress the activity of carbonic anhydrase. Their clinical use has been established as antiglaucoma agents, diuretics, antiepileptics in the management of mountain sickness, gastric and duodenal ulcers, neurological disorders, or osteoporosis. The connections between carbonic anhydrase and cancer is known for approximately 15 years, since two tumor-associated membrane carbonic anhydrase isozymes

(CA IX and CA XII) have been identified cloned, and sequenced⁸⁻¹⁰. Many reports have been published on the role of CA IX in tumor physiology. In addition to its role in the control of tumor pH, there are evidences that CA IX can also influence other processes in the cell microenvironment that promote cell proliferation, invasion, and metastasis¹¹⁻¹³.

Other CA isoforms are found in a variety of tissues where they participate in several important biological processes such as acid-base balance, respiration, carbon dioxide and ion transport, bone resorption, ureagenesis, gluconeogenesis, lipogenesis and electrolyte secretion^{4,14}.

As the inhibitors of CA enzymes are very important leads for medicinal chemistry applications¹⁵⁻¹⁷, in this paper we report the inhibitory effects of compounds 1–3, sulfonamidesethoxzolamide (EZA), zonisamide (ZNA) and acetazolamide (AZA) against human CA I, II and VI isoforms. The reason for selection of these compounds is that 2-hydroxyethyl-4-methylthiazolium derivatives were shown to be bioactive molecules¹⁸.

Address for Correspondence: Murat Şentürk, Department of Chemistry, Art and Science Faculty, Ağrı İbrahim Çeçen University, Ağrı, Turkey. Tel: +90-442-2314414. Fax: +90-442-2360948. E-mail: senturkm36@gmail.com; Deniz Ekinci, Department of Agricultural Biotechnology, Faculty of Agriculture, Ondokuz Mayıs University, Samsun, Turkey. Tel: +90-362-3121919. Fax: +90-362-3576034. E-mail: deniz.ekinci@omu.edu.tr

(Received 28 September 2011; revised 28 October 2011; accepted 28 October 2011)

Materials and methods

5-(2-hydroxyethyl)-3,4-dimethylthiazolium iodide, 3-Benzyl-5-(2-hydroxyethyl)-4-methylthiazolium chloride, thiamine (2-[3-[(4-Amino-2-methyl-pyrimidin-5-yl)methyl]-4-methyl-thiazol-5-yl] ethanol), Sepharose 4B, protein assay reagents, 4-nitrophenylacetate were obtained from Sigma-Aldrich Co. (Germany). All other chemicals were analytical grade and obtained from Merck.

Purification of carbonic anhydrase isozymes by affinity chromatography

Purification of hCA I and hCAII were previously described^{14–16}. Fresh non-citrated human whole blood obtained from the Blood Center of the Research Hospital at Atatürk University. The blood samples were centrifuged at 5000 rpm for 15 min and precipitant were removed. The serum was isolated. The pH was adjusted to 8.7 with solid Tris. Sepharose-4B-aniline-sulfanilamide affinity column equilibrated with 25 mM Tris-HCl/0.1 M Na₂SO₄ (pH 8.7). The affinity gel was washed with 25 mM Tris-HCl/22 mM Na₂SO₄ (pH 8.7). The human carbonic anhydrase (hCA VI) isozyme was eluted with 0.25 M H₂NSO₃H/25 mM Na₂HPO₄ (pH=6.7^{19,20}).

CA inhibition assay

The effect of increasing concentrations of compounds 1–3, AZA, ZNA, and EZA on HCA isozyme activity was determined colorimetrically using CO₂-hydration method of Khalifah²¹. Phenol red (at a concentration of 0.2 mM) has been used as indicator, working at the absorbance maximum of 557 nm, with 10 mM Tris-HCl (pH 7.5) as buffer, 0.1 M Na₂SO₄ (for maintaining constant the ionic strength), at 25°C, following the CA-catalyzed CO₂-hydration reaction for a period of 10–80 s (the uncatalyzed reaction needs around 60–100 s in the assay conditions, whereas the catalyzed ones are of around 6–10 s). The CO₂ concentrations ranged from 1.7 to 17 mM for the determination of kinetic parameters. For each inhibitor, tested in the concentration range between 0.01 and 100 µM, at least six traces of the initial 5–10% of the reaction have been used for determining the initial velocity. The uncatalyzed rates were determined in the same manner and subtracted from the total observed rates. Stock solutions of inhibitor (1 mM) were prepared in distilled-deionized water with 10–20% (v/v) dimethyl sulfoxide (which is not inhibitory at these concentrations), and dilutions up to 0.001 µM were done thereafter with distilled-deionized water. Inhibitor and enzyme solutions were preincubated together for 15 min at room temperature prior to assay in order to allow for the formation of the E-I complex. The curve-fitting algorithm allowed us to obtain the IC₅₀ values, working at the lowest concentration of substrate of 1.7 mM, from which K_i values were calculated by using the Cheng-Prusoff equation²². The catalytic activity (in the absence of inhibitors) of these enzymes was calculated from Lineweaver-Burk plots, as reported earlier, and represent the mean from at least

three different determinations. Enzyme concentrations in the assay system were 9.2 nM for hCA I, 7.3 nM for hCA II, and 7.5 nM for hCA VI. Enzymes used here were purified from human blood as described earlier.

Protein determination

Protein quantity was determined spectrophotometrically at 595 nm during the purification steps according to the Bradford method, using bovine serum albumin as a standard²³.

Sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis

SDS polyacrylamide gel electrophoresis was performed after purification of the enzymes. It was carried out in 10% and 3% acrylamide for the running and the stacking gel, respectively, containing 0.1% SDS according to Laemmli procedure²⁴.

Results and discussion

Purification of the two CA isozymes used here was performed with a simple one-step method by a Sepharose-4B-aniline-sulfanilamide affinity column chromatography^{16–19}. hCA I was purified approx. 102-fold with a specific activity of 874.1 EUmg⁻¹ and an overall yield of 62.4%, hCA II was purified approx. 685-fold with a specific activity of 5863.3 EUmg⁻¹ and an overall yield of 58.7% and 63-fold with a specific activity of 248 EUmg⁻¹ and an overall yield of 29.2%. Inhibitory effects of compounds 1–3 and EZA, ZNA and AZA on enzyme activities were tested under *in vitro* conditions; K_i values were calculated from the Cheng-Prusoff equation and are given in Table 1²².

We report here the first study on the inhibitory effects of compounds 1–3 of type on the hydratase activity of hCA I, II and VI. EZA, ZNA and AZA, CAI were used as negative controls in our experiments, and for comparison purposes. The data in Table 1 show the following regarding inhibition of hCA I, II and VI with compounds 1–3 and EZA, ZNA and AZA:

1. Against the slow cytosolic isozyme hCA I, compound **1** behaves as a moderate inhibitor, with a K_i value of 2.27 µM. Compound **2** and thiamine **3** showed better

Table 1. K_i values obtained from regression analysis graphs for hCA I, hCA II and hCA VI in the presence of different inhibitors concentrations (µM).

Compound	hCA I	hCA II	hCA VI
1	2.27	0.784	0.593
2	1.15	0.105	0.128
3	0.38	0.085	0.062
EZA	0.03	0.011	0.050
ZNA	0.06	0.041	0.095
AZA	0.27	0.015	0.015

AZA, acetazolamide; EZA, sulfonamides ethoxzolamide; hCA, human erythrocyte carbonic anhydrase; ZNA, zonisamide.




Figure 1. Structures of tested compounds and some clinical used sulfonamides.

inhibitory activity when compared to the previously mentioned compound **1**, with K_i values of 1.15 and 0.38 μM (Table 1). Thus, the 4-amino-2-methyl-pyrimidin-5-yl moiety improves hCA I inhibitory activity. AZA is also a strong hCA I inhibitor with this assay and K_i of 0.27 μM , whereas thiamine, ZNA and EZA were more powerful inhibitors than AZA (Table 1).

2. A better inhibitory activity has been observed with compounds **2** and **3** for the inhibition of the rapid cytosolic isozyme hCA II (Table 1). Compound **1** showed moderate hCA II inhibitory activity with a K_i value of 0.784 μM (Table 1), whereas thiamine was quite an effective hCA II inhibitor, with a K_i value of 0.085 μM , (Table 1). Similar to hCA I, 4-amino-2-methyl-pyrimidin-5-yl moiety strongly influences hCA II inhibitory activity as well.
3. Compounds **1** and **2** were relatively weak inhibitors for hCA VI, whereas, **3**, EZA and ZNA were moderate inhibitors of the secreted isozyme, with K_i of 0.050–0.095 μM . A better inhibitory activity has been observed with AZA for the inhibition of the secreted isozyme hCA VI (Table 1).

There are many CA inhibitors in the literature^{25–34}, but it is still critically important to carry out further CA inhibition studies to find novel applications for the inhibitors of these widespread enzymes.

Conclusions

Compounds **1–3** used in this study affect the activity of CA isozymes due to the presence of the different functional groups present in their scaffold. 4-amino-2-methyl-pyrimidin-5-yl moiety improves the inhibition capacity of the tested compounds. Our findings here contribute to discovery of novel possible CAIs of interest, in addition to the well-known sulfonamides/sulfamates/sulfamides. In addition, usage of natural molecules as inhibitors of CA or other enzymes is of particular interest for medicinal chemists. Therefore, this study may be very useful for design and biological evaluation of novel agents effective as enzyme inhibitors.

Acknowledgement

The authors thank to Prof. Claudiu T. supuran for several helpful publications on CA inhibitors.

Declaration of interest

The authors report no conflicts of interest.

References

1. Parui R, Gambhir KK, Mehrotra PP. Changes in carbonic anhydrase may be the initial step of altered metabolism in hypertension. *Biochem Int* 1991;23:779–789.
2. Tashian RE, Hewett-Emmet D, Venta PJ. Diversity and evolution in the carbonic anhydrase gene family. In: Botr'e F, Gros G, Storey BT, eds. *Carbonic anhydrase: From biochemistry and genetics to physiology and clinical medicine*. Weinheim: VCH Verlagsgesellschaft; 1991, 151–161.
3. Supuran CT, Scozzafava A. Carbonic anhydrases as targets for medicinal chemistry. *Bioorg Med Chem* 2007;15:4336–4350.
4. Supuran CT. Carbonic anhydrases: Novel therapeutic applications for inhibitors and activators. *Nat Rev Drug Discov* 2008;7:168–181.
5. Bayram E, Senturk M, Kufrevioglu OI, Supuran CT. *In vitro* inhibition of salicylic acid derivatives on human cytosolic carbonic anhydrase isozymes I and II. *Bioorg Med Chem* 2008;16:9101–9105.
6. Ekinci D, Ceyhan SB, Senturk M, Erdem D, Kufrevioglu OI, Supuran CT. Characterization and anions inhibition studies of an α -carbonic anhydrase from the teleost fish *Dicentrarchus labrax*. *Bioorg Med Chem* 2011;19:744–748.
7. Sly WS, Hu PY. Human carbonic anhydrases and carbonic anhydrase deficiencies. *Annu Rev Biochem* 1995;64:375–401.
8. Winum JY, Rami M, Scozzafava A, Montero JL, Supuran C. Carbonic anhydrase IX: A new druggable target for the design of antitumor agents. *Med Res Rev* 2008;28:445–463.
9. Güzel O, Innocenti A, Scozzafava A, Salman A, Parkkila S, Hilvo M et al. Carbonic anhydrase inhibitors: Synthesis and inhibition studies against mammalian isoforms I–XV with a series of 2-(hydrazinocarbonyl)-3-substituted-phenyl-1H-indole-5-sulfonamides. *Bioorg Med Chem* 2008;16:9113–9120.
10. Nuti E, Orlandini E, Nencetti S, Rossello A, Innocenti A, Scozzafava A et al. Carbonic anhydrase and matrix metalloproteinase inhibitors. Inhibition of human tumor-associated isozymes IX and cytosolic isozyme I and II with sulfonylated hydroxamates. *Bioorg Med Chem* 2007;15:2298–2311.
11. Rami M, Innocenti A, Montero JL, Scozzafava A, Winum JY, Supuran CT. Synthesis of rhodamine B-benzenesulfonamide conjugates and their inhibitory activity against human α - and

- bacterial/fungal β -carbonic anhydrases. *Bioorg Med Chem Lett* 2011;21:5210–5213.
12. Rami M, Maresca A, Smaine FZ, Montero JL, Scozzafava A, Winum JY et al. Sulfonamides incorporating boroxazolidone moieties are potent inhibitors of the transmembrane, tumor-associated carbonic anhydrase isoforms IX and XII. *Bioorg Med Chem Lett* 2011;21:2975–2979.
 13. Lou Y, McDonald PC, Oloumi A, Chia S, Ostlund C, Ahmadi A et al. Targeting Tumor Hypoxia: Suppression of breast tumor growth and metastasis by novel carbonic anhydrase IX inhibitors. *Cancer Res* 2011;71:3364–3376.
 14. Ekinci D, Cavdar H, Talaz O, Sentürk M, Supuran CT. NO-releasing esters show carbonic anhydrase inhibitory action against human isoforms I and II. *Bioorg Med Chem* 2010;18:3559–3563.
 15. Senturk M, Gulcin I, Dastan A, Küfrevioglu OI, Supuran CT. Carbonic anhydrase inhibitors: Inhibitor of human erythrocyte I and II isoenzymes with antioxidant phenolic compounds. *Bioorg Med Chem* 2009;17:3207–3211.
 16. Sentürk M, Gülçin I, Beydemir S, Küfrevioglu OI, Supuran CT. *In Vitro* inhibition of human carbonic anhydrase I and II isozymes with natural phenolic compounds. *Chem Biol Drug Des* 2011;77:494–499.
 17. Durdagi S, Sentürk M, Ekinci D, Balaydin HT, Göksu S, Küfrevioglu ÖI, et al. Kinetic and docking studies of phenol-based inhibitors of carbonic anhydrase isoforms I, II, IX and XII evidence a new binding mode within the enzyme active site. *Bioorg Med Chem* 2011;19:1381–1389.
 18. Valverde LF, Cedillo FD, Reyes GC, Ramos ML. Synthesis and antibacterial activity of pregnenolone-vitamin B1 conjugate. *J Mex Chem Soc* 2008;52:130–135.
 19. Ozturk Sarikaya SB, Topal F, Senturk M, Gulcin I, Supuran CT. *In vitro* inhibition of some phenolic compounds on some α -carbonic anhydrase isozymes. *Bioorg Med Chem Lett* 2011;21:4259–4262.
 20. Kivela J, Parkkila S, Waheed A, Parkkila AK, Sly WS, Rajaniemi H. Secretory carbonic anhydrase isoenzyme (VI) in human serum. *Clin Chem* 1997;43:2318–2322.
 21. Khalifah RG. The carbon dioxide hydration activity of carbonic anhydrase. *J Biol Chem* 1971;246:2561–2573.
 22. Cheng Y, Prusoff WH. Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction. *Biochem Pharmacol* 1973;22:3099–3108.
 23. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–254.
 24. Laemmli DK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680–685.
 25. Renzi G, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors: topical sulfonamide antiglaucoma agents incorporating secondary amine moieties. *Bioorg Med Chem Lett* 2000;10:673–676.
 26. Nair SK, Ludwig PA, Christianson DW. Two-site binding of phenol in the active site of human carbonic anhydrase II: Structural implications for substrate association. *J Am Chem Soc* 1994;116:3659–3660.
 27. Ekinci D, Beydemir S, Alim Z. Some drugs inhibit *in vitro* hydratase and esterase activities of human carbonic anhydrase-I and II. *Pharmacol Rep* 2007;59:580–587.
 28. Supuran CT, Mincione F, Scozzafava A, Briganti F, Mincione G, Ilies MA. Carbonic anhydrase inhibitors. Part 52. Metal complexes of heterocyclic sulfonamides: A new class of strong topical intraocular pressure lowering agents with potential use as antiglaucoma drugs. *Eur J Med Chem* 1998;33:247–254.
 29. Innocenti A, Maresca A, Scozzafava A, Supuran CT. Carbonic anhydrase inhibitors: Thioxolone versus sulfonamides for obtaining isozyme-selective inhibitors? *Bioorg Med Chem Lett* 2008;18:3938–3941.
 30. Senturk M, Ekinci D, Goksu S, Supuran CT. Effects of dopaminergic compounds on carbonic anhydrase isozymes I, II, and VI. *J Enzym Inhib Med Ch* 2011. DOI: 10.3109/14756366.2011.591290.
 31. Alp C, Ekinci D, Gültekin MS, Sentürk M, Sahin E, Küfrevioglu OI. A novel and one-pot synthesis of new 1-tosyl pyrrol-2-one derivatives and analysis of carbonic anhydrase inhibitory potencies. *Bioorg Med Chem* 2010;18:4468–4474.
 32. Balaydin HT, Durdagi S, Ekinci D, Senturk M, Goksu S, Menzek A. Inhibition of human carbonic anhydrase isozymes I, II and VI with a series of bisphenol, methoxy and bromophenol compounds. *J Enzym Inhib Med Ch* 2011. DOI:10.3109/14756366.2011.596836.33. Balaydin HT, Soyut H, Ekinci D, Goksu S, Beydemir S, Menzek A, Sahin E. Synthesis and carbonic anhydrase inhibitory properties of novel bromophenols including natural products. *J Enzym Inhib Med Ch* 2011. DOI: 10.3109/14756366.2011.574131.
 34. Ceyhun SB, Sentürk M, Yerlikaya E, Erdogan O, Küfrevioglu OI, Ekinci D. Purification and characterization of carbonic anhydrase from the teleost fish *Dicentrarchus labrax* (European seabass) liver and toxicological effects of metals on enzyme activity. *Environ Toxicol Pharmacol* 2011;32:69–74.