



Journal of Enzyme Inhibition and Medicinal Chemistry

ISSN: 1475-6366 (Print) 1475-6374 (Online) Journal homepage: informahealthcare.com/journals/ienz20

# Stabilization of Aspergillus parasiticus cytosine deaminase by immobilization on calcium alginate beads improved enzyme operational stability

H. Zanna, A. J. Nok, S. Ibrahim & H. M. Inuwa

To cite this article: H. Zanna, A. J. Nok, S. Ibrahim & H. M. Inuwa (2013) Stabilization of Aspergillus parasiticus cytosine deaminase by immobilization on calcium alginate beads improved enzyme operational stability, Journal of Enzyme Inhibition and Medicinal Chemistry, 28:6, 1217-1220, DOI: 10.3109/14756366.2012.724406

To link to this article: https://doi.org/10.3109/14756366.2012.724406



Published online: 03 Oct 2012.

Submit your article to this journal 🖸

Article views: 473



View related articles 🗹



Citing articles: 1 View citing articles

# Stabilization of Aspergillus parasiticus cytosine deaminase by immobilization on calcium alginate beads improved enzyme operational stability

H. Zanna<sup>1</sup>, A. J. Nok<sup>2</sup>, S. Ibrahim<sup>2</sup>, and H. M. Inuwa<sup>2</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Science, University of Maiduguri, Maiduguri, Nigeria and <sup>2</sup>Department of Biochemistry, Faculty of Science, Ahmadu Bello University, Zaria, Nigeria

#### Abstract

Cytosine deaminase (CD) from Aspergillus parasiticus, which has half-life of 1.10h at 37°C, was stabilized by immobilization on calcium alginate beads. The immobilized CD had pH and temperature optimum of 5 and 50°C respectively. The immobilized enzyme also stoichiometrically deaminated Cytosine and 5-fluorocytosine (5-FC) with the apparent  $K_{\rm M}$  values of 0.60 mM and 0.65 mM respectively, displaying activation energy of 10.72 KJ/mol. The immobilization of native CD on calcium alginate beads gave the highest yield of apparent enzymatic activity of 51.60% of the original activity and the enzymatic activity was lost exponentially at 37°C over 12 h with a half-life of 5.80 h. Hence, the operational stability of native CD can be improved by immobilization on calcium alginate beads.

Keywords: Cytosine, 5-fluorocytosine, half-life orcont

**M**C

# Introduction

Cytosine deaminase (CD) can deaminate Cytosine to uracil and 5-fluorocytosine (5-FC) to 5-fluorouracil (5-FU)<sup>1</sup>. The ability of CD to convert nontoxic 5FC into toxic 5FU antimetabolite and its absence in mammalian cells has been exploited in enzyme-prodrug gene therapy and shown to control tumour growth<sup>2</sup>. The Escherichia coli CD although stable, has high  $K_{\rm M}$ for 5-FC while yeast CD has low  $K_{\rm M}$  for 5-FC but is less stable<sup>3</sup>. Cytosine deaminase from Baker's yeast was stabilized through immobilization by various techniques. When the yeast CD was immobilized on epoxy-acrylic resin, the highest yield of apparent enzymatic activity was about 80% of the starting activity, and the half-life became of 10 days, though the enzyme lost activity exponentially at 37°C over a few weeks. With lower apparent enzymatic activity of 20-67% and 21%, and half-lives of 15 and 24 days respectively, formyl-cellulofine and CNBr-activated sepharose 4B gave a good stability. The apparent  $K'_{M}$  for Cytosine and 5FC of the

And Hand Student Junification CD immobilized on epoxy-acrylic resin were 30 and 4 mM respectively which were much larger than those of the free enzyme, about 3 and 1 mM respectively<sup>4</sup>.

Intratumoural conversion of 5FC into 5-fluorouracil (5FU) by locally implanted capsules containing CD followed by systemic administration of 5FC to induce antineoplastic activity at a local site with minimal toxicity was investigated by Nishiyama et al.<sup>5</sup> In vitro studies showed that 5FC combined with CD capsule, made of cellulose tubing, induced significant growthinhibitory effects on the cultured glioma cells. There was a sufficient delivery of 5FC to the tumour cells of the rats with 5FU appearing in the tumour reaching the level of 8.0  $\mu$ g/g at 2 h and staying at more than 1.0  $\mu$ g/g at between 1 and 6 h resulting in significant reduction in the tumour growth<sup>5</sup>.

However, the use of CD from *E. coli* has two problems: first, the, high  $K_{\rm M}$  for 5-FC, second, the difficulty of culturing the bacteria on a large scale to obtain sufficient activity<sup>4</sup>. Cytosine deaminase from A. parasiticus was

Address for Correspondence: H. Zanna, Department of Biochemistry, University of Maiduguri, Maiduguri, Nigeria. Tel.: +2348062210711. E-mail: zannahassan@yahoo.co.uk

<sup>(</sup>Received 23 July 2012; revised 19 August 2012; accepted 19 August 2012)

### 1218 H. Zanna et al.

purified 387.73 folds with an overall yield of 13%, the halflife being 1.10 h at  $37^{\circ}C^{6}$ . This study is aimed at improving the half-life of *A. parasiticus* CD by immobilization on calcium alginate beads and to characterize the immobilized CD.

# **Materials and methods**

## Materials

#### Chemicals

All the chemicals used in this study were of analytical grade and purchased from various sources.

#### Organism

The *A. parasiticus* used in this study was isolated and identified by a mycologist using internet resources.

## Methods

#### Inoculation

Spores of *A. parasiticus* were harvested from 5 day old potato-dextrose agar (PDA) slants by washing with sterile 0.2% tween 80. The spores were used to inoculate the mineral salt medium containing: distilled water 1.0 L, glucose 10 g,  $\text{KH}_2\text{PO}_4$  5.0 g,  $\text{MgSO}_4$ -7H<sub>2</sub>O 1.0 g, NaCl 0.5 g, NaNO<sub>3</sub> 2 g, 0.5 g peptone and pH adjusted to 5.5 with 0.1 M HCl/NaOH as described by refs 7,8.

#### Enzyme assay

Cytosine deaminase activity was assayed as described by Ipata and Cercignani<sup>9</sup>. Enzyme activity was measured by direct spectrophotometric assay from the fall in absorbance at 286 nm following conversion of 4-amino to 4-keto compounds. The following aliquots (0.3 mL of 0.5 mM Tris-HCl buffer pH 7.2, 0.5 mL 0f 0.5 mM cytosine) were dispensed into 1 cm quartz cuvettes. This was followed by the addition of 0.2 mL of enzyme preparation. The solutions were mixed and incubated in a spectrophotometer at 286 nm. Change in absorbance was recorded for 5-10 mins as a function of the activity of CD.

*Cytosine Deaminase Assay Using 5-Fluorocytosine as Substrate* This was done as described by Nishiyama et al.<sup>5</sup>; Mahan et al.<sup>10</sup>

Enzyme activity was measured spectrophotometrically at 255 and 290 nm following conversion of 5-FC to 5-FU.

The following concentrations; 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5 and 4.0 mM 5-fluorocytosine were dispensed into separate tubes. This was followed by the addition of 0.3 mL of 0.5 mM Tris–Hcl buffer, pH7.2, 0.5 mL of 5FC, and 0.2 mL of enzyme. The mixture was shaken thoroughly and incubated at 37°C. One aliquot (50  $\mu$ L) of the reaction mix was taken every minute over a 15 min time period and the reaction terminated in 2.45 mL 0.1 N HCl. Spectrophotometric readings were taken at OD<sub>255</sub> (OD at which 5FU absorbs) and OD<sub>290</sub> (OD at which 5FC absorbs). Then absorbance ratio of OD255/290 was used to measure amount of 5FU formed.

#### Immobilization of CD on calcium alginate

Cytosine deaminase (0.006 mg/mL) was immobilized on calcium alginate as described below. Exactly 1.5 g of Sodium alginate was suspended in 50 mL of distilled water and autoclaved at 121°C for 15 min. After cooling to room temperature 3.5 mL of the enzyme were added and mixed thoroughly and allowed to stand for 10 mins. The enzyme-alginate mixture was carefully pumped through a sterile syringe dropwise into a beaker containing 250 mL of sterile 0.12 M Calcium chloride for the formation of the beads. The beads were kept in solution for 1 h at 4°C to ensure complete precipitation.

#### Kinetics and characterization of immobilized CD

 $V'_{\text{max}}$ ,  $K'_{\text{M}}$ , pH optimum and temperature stability studies of the immobilized enzyme were performed.

## Statiscal analysis

Results are mean  $\pm$  standard deviation for triplicate determination.

# Results

### Native and immobilized CD activities

The specific activities of native and immobilized CD is presented in Figure 1. The native enzyme had a specific activity of 937.30  $\mu$ mol/min/mg. Following immobilization on calcium alginate, the specific activity dropped to 483.33  $\mu$ mol/min/mg representing 51.60% decrease of apparent enzymatic activity.

## pH optimum of immobilized cytosine deaminase

The effect of various pHs on immobilized CD is as shown in Figure 2. The calcium alginate immobilized enzyme had an optimum pH of 5 as opposed to the native enzyme that had a pH optimum of 7.2.



Figure 1. Specific activity of free (0.0102 mg/mL) and immobilized (0.006 mg/mL) *A. parasiticus* cytosine deaminase.

### Effects of temperature on CD catalyzed reactions Temperature optimum of immobilized CD

Figure 3 depicts the effect of temperature on immobilized enzyme. Following the exposure of the immobilized enzyme to various degrees of temperature, maximum apparent enzymatic activity was recorded at the temperature of 50°C.

### Operational stability for immobilized CD

The result of operational stability for the immobilized CD at 37°C is presented in Figure 4. Percentage residual activity following 2 h. of incubation was about 56%. This dropped to about 26% after 5 h. The  $t_{1/2}$  for the immobilized CD was 5.8 h.

#### Apparent activation energy (Ea') for immobilized CD

Figure 5 represents the Arrhenius plot for the determination of Ea' of the immobilized *A. parasiticus* CD. The



Figure 2. Effect of pH on immobilized *A. parasiticus* cytosine deaminase activity.

Ea' of the immobilized CD was calculated to be 10.72 KJ/mole.

# Kinetics of immobilized CD with cytosine and 5-FC as substrates

Table 1 describes the kinetic properties of CD immobilized on calcium alginate beads. Apparent  $K'_{\rm M}$  and  $V'_{\rm max}$  of the immobilized enzyme using cytosine as substrate was 0.60 mM and 3.27 µmol/min with  $K_{\rm cat}$  of 545±5.60 S<sup>-1</sup> while the apparent  $K'_{\rm M}$  and  $V'_{\rm max}$  of the immobilized CD using 5-FC as substrate was 0.65 mM and 0.024 µmol/min with  $K_{\rm cat}$  of 4.00 S<sup>-1</sup>.

#### Discussion

Earlier, we were able to purify and characterize CD from *A. parasiticus*. The purified CD had pH and temperature optimum of 7.2 and 45°C respectively. The native enzyme also had an Ea of 8.4 KJ/mole with  $K_{\rm M}$  values of 0.19 mM and 0.30 mM for cytosine and 5-FC respectively. The operational stability of CD at 37°C was also determined. The percentage residual activity after 2h was 33% with  $t_{1/2}$  of 1.10 h<sup>6</sup>.

Immobilization of the native CD on calcium alginate produced beads of 141.0 mm<sup>3</sup>. The yield of apparent enzymatic activity following immobilization was 54.7%. The aim of the immobilization was to improve the stability of the native enzyme. The result of the operational stability of immobilized CD at 37°C reveals that the immobilized enzyme retained residual activity of 56% after 2h of incubation with  $t_{1/2}$  of 5.8h. The  $t_{1/2}$  of 5.8h for the immobilized CD compared to that of the native enzyme (1.1 h) shows that the native CD can



Figure 3. Effect of temperature on immobilized A. parasiticus CD (0.006 mg/mL/bead) activity showing an optimum temperature of 50°C.



Figure 4. Operational stability of immobilized *A. parasiticus* CD at 37°C. Residual activity of CD was 56% after 2h. Of incubation and it dropped to zero at 12h.



Figure 5. Arrhenius plot for immobilized *A. parasiticus* cytosine deaminase.

Table 1. Kinetics of immobilized *A. parasiticus* CD with cytosine and 5-fluorocytosine as substrates.

| Cytosine        | 5-fluorocytosine                   |
|-----------------|------------------------------------|
| $0.60 \pm 0.02$ | $0.65 \pm 0.03$                    |
| $3.27\pm0.32$   | $0.024 \pm 0.01$                   |
| $545\pm5.60$    | $4.00 \pm 0.50$                    |
|                 | 0.60±0.02<br>3.27±0.32<br>545±5.60 |

Values are mean ± SD for triplicate determination.

be stabilized by immobilization. The buffering effect of the carrier against the conditions of the microenvironment might have contributed to the stabilization of the enzyme activity. The pH optimum of the immobilized CD was 5 as opposed to 7.2 for the native enzyme. The drop in pH could be as a result of the polycationic nature of the carrier. Polycationic supports expel protons from the enzyme's microenvironments and thereby lowering the pH of the bulk phase from which pH measurements are usually done.

The temperature optimum of the immobilized CD was 50°C compared with 40–45°C for the free enzyme. The increase in the optimum temperature following immobilization could be due to the enhanced physical protection provided by the solid support. Increased stability following immobilization is advantageous since this enzyme is intended for use in cancer chemotherapy. The immobilized enzyme had an activation energy of 10.72 KJ/mole, higher than that of the free enzyme (8.4 KJ/mole). This implies that substrate molecules reacting with the immobilized CD must possess higher Ea than when reacting with the free CD. The activation energy following immobilization may be less than, same as or greater than that of the native enzyme.

The apparent Michaelis constant,  $K'_{M'}$  for cytosine and 5FC of the immobilized *A. parasiticus* cytosine deaminase were 0.60 mM and 0.65 mM respectively, which were higher than those of the free enzyme, about 0.19 mM and 0.30 mM respectively. Katsuragi et al.<sup>4</sup>, also reported larger  $K'_{M}$  for cytosine and 5FC (30 and 4 mM respectively) when

cytosine deaminase from Baker's yeast was immobilized on Eupergit C compared with those of the free enzyme (3 and 1 mM respectively). The increase in  $K'_{\rm M}$  of the immobilized enzyme could be due to the electrostatic field of the carrier, diffusional restrictions and interaction with carrier or deactivation due to immobilization. Apparent  $K'_{\rm M}$  decreases when the carrier has opposite charge with that of the substrate and increases when the carrier and the substrate have similar charge.

# Conclusion

Immobilization of the *A. parasiticus* CD on calcium alginate beads improved the operational stability of the enzyme at 37°C by about 8-fold with superior kinetic properties than both yeast and *E. coli* CD. Therefore, cytosine deaminase from *A. parasiticus* is a good candidate for comparison with yeast and *E. coli* CD; and for subsequent deployment in cancer therapy either as capsule implant at the site of the tumour or in suicide gene therapy.

#### **Declaration of interest**

The authors report no declarations of interest.

## References

- 1. Ireton GC, McDermott G, Black ME, Stoddard BL. The structure of *Escherichia coli* cytosine deaminase. J Mol Biol 2002;315:687-697.
- Hamaji Y, Fujimori M, Sasaki T, Matsuhashi H, Matsui-Seki K, Shimatani-Shibata Y et al. Strong enhancement of recombinant cytosine deaminase activity in Bifidobacterium longum for tumortargeting enzyme/prodrug therapy. Biosci Biotechnol Biochem 2007;71:874–883.
- Kievit E, Bershad E, Ng E, Sethna P, Dev I, Lawrence TS et al. Superiority of yeast over bacterial cytosine deaminase for enzyme/ prodrug gene therapy in colon cancer xenografts. Cancer Res 1999;59:1417–1421.
- Katsuragi T, Shibata M, Sakai T, Tonomura K. Stabiliation of cytosine deaminase from Baker's yeast by immobilization. Agric Biol Chem 1989;53:1515–1523.
- Nishiyama T, Kawamura Y, Kawamoto K, Matsumura H, Yamamoto N, Ito T et al. Antineoplastic effects in rats of 5-fluorocytosine in combination with cytosine deaminase capsules. Cancer Res 1985;45:1753–1761.
- Zanna H, Nok AJ, Ibrahim S, Inuwa HM. Purification and characterization of *Aspergillus parasiticus* cytosine deaminase for possible deployment in suicide gene therapy. Adv Biol Chem 2012;2:152–159.
- Haq IU, Ali S, Qadeer MA. Influence of dissolved oxygen concentration on intracellular pH for regulation of *Aspergillus* niger growth rate during citric acid fermentation in a stirred tank bioreactor. Int J Biol Sci 2005;1:34–41.
- Zanna H, Nok AJ, Ibrahim S, Inuwa HM. Screening some *Aspergillus* fungi for cytosine deaminase activity. Int J Biotechnol Biochem 2011;7:397–403.
- 9. Ipata PL, Cercignani G. Cytosine and cytidine deaminase from yeast. Meth Enzymol 1978;51:394–400.
- Mahan SD, Ireton GC, Knoeber C, Stoddard BL, Black ME. Random mutagenesis and selection of *Escherichia coli* cytosine deaminase for cancer gene therapy. Protein Eng Des Sel 2004;17:625–633.