



Efficacy and Safety of Aclidinium Bromide Compared with Placebo and Tiotropium in Patients with Moderate-to-Severe Chronic Obstructive Pulmonary Disease: Results from a 6-week, Randomized, Controlled Phase Iiib Study

Jutta Beier, Anne-Marie Kirsten, Robert Mróz, Rosa Segarra, Ferran Chuecos, Cynthia Caracta & Esther Garcia Gil

To cite this article: Jutta Beier, Anne-Marie Kirsten, Robert Mróz, Rosa Segarra, Ferran Chuecos, Cynthia Caracta & Esther Garcia Gil (2013) Efficacy and Safety of Aclidinium Bromide Compared with Placebo and Tiotropium in Patients with Moderate-to-Severe Chronic Obstructive Pulmonary Disease: Results from a 6-week, Randomized, Controlled Phase Iiib Study, COPD: Journal of Chronic Obstructive Pulmonary Disease, 10:4, 511-522, DOI: [10.3109/15412555.2013.814626](https://doi.org/10.3109/15412555.2013.814626)

To link to this article: <https://doi.org/10.3109/15412555.2013.814626>



© 2013 The Author(s). Published by Taylor & Francis.



Published online: 02 Jul 2013.



Submit your article to this journal [↗](#)



Article views: 5630



View related articles [↗](#)



Citing articles: 5 View citing articles [↗](#)

ORIGINAL RESEARCH

Efficacy and Safety of Aclidinium Bromide Compared with Placebo and Tiotropium in Patients with Moderate-to-Severe Chronic Obstructive Pulmonary Disease: Results from a 6-week, Randomized, Controlled Phase IIIb Study

Jutta Beier,¹ Anne-Marie Kirsten,² Robert Mróz,³ Rosa Segarra,⁴ Ferran Chuecos,⁴ Cynthia Caracta,⁵ and Esther Garcia Gil⁴

¹ insaf Respiratory Research Institute, Wiesbaden, Germany

² Pulmonary Research Institute at Hospital Grosshansdorf, Grosshansdorf, Germany

³ ISPL Centrum Medyczne and Department of Lung Diseases and Tuberculosis, Medical University of Białystok, Białystok, Poland

⁴ Almirall, Barcelona, Spain

⁵ Forest Research Institute, Jersey City, New Jersey, USA

Abstract

Background: This randomized, double-blind, Phase IIIb study evaluated the 24-hour bronchodilatory efficacy of aclidinium bromide versus placebo and tiotropium in patients with moderate-to-severe chronic obstructive pulmonary disease (COPD). **Methods:** Patients received aclidinium 400 µg twice daily (morning and evening), tiotropium 18 µg once daily (morning), or placebo for 6 weeks. The primary endpoint was change from baseline in forced expiratory volume in 1 second area under the curve for the 24-hour period post-morning dose (FEV₁ AUC_{0–24}) at week 6. Secondary and additional endpoints included FEV₁ AUC_{12–24}, COPD symptoms (EXAcerbations of chronic pulmonary disease Tool-Respiratory Symptoms [E-RS] total score and additional symptoms questionnaire), and safety. **Results:** Overall, 414 patients were randomized and treated (FEV₁ 1.63 L [55.8% predicted]). Compared with placebo, FEV₁ AUC_{0–24} and FEV₁ AUC_{12–24} were significantly increased from baseline with aclidinium (Δ = 150 mL and 160 mL, respectively; p < 0.0001) and tiotropium (Δ = 140 mL and 123 mL, respectively; p < 0.0001) at week 6. Significant improvements in E-RS total scores over 6 weeks were numerically greater with aclidinium (p < 0.0001) than tiotropium (p < 0.05) versus placebo. Only aclidinium significantly reduced the severity of early-morning cough, wheeze, shortness of breath, and phlegm, and of nighttime symptoms versus placebo (p < 0.05). Adverse-event (AE) incidence (28%) was similar between treatments. Few anticholinergic AEs (<1.5%) or serious AEs (<3%) occurred in any group. **Conclusions:** Aclidinium provided significant 24-hour bronchodilation versus placebo from day 1 with comparable efficacy to tiotropium after 6 weeks. Improvements in COPD symptoms were consistently numerically greater with aclidinium versus tiotropium. Aclidinium was generally well tolerated.

Keywords: 24-hour, bronchodilation, long-acting muscarinic antagonist, nighttime, symptoms

Clinical trial registration: This trial was registered on clinicaltrials.gov (NCT01462929).

Correspondence to: Dr Jutta Beier, insaf Respiratory Research Institute, Biebricher Allee 34, 65187, Wiesbaden, Germany, phone: +49 611 985 4410, email: j.beier@insaf-wi.de

Introduction

Circadian variation in lung function, driven in part by changes in cholinergic tone, has been documented in patients with chronic obstructive pulmonary disease (COPD) (1–3). Variation in COPD daily symptoms has also been reported, with the most severe symptoms generally experienced in the morning followed by during the nighttime (4, 5). Consequently, the importance of identifying and managing early-morning symptoms is generally accepted. However, unlike in asthma where variation in lung function

and symptoms has been better characterized, the clinical relevance of nighttime symptoms can sometimes be underestimated in COPD and a lack of routine assessment means they can be under-reported (6). As symptoms throughout the day impact on patient quality of life (5), maintaining significant bronchodilation and symptom control over 24 hours should be an important goal of therapy.

Inhaled bronchodilatory therapies, including long-acting β_2 -agonists and long-acting muscarinic antagonists (LAMAs), are central to COPD management (7); however, until recently, tiotropium bromide was the only available agent in the LAMA class (2, 8). Aclidinium bromide is a LAMA that has recently been approved as a maintenance bronchodilator treatment for patients with COPD (9, 10). In a Phase IIa study, twice-daily (BID) treatment with aclidinium 400 μ g was demonstrated to provide significant improvements in 24-hour bronchodilation versus placebo that were generally similar to once-daily (QD) treatment with tiotropium 18 μ g after 2 weeks, although significant differences in favor of aclidinium were observed for the nighttime period (11).

The Phase IIIb study reported in this paper was conducted to confirm the 24-hour bronchodilatory efficacy of aclidinium versus placebo and tiotropium over a longer treatment period (6 weeks) and in a larger population. The effects of treatment on COPD symptoms, inhaler preference, and safety were also evaluated in this Phase IIIb study.

Methods

Study design and treatment

This randomized, double-blind, double-dummy, placebo- and active comparator-controlled, multicenter Phase IIIb study was conducted in the Czech Republic, Germany, Hungary, and Poland between October 2011 and March 2012 (clinicaltrials.gov identifier NCT01462929).

Following screening and a 2- to 3-week run-in period, during which disease stability was assessed, patients were randomized (2:2:1) via an interactive voice-response system and computer-generated schedule to receive aclidinium bromide 400 μ g (metered dose; equivalent to aclidinium 322 μ g delivered dose) BID in the morning and evening via the Genuair[®]/Pressair[™] multidose dry powder inhaler, tiotropium 18 μ g QD in the morning via the HandiHaler[®], or placebo for 6 weeks.

Two Genuair inhalers (pre-loaded with 1 month's supply [60 doses] of either aclidinium or matched placebo) and one HandiHaler (with 60 capsules of tiotropium or matched placebo) were supplied. To maintain blinding, patients were instructed to use both inhalers each morning (9:00 $\pm$ 1 hour) and Genuair only each evening (21:00 $\pm$ 1 hour). Patients and study personnel remained blinded to treatment allocation throughout the study. Training on the correct use of the inhalers was provided at screening and before randomization on day 1.

Relief medication (salbutamol pressurized metered-dose inhaler 100 μ g/puff) was provided for additional symptom control as needed (except $\leq$ 6 hours before each visit). Patients were permitted to continue stable use of oral sustained-release theophylline (use of other methylxanthines was not permitted), inhaled corticosteroids, and oral or parenteral corticosteroids (prednisone $\leq$ 10 mg/day or 20 mg/every other day, or equivalent), except $\leq$ 6 hours before each visit. Oxygen therapy (except $\leq$ 2 hours before each visit) was allowed.

The study was approved by an independent ethics committee at each site and was conducted in accordance with the Declaration of Helsinki, the International Conference on Harmonization and Good Clinical Practice. Patients provided written informed consent before participating in any study procedures.

Patients

Eligible patients were aged $\geq$ 40 years with a clinical diagnosis of stable moderate-to-severe COPD (post-bronchodilator forced expiratory volume in 1 second [FEV₁]/forced vital capacity [FVC] $<$ 70%, and FEV₁ $\geq$ 30% and $<$ 80%) (12) and were either current or former cigarette smokers (smoking history of $\geq$ 10 pack-years). Patients with a history or current diagnosis of asthma or other clinically significant respiratory or cardiovascular conditions were excluded, as were those who had experienced any respiratory tract infection or COPD exacerbation $\leq$ 6 weeks before screening ($\leq$ 3 months if exacerbation resulted in hospitalization), or for whom the use of muscarinic antagonists was contraindicated. Additional relevant exclusion criteria included hypersensitivity to inhaled muscarinic antagonists and inability to use the study inhalers properly.

Patients could be discontinued from the study at any time at their own request or in the event of ineligibility, non-compliance, lack of efficacy, loss to follow-up, safety concerns (including moderate or severe COPD exacerbation), or any other reason at the investigator's discretion.

Study Assessments

Lung function

Lung function was assessed over 24 hours following morning treatment on day 1 and at week 6. FEV₁ and FVC were measured using procedures and spirometers that met European Respiratory Society (ERS) and American Thoracic Society (ATS) standards (13). Three manoeuvres were performed at each time point to provide three measurements that met ATS/ERS acceptability and repeatability criteria. Additional measurements could be made (up to a total of eight tests) if the first three were not acceptable.

The primary endpoint was change from baseline in normalized FEV₁ area under the curve over the 24-hour period post-morning dose (AUC₀₋₂₄) at week 6 (tiotropium administered QD in the morning; aclidinium administered BID in the morning and evening).

Change from baseline in normalized FEV₁ AUC over the nighttime period (AUC₁₂₋₂₄) at week 6 was the secondary endpoint. Changes from baseline in normalized FEV₁ AUC for the 12-hour period post-morning treatment (AUC₀₋₁₂), morning pre-dose (trough) and peak FEV₁ and FVC were additional endpoints.

COPD daily symptoms and relief medication use

Patients completed the 14-item EXacerbations of Chronic pulmonary disease Tool (EXACT) each evening before bedtime (recall period of 'today') via electronic diaries. Responses to 11 of 14 EXACT questions that captured changes in specific respiratory symptoms (grouped into breathlessness, cough and sputum, and chest symptom domains) were used to calculate an EXACT-Respiratory Symptoms (E-RS) (14) total score (range 0–40; a higher score indicates more severe symptoms).

An additional COPD symptoms questionnaire (developed by the study sponsor) was completed by patients each morning via electronic diaries to capture the severity of early-morning and nighttime symptoms (5-point scale: 1 = 'did not experience symptoms'; 5 = 'very severe'), and individual morning symptoms of cough, wheeze, shortness of breath, and phlegm (5-point scale: 0 = 'no symptoms'; 4 = 'very severe symptoms'), as well as limitation of morning activities (5-point scale: 1 = 'not at all'; 5 = 'a very good deal') and frequency of nocturnal awakenings as a result of COPD symptoms. Relief medication use was also recorded daily via electronic diaries.

Inhaler preference and willingness to continue

After 6 weeks, overall inhaler preference and preference based on a number of specific inhaler attributes were assessed. Patients were first asked "which inhaler do you prefer?" (Genuair, HandiHaler, or no preference) then, "which device do you prefer in terms of the following attributes: ease of use, convenience, ease of learning to use, ease of holding, ease of operating, ease of preparation of the dose, and feedback to indicate correct inhalation?" Willingness to continue using each inhaler was also rated on a scale from 0 = 'not willing' to 100 = 'definitely willing'.

Safety

Adverse events (AEs) were recorded throughout the study and assessed for severity and relationship to study treatment by the investigator. Other safety assessments included a physical examination and vital signs measurement at screening and week 6.

Statistical Analysis

A target population of approximately 405 patients was planned to provide a sample size of 385 patients, taking into account a 5% dropout rate. This provided >90% power to detect a 130 mL difference between acclidinium and placebo for the primary and secondary

endpoints, and >80% power to detect a 70 mL difference between acclidinium and tiotropium for the secondary endpoint, with a two-sided significance level of $p < 0.05$. Assumptions about treatment group differences were made based on observations from the previous Phase IIa study that compared the 24-hour bronchodilatory efficacy of acclidinium with placebo and tiotropium (11).

Efficacy analyses were based on the intent-to-treat population, which included all treated patients (≥ 1 dose) who had ≥ 1 baseline and post-baseline FEV₁ value. Endpoints were assessed using an analysis of covariance model with treatment and sex as factors, and age and baseline values as covariates. Between-group least squares mean differences and 95% confidence intervals were calculated for all treatment-group comparisons. The primary and secondary endpoint analyses were conducted in a stepwise manner to control for multiplicity: for FEV₁ AUC₀₋₂₄, the primary comparison was acclidinium versus placebo; for FEV₁ AUC₁₂₋₂₄, the primary comparison was acclidinium versus placebo, and the secondary comparison was acclidinium versus tiotropium. Other comparisons were considered additional.


Inhaler preference was summarized descriptively and the percentage of patients preferring Genuair was assessed using an exact binomial test. Willingness to continue inhaler use was assessed using an analysis of variance. Safety analyses included all treated patients (safety population) and were descriptive in nature.

Results

Patients

Of 485 patients screened, 414 patients from 41 sites (2.7% from 3 sites in the Czech Republic, 49.5% from 20 sites in Germany, 10.6% from 5 sites in Hungary, and 37.2% from 13 sites in Poland) were randomized, treated, and included in the study analyses, and 400 completed the study (Figure 1). The discontinuation rate was slightly higher in the placebo group (5.9%) compared with the acclidinium (2.9%) or tiotropium (2.5%) groups. In the placebo group, reasons for discontinuation were AEs and lack of efficacy, whereas in the acclidinium and tiotropium groups, reasons for discontinuation were AEs and patient's personal request (Figure 1). No patients were lost to follow-up. Treatment compliance, as assessed by the investigator based on information provided in patients' electronic diaries and the Genuair dose indicator or number of tiotropium pierced capsules, was 94.1%, 98.2%, and 96.8% in the placebo, acclidinium, and tiotropium groups, respectively.

Demographics and baseline characteristics were similar across treatment groups, with the exception of higher proportions of male patients in the acclidinium and tiotropium groups than in the placebo group (Table 1). Mean post-bronchodilator percent predicted FEV₁ at screening (adjusted for gender-related differences) was similar in each treatment group but, reflective of the higher proportion of male patients in the active treatment groups, mean



FEV₁ was slightly higher in these groups versus the placebo group. In total, 86.7% of patients had used COPD medications prior to the start of the study. Anticholinergic therapies (LAMA, short-acting muscarinic antagonist [SAMA],

or short-acting β -2 agonist plus SAMA) had been used by 25.6%, 18.8%, and 5.3% of patients, respectively.

Table 1. Demographics and baseline characteristics of the study population (safety population)

	Placebo (N = 85)	Acclidinium 400 µg BID (N = 171)	Tiotropium 18 µg QD (N = 158)
Gender (male), n (%)	48 (56.5)	114 (66.7)	116 (73.4)
Age (years), mean (SD)	62.2 (8.2)	61.8 (8.2)	62.8 (7.9)
Race, n (%)			
White	84 (98.8)	171 (100)	158 (100)
Asian	1 (1.2)	0 (0)	0 (0)
BMI (kg/m ²), mean (SD)	26.7 (4.9)	27.5 (4.9)	27.6 (4.8)
Current smoker, n (%)	47 (53.3)	93 (54.4)	84 (53.2)
Smoking consumption (pack-years), mean (SD)	39.6 (15.4)	41.5 (22.4)	45.0 (21.8)
COPD duration (years), mean (SD)	9.6 (6.7)	8.8 (5.9)	8.2 (6.0)
Post-bronchodilator FEV ₁ (L)			
Mean (SD)	1.57 (0.52)	1.61 (0.50)	1.67 (0.54)
% predicted, mean (SD)	55.5 (11.8)	55.8 (13.3)	56.0 (13.2)
COPD severity, ^{a,b} n (%)			
Moderate	58 (68.2)	108 (63.2)	104 (66.2)
Severe	27 (31.8)	63 (36.8)	53 (33.8)

^aGOLD Stage II (moderate): FEV₁/FVC < 0.70, and post-bronchodilator FEV₁ ≥ 50% and < 80% predicted; GOLD Stage III (severe): FEV₁/FVC < 0.70, and post-bronchodilator FEV₁ ≥ 30% and < 50% predicted.

^bCOPD severity was missing for one patient in the tiotropium treatment group at screening (pre- and post-salbutamol values determined to be unacceptable following review).

BID, twice daily; BMI, body mass index; COPD, chronic obstructive pulmonary disease; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; GOLD, Global initiative for chronic Obstructive Lung Disease; QD, once daily; SD, standard deviation.

Lung function

On day 1 of treatment, both acclidinium and tiotropium improved 24-hour, nighttime, and day time bronchodilation, as demonstrated by statistically significant increases in FEV_1 AUC_{0-24} , FEV_1 AUC_{12-24}

and FEV_1 AUC_{0-12} from baseline versus placebo ($p < 0.001$; Figure 2A). Improvements in FEV_1 AUC_{0-24} and FEV_1 AUC_{12-24} were significantly greater with acclidinium versus tiotropium ($p < 0.05$ and $p < 0.01$, respectively) on day 1.




Figure 2. Change from baseline in FEV_1 AUC_{0-24} , FEV_1 AUC_{12-24} , and FEV_1 AUC_{0-12} compared with placebo (A) on day 1 and (B) at week 6 (ITT population). Data reported as LS mean (95% CI) differences from placebo (ANCOVA). † $p < 0.05$; § $p < 0.001$ for acclidinium or tiotropium versus placebo. ‡ $p < 0.001$ for acclidinium versus tiotropium. ANCOVA, analysis of covariance; AUC, area under the curve; AUC_{0-24} , AUC over the 24-hour period post-morning treatment; AUC_{12-24} , AUC over the nighttime period; AUC_{0-12} , AUC for the 12-hour period post-morning treatment; BID, twice daily; CI, confidence interval; FEV_1 , forced expiratory volume in 1 second; ITT, intent-to-treat; LS, least squares; QD, once daily.

In the primary endpoint analysis, FEV_1 AUC_{0-24} was significantly improved with acclidinium compared with placebo at week 6 ($p < 0.0001$; Figure 2B). Compared with placebo, tiotropium also significantly increased FEV_1 AUC_{0-24} from baseline to week 6 ($p < 0.0001$; Figure 2B); the effects of acclidinium and tiotropium over 6 weeks were similar.

Over 6 weeks, FEV_1 AUC_{12-24} and FEV_1 AUC_{0-12} were also significantly increased from baseline with both acclidinium and tiotropium versus placebo ($p < 0.0001$) (Figure 2B). Although the improvement in FEV_1 AUC_{12-24} was numerically greater with acclidinium versus tiotropium, and the improvement in FEV_1 AUC_{0-12} was numerically greater with tiotropium versus acclidinium, there were no significant differences between active treatments ($p = 0.12$ and $p = 0.48$, respectively).

Both acclidinium and tiotropium produced significant increases from baseline in FEV_1 versus placebo at each observation time point from 15 minutes to 24 hours post-dose on day 1 and at week 6 (Figure 3). Both acclidinium and tiotropium also significantly improved morning pre-dose (trough) and peak FEV_1 and FVC compared with placebo on day 1 and at week 6 (Table 2). Improvements were numerically greater with acclidinium versus tiotropium, with significant differences in favor of acclidinium for trough FEV_1 and FVC on day 1.

COPD Symptoms and Relief Medication Use

Over 6 weeks, E-RS total scores were significantly reduced from baseline with both acclidinium ($p < 0.0001$) and tiotropium ($p < 0.05$) versus placebo (Figure 4). Improvements in individual domain scores were numerically greater with acclidinium than tiotropium, and the improvement for cough and sputum was significant versus placebo for acclidinium only ($p < 0.05$). Comparisons of acclidinium versus tiotropium yielded no significant differences.

The severity of early-morning symptoms, overall, was significantly reduced over 6 weeks with acclidinium ($p < 0.001$) and tiotropium ($p < 0.05$) versus placebo (Figure 5A); the difference between active treatments was not statistically significant. Only acclidinium produced significant improvements in individual early-morning symptoms of phlegm, shortness of breath, wheeze, and cough compared with placebo at week 6. Night-time symptom severity was significantly reduced from baseline over 6 weeks with acclidinium versus placebo but not with tiotropium versus placebo (Figure 5B); the difference between active treatments was not statistically significant. There were no significant changes from baseline in number of nocturnal awakenings in any treatment group. Limitation of activity caused by COPD symptoms was also significantly reduced from baseline over 6 weeks with acclidinium versus placebo but not with tiotropium versus placebo (Figure 5). This improvement in limitation of activity was significantly greater for acclidinium versus tiotropium ($p < 0.05$).

Over 6 weeks, there was a significant increase in relief medication-free days with acclidinium and tiotropium versus placebo (difference of 9.6% and 8.9%, respectively; $p < 0.05$).

Inhaler preference

When asked "which device do you prefer?" at week 6, significantly more patients overall preferred Genuair to HandiHaler (80.1% vs 10.7%; $p < 0.0001$). The option of 'no preference' was chosen by 9.2% of patients. Additionally, >75% of patients preferred Genuair to HandiHaler for each of the individual inhaler attributes assessed (Table 3). Inhaler preference appeared to be independent of whether active medication or placebo was administered via the inhalers: when the acclidinium, tiotropium, and placebo groups were analyzed separately, $\geq 79\%$ of patients preferred Genuair compared with $\leq 13\%$ who preferred HandiHaler in each group. Patients were more willing to continue using Genuair than HandiHaler, as indicated by a significant difference in mean ratings at week 6 (88.8 vs 45.4; $p < 0.0001$).

Safety

AE incidence was similar in the placebo (25.9%), acclidinium (27.5%), and tiotropium (29.7%) groups. Nasopharyngitis and headache were most common, each reported by 5.1% of patients overall. Nasopharyngitis occurred more frequently with acclidinium and tiotropium versus placebo (5.8% and 5.7% vs 2.4%, respectively); headache occurred more frequently with acclidinium than with either placebo or tiotropium (7.0% vs 3.5% and 3.8%, respectively). Other common AEs ($\geq 2\%$ of patients overall) were COPD exacerbation (2.4%) and cough (2.2%). The majority of AEs were mild or moderate in intensity. There were few serious AEs (1.7% overall) and no deaths. In total, 8 patients (1.9%) discontinued due to AEs, with COPD exacerbation being the most common cause ($n = 2$ in each treatment group). No AEs resulting in discontinuation were considered to be treatment-related.

With the exception of dry mouth, which was reported by three patients (0.7%) in total (acclidinium $n = 1$; tiotropium $n = 2$), no other treatment-related AEs occurred in >1 patient overall. The incidence of potentially anticholinergic AEs was similarly low across treatment groups ($< 1.5\%$ in any group). Only dry mouth, pharyngitis (placebo $n = 1$; acclidinium $n = 1$; tiotropium $n = 2$), and constipation (tiotropium $n = 2$) occurred in >1 patient in any group. No potentially anticholinergic AEs were serious or led to discontinuation. There were no clinically significant physical examination or vital signs findings.

Discussion

This Phase IIIb study was designed to evaluate the 24-hour bronchodilatory efficacy of acclidinium 400 μ g BID in patients with moderate-to-severe COPD. Findings from our primary endpoint analysis of change from baseline in FEV_1 AUC_{0-24} after 6 weeks of treatment




Figure 3. Change from baseline in FEV₁ over 24 hours (A) on day 1 and (B) at week 6. Data reported as LS mean change from baseline (ANCOVA). $p < 0.01$ for acclidinium and tiotropium versus placebo at each time point. * $p < 0.05$; # $p < 0.01$; $^{\dagger}p < 0.001$; $^{\ddagger}p < 0.0001$ for acclidinium versus tiotropium. ANCOVA, analysis of covariance; BID, twice daily; FEV₁, forced expiratory volume in 1 second; LS, least squares; QD, once daily.

demonstrated a statistically significant improvement with acclidinium versus placebo, and changes from baseline in trough FEV₁ with acclidinium exceeded the proposed minimally important difference for this parameter (100–140 mL) (15, 16). Furthermore, acclidinium significantly improved day time and nighttime bronchodilation over placebo. These findings are consistent with an earlier 15-day Phase IIa cross-over study ($n = 30$) that

also evaluated 24-hour bronchodilation with acclidinium versus placebo and tiotropium, although a larger difference between acclidinium and placebo for the change from baseline in FEV₁ AUC_{0–24} (232 mL) was reported previously, potentially as a result of differences in trial design and duration (11).

Also consistent with previous findings are the numerically greater improvements in nighttime bronchodilation

Table 2. Change from baseline (difference from placebo) in additional lung-function variables (ITT population)

	Acclidinium 400 µg BID (N = 171)	Tiotropium 18 µg QD (N = 158)	Δ (aclidinium vs tiotropium)
<i>Morning pre-dose (trough) FEV₁ (mL)</i>			
Day 1	141 [§]	93 [‡]	48 [¶]
Week 6	141 [§]	102 [‡]	38
<i>Peak FEV₁ (mL)</i>			
Day 1	154 [§]	139 [§]	14
Week 6	180 [§]	172 [§]	8
<i>Morning pre-dose (trough) FVC (mL)</i>			
Day 1	212 [§]	127 [†]	84 [¶]
Week 6	223 [§]	144 [†]	79
<i>Peak FVC (mL)</i>			
Day 1	201 [§]	146 [§]	54
Week 6	212 [§]	170 [†]	42

Data reported as LS mean differences from placebo (ANCOVA).

[†]p < 0.01; [‡]p < 0.001; [§]p < 0.0001 for aclidinium or tiotropium versus placebo.

[¶]p < 0.05 for aclidinium versus tiotropium.

ANCOVA, analysis of covariance; BID, twice daily; FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; ITT, intent-to-treat; LS, least squares; QD, once daily.

achieved with aclidinium versus tiotropium in this study, with statistically significant improvements favoring aclidinium on day 1 of treatment. These day 1 findings may be explained by differences in the pharmacokinetics of aclidinium and tiotropium, whereby aclidinium reaches steady state more quickly than tiotropium (17, 18). After 6 weeks of treatment, the 24-hour bronchodilatory

Table 3. Patient preference for each inhaler based on specific inhaler attributes at week 6, n (%)

Attribute	No. of patients (%)		
	Genuair	HandiHaler	No preference
Ease of use	357 (86.7)	37 (9.0)	18 (4.4)
Convenience	360 (87.4)	32 (7.8)	20 (4.9)
Ease of learning to use	326 (79.1)	33 (8.0)	53 (12.9)
Ease of holding	329 (79.9)	35 (8.5)	48 (11.7)
Ease of operating	334 (81.1)	38 (9.2)	40 (9.7)
Ease of dose preparation	356 (86.4)	32 (7.8)	24 (5.8)
Feedback to indicate correct inhalation	314 (76.2)	39 (9.5)	59 (14.3)

For all attributes, p < 0.0001 for Genuair versus HandiHaler (exact binomial test).

efficacy of aclidinium and tiotropium was considered to be comparable.

Many patients with COPD experience early-morning or nighttime peaks in symptom severity (5). Although as many as 80%–90% of patients experience nighttime symptoms (19) and sleep disturbance (6), there is a lack of therapeutic options for their management. To date, a small number of studies have evaluated the efficacy of bronchodilatory therapies to improve nighttime lung function and/or sleep quality with mixed findings (2, 20–23). For example, a 6-week study in patients with stable COPD found that evening dosing of tiotropium did not result in improved nighttime bronchodilation compared with morning dosing (2).




Figure 4. Change from baseline in E-RS total and domain scores over 6 weeks. Data reported as LS mean (95% CI) difference from placebo (ANCOVA). *p < 0.05; [†]p < 0.01; [§]p < 0.0001 for aclidinium or tiotropium versus placebo. ANCOVA, analysis of covariance; BID, twice daily; CI, confidence interval; E-RS, EXAcacerbations of Chronic pulmonary disease Tool (EXACT)-Respiratory Symptoms; LS, least squares; QD, once daily.




Figure 5. Difference from placebo in change from baseline in (A) severity of early-morning symptoms, (B) severity of nighttime symptoms and number of nocturnal awakenings due to COPD symptoms, and (C) limitation of activity caused by COPD symptoms (COPD additional symptoms questionnaire) over 6 weeks (ITT population). Data reported as LS mean differences from placebo (ANCOVA). * $p < 0.05$; [†] $p < 0.01$; [‡] $p < 0.001$ for acclidinium or tiotropium versus placebo. [†] $p < 0.05$ for acclidinium versus tiotropium. Severity of overall early-morning and nighttime symptoms rated on a 5-point scale from 1 = 'did not experience any symptoms' to 5 = 'very severe'; individual morning symptoms rated on a 5-point scale from 0 = 'no symptoms' to 4 = 'very severe'; limitation of activity rated on a 5-point scale from 1 = 'not at all' to 5 = 'a very good deal'. ANCOVA, analysis of covariance; BID, twice daily; COPD, chronic obstructive pulmonary disease; ITT, intent-to-treat; LS, least squares; QD, once daily.

Additionally, reduced daily variation in lung function and improved nighttime oxygen saturation achieved with evening administration of tiotropium in a separate 4-week study in patients with severe COPD did not translate into improved sleep quality (21). Conversely, four-times-daily treatment with ipratropium (a SAMA) has been shown to improve oxygen saturation and sleep quality over 4 weeks in patients with moderate-to-severe COPD (20).

The impact of treatment with acclidinium on sleep quality has not been evaluated in a sleep laboratory setting; however, it has been demonstrated to improve health status and early-morning and nighttime symptoms, and to reduce the frequency of nocturnal awakenings in Phase III studies (ATTAIN and ACCORD COPD I) (24–27). In the present study, changes in COPD daily symptoms following BID dosing of acclidinium and QD dosing of tiotropium compared with placebo were assessed using the E-RS tool (14, 28), and an additional symptoms questionnaire (currently undergoing validation), that was developed by the sponsor in the absence of a validated tool to capture the severity and impact of early-morning and nighttime symptoms.

Our results suggest that acclidinium provides statistically significant improvements in early-morning and nighttime symptoms compared with placebo that were consistently numerically greater than those observed with tiotropium. Improvements in nighttime symptom severity were significantly different versus placebo for acclidinium only, which could suggest that the numerical advantage of acclidinium over tiotropium for greater nighttime bronchodilation may translate into statistically significant changes in patient-reported outcomes. Furthermore, only acclidinium significantly reduced the limitation of activity caused by COPD symptoms compared with placebo. This is the first report of this therapeutic effect with a LAMA but should, along with the other COPD additional symptoms findings, be interpreted with some caution given the generally mild symptoms that patients reported at baseline, the small magnitude of the reductions observed, and the as yet unvalidated state of this tool.

The trend towards greater symptomatic improvement with acclidinium over tiotropium observed in this study may be related to differences in dosing frequency. However, while a second evening dose of acclidinium may be beneficial in terms of improving nighttime and early-morning symptoms under clinical trial conditions, the potential disadvantages of BID versus QD dosing should also be considered. Findings from an observational study have suggested that treatment adherence among patients with COPD declines with increasing dosing frequency (29), but this appears to be a greater concern for three- and four-times-daily regimens, and there is a lack of evidence to support greater adherence to QD versus BID treatment in practice.

Furthermore, poor adherence to prescribed treatment in COPD is multifactorial: it can also be influenced

by perceived efficacy of treatment, side effects, number of concomitant medications, patient understanding, inhaler satisfaction or preference, and a number of other issues (30–32). Due to the double-blind, double-dummy design of this study, adherence to tiotropium QD compared with acclidinium BID could not be assessed but may be of interest for future work.

Although not powered for this purpose, patients' preference for Genuair compared with HandiHaler was assessed as an additional endpoint in this study and our findings were consistent with those of a previous study that reported greater preference for, greater satisfaction with, and fewer errors with, Genuair versus HandiHaler after 2 weeks' daily practice with placebo-containing devices (33).

However, in contrast with the previous study, patients inhaled both acclidinium and placebo through Genuair, and both tiotropium and placebo via HandiHaler in the present study. Patients were not asked to consider perceived efficacy when indicating their preferred device and preference for Genuair was maintained regardless of whether it was used to deliver placebo or acclidinium.

Finally, acclidinium was generally well tolerated over 6 weeks in this study, with a similar safety profile to tiotropium. Our findings are consistent with previous Phase III studies in terms of similar incidences of AEs and serious AEs in patients treated with acclidinium or placebo (24, 25). The potential for anticholinergic AEs, such as dry mouth, constipation, urinary retention, and cardiovascular events, is a risk associated with LAMAs.

Acclidinium, however, has been shown to undergo more rapid hydrolysis than tiotropium in human plasma (34, 35) and could, therefore, be considered to have the potential for fewer systemic side effects than tiotropium. Tiotropium has been associated with an increased risk of all-cause and cardiovascular-related mortality when administered via soft mist inhaler (36–38) but not when administered via HandiHaler (39).

In this 6-week study, the incidence of dry mouth, constipation, and other anticholinergic AEs (<1.5% for any event) was somewhat lower than has been observed in a pooled analysis of tiotropium safety data from 26 Phase III and IV studies (39), possibly as a result of the short trial duration and patients' previous exposure to tiotropium. The incidence of anticholinergic AEs was similarly low in the acclidinium treatment group, consistent with findings from other Phase III studies (24, 25).

Conclusions

These findings suggest that in patients with moderate-to-severe COPD, acclidinium 400 µg BID provides significant 24-hour bronchodilation compared with placebo from day 1 and over 6 weeks of treatment. At 6 weeks, the bronchodilatory effects of acclidinium and tiotropium are generally comparable. Acclidinium provides consistently numerically greater improvements in COPD symptoms, including early-morning and

nighttime symptoms, than tiotropium and is well tolerated, with a similar safety profile. Patients in this study preferred Genuair over HandiHaler for the administration of their inhaled medications.

Acknowledgment

This study was supported by Almirall, S.A., Barcelona, Spain, and Forest Laboratories, Inc., New York, NY, USA. Editorial support, funded by Almirall, S.A., was provided by Lynsey Stevenson of Complete Medical Communications.

Declaration of Interest Statement

Jutta Beier has received consulting fees, speaker's fees, and travel expenses from Almirall. Anne-Marie Kirsten has received research grants from Almirall. Rosa Segarra, Ferran Chuecos, and Esther Garcia Gill are employees of Almirall and have stock/stock options in Almirall. Cynthia Caracta is an employee of Forest Research Institute. Robert Mróz declares that he has no conflicts of interest to disclose.

Genuair[®] is the registered trademark of Almirall S.A., Barcelona, Spain for use within the European Union, Iceland, and Norway, and within the United States as Pressair[™].

The authors alone are responsible for the content and [AQ3] writing of the paper.

References

- Postma DS, Koëter GH, vd Mark TW, Reig RP, Sluiter HJ. The effects of oral slow-release terbutaline on the circadian variation in spirometry and arterial blood gas levels in patients with chronic airflow obstruction. *Chest* 1985; 87:653–657.
- Calverley PM, Lee A, Towse L, van Noord J, Witek TJ, Kelsen S. Effect of tiotropium bromide on circadian variation in airflow limitation in chronic obstructive pulmonary disease. *Thorax* 2003; 58:855–860.
- van Noord JA, Aumann JL, Janssens E, Verhaert J, Smeets JJ, Mueller A, Cornelissen PJ. Effects of tiotropium with and without formoterol on airflow obstruction and resting hyperinflation in patients with COPD. *Chest* 2006; 129:509–517.
- Kessler R, Partridge MR, Miravittles M, Cazzola M, Vogelmeier C, Leynaud D, Ostinelli J. Symptom variability in patients with severe COPD: a pan-European cross-sectional study. *Eur Respir J* 2011; 37:264–272.
- Partridge MR, Karlsson N, Small IR. Patient insight into the impact of chronic obstructive pulmonary disease in the morning: an internet survey. *Curr Med Res Opin* 2009; 25:2043–2048.
- Agusti A, Hedner J, Marin JM, Barbé F, Cazzola M, Rennard S. Nighttime symptoms: a forgotten dimension of COPD. *Eur Respir Rev* 2011; 20:183–194.
- <http://www.goldcopd.org/guidelines-global-strategy-for-diagnosis-management.html> (accessed February 2013).
- Donohue JF, van Noord JA, Bateman ED, Langley SJ, Lee A, Witek TJ, Jr., Kesten S, Towse L. A 6-month, placebo-controlled study comparing lung function and health status changes in COPD patients treated with tiotropium or salmeterol. *Chest* 2002; 122:47–55.
- http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002211/human_med_001571.jsp&mid=WC0b01ac058001d124 (accessed March 2013).
- <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=SearchDrugDetails> (accessed March 2013).
- Fuhr R, Magnussen H, Sarem K, Llovera AR, Kirsten AM, Falqués M, Caracta CF, Garcia Gil E. Efficacy of acclidinium bromide 400 µg twice daily compared with placebo and tiotropium in patients with moderate to severe COPD. *Chest* 2012; 141:745–752.
- Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. www.goldcopd.com 2010.
- Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. Standardisation of spirometry. *Eur Respir J* 2005; 26:319–338.
- Murray L, Houle C, Stolar M, Hsieh R, Jones PW, Monz BU, Ramachandran S, Goldman M, Sethi S, Leidy NK. Quantifying the severity of respiratory symptoms of chronic obstructive pulmonary disease (COPD): performance properties of the EXacerbations of Chronic pulmonary disease Tool - Respiratory Symptoms (E-RS) in 3 randomized controlled trials. *Am J Resp Crit Care Med* 2012; 185:A1515.
- Cazzola M, MacNee W, Martinez FJ, Rabe KF, Franciosi LG, Barnes PJ, Brusasco V, Burge PS, Calverley PM, Celli BR, Jones PW, Mahler DA, Make B, Miravittles M, Page CP, Palange P, Parr D, Pistolesi M, Rennard SI, Rutten-van Mölken MP, Stockley R, Sullivan SD, Wedzicha JA, Wouters EF. Outcomes for COPD pharmacological trials: from lung function to biomarkers. *Eur Respir J* 2008; 31:416–469.
- Donohue JF. Minimal clinically important differences in COPD lung function. *COPD* 2005; 2:111–124.
- <http://www.spiriva.com/> (accessed March 2013).
- Lasseter KC, Dilzer S, Jansat JM, Garcia Gil E, Caracta C, Ortiz S. Safety and pharmacokinetics of multiple doses of acclidinium bromide administered twice daily in healthy volunteers. *Pulm Pharmacol Ther* 2012; 25:193–199.
- Mocarski M, Caracta C, Tourkodimitris S, Park G, Garcia Gil E, Setyawan J. Nighttime symptoms of COPD in a clinical trial population: prevalence and impact. *Am J Resp Crit Care Med* 2011; 183, abs A1495.
- Martin RJ, Bartelson BL, Smith P, Hudgel DW, Lewis D, Pohl G, Koker P, Souhrada JF. Effect of ipratropium bromide treatment on oxygen saturation and sleep quality in COPD. *Chest* 1999; 115:1338–1345.
- McNicholas WT, Calverley PM, Lee A, Edwards JC. Long-acting inhaled anticholinergic therapy improves sleeping oxygen saturation in COPD. *Eur Respir J* 2004; 23:825–831.
- Mulloy E, McNicholas WT. Theophylline improves gas exchange during rest, exercise, and sleep in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1993; 148:1030–1036.
- Ryan S, Doherty LS, Rock C, Nolan GM, McNicholas WT. Effects of salmeterol on sleeping oxygen saturation in chronic obstructive pulmonary disease. *Respiration* 2010; 79:475–481.
- Jones PW, Singh D, Bateman ED, Agusti A, Lamarca R, de Miquel G, Segarra R, Caracta C, Garcia Gil E. Efficacy and safety of twice-daily acclidinium bromide in COPD patients: The ATTAIN study. *Eur Respir J* 2012; 40:830–836.
- Kerwin EM, D'Urzo AD, Gelb AF, Lakkis H, Garcia Gil E, Caracta CF, ACCORD I study investigators. Efficacy and safety of a 12-week treatment with twice-daily acclidinium bromide in COPD patients (ACCORD COPD I). *COPD* 2012; 9:90–101.
- Agusti A, Jones PW, Bateman ED, Singh D, Lamarca R, de Miquel G, Caracta C, Garcia Gil E. Improvement in symptoms

- and rescue medication use with aclidinium bromide in patients with chronic obstructive pulmonary disease: results from ATTAIn. *Eur Respir J* 2011; 38:149.
27. Kerwin E, Rennard S, Gelb AE, Rekeda L, Garcia Gil E, Caracta C. ACCORD COPD I: improvements in nighttime symptoms and rescue medication use in COPD with twice-daily aclidinium bromide. Poster presented at the European Respiratory Society Annual Congress, Amsterdam, The Netherlands, 24–28 September, 2011.
 28. <http://www.exactproinitiative.com/default.php> (accessed April 2013).
 29. Toy EL, Beaulieu NU, McHale JM, Welland TR, Plauschinat CA, Swensen A, Duh MS. Treatment of COPD: relationships between daily dosing frequency, adherence, resource use, and costs. *Respir Med* 2011; 105:435–441.
 30. Chrystyn H, Small M, Milligan G, Higgins V, Garcia Gil E, Karakurum C. Impact of patient satisfaction with their maintenance inhaler on treatment compliance and health outcomes in chronic obstructive pulmonary disease: an analysis of real-world clinical practice in Europe. Abstract presented at the British Thoracic Society Winter Meeting, London, UK, 5–7 December 2012.
 31. Lareau SC, Yawn BP. Improving adherence with inhaler therapy in COPD. *Int J Chron Obstruct Pulmon Dis* 2010; 5:401–406.
 32. Restrepo RD, Alvarez MT, Wittnebel LD, Sorenson H, Wettstein R, Vines DL, Sikkema-Ortiz J, Gardner DD, Wilkins RL. Medication adherence issues in patients treated for COPD. *Int J Chron Obstruct Pulmon Dis* 2008; 3:371–384.
 33. van der Palen J, Ginko T, Kroker A, van der Valk P, Goossens M, Padulles L, Seoane B, Rekeda L, Garcia-Gil E. Preference, satisfaction and critical errors with Genuair® and HandiHaler® in patients with COPD. *Eur Respir J* 2012; 40 (Suppl 56): 389s.
 34. Gavalda A, Sentellas S, Albertí J, Miralpeix M, Antón F, Salvà M, Beleta J. Rapid plasma hydrolysis of aclidinium bromide, a novel long-acting anticholinergic drug. Poster presented at the European Respiratory Society Annual Congress, Berlin, Germany, 4–8 October, 2008.
 35. Sentellas S, Ramos I, Albertí J, Salvà M, Antón F, Miralpeix M, Beleta J, Gavalda A. Aclidinium bromide, a new, long-acting, inhaled muscarinic antagonist: *in vitro* plasma inactivation and pharmacological activity of its main metabolites. *Eur J Pharm Sci* 2010; 39:283–290.
 36. Dong YH, Lin HH, Shau WY, Wu YC, Chang CH, Lai MS. Comparative safety of inhaled medications in patients with chronic obstructive pulmonary disease: systematic review and mixed treatment comparison meta-analysis of randomised controlled trials. *Thorax* 2013; 68:48–56.
 37. Karner C, Chong J, Poole P. Tiotropium versus placebo for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012; 7:CD009285.
 38. Singh S, Loke YK, Enright PL, Furberg CD. Mortality associated with tiotropium mist inhaler in patients with chronic obstructive pulmonary disease: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2011; 342:d3215.
 39. Kesten S, Jara M, Wentworth C, Lanes S. Pooled clinical trial analysis of tiotropium safety. *Chest* 2006; 130:1695–1703.